

Linear Factors That Repeat

$$\frac{5x^2 + 20x + 6}{x^3 + 2x^2 + x}$$

$$\frac{5x^2 + 20x + 6}{x(x^2 + 2x + 1)}$$

$$\frac{5x^2 + 20x + 6}{(x)(x+1)(x+1)}$$

$$\frac{A}{x} + \frac{B}{x+1} + \frac{C}{(x+1)^2}$$

note: square

repeat

$$[CD: (x)(x+1)^2]$$

note ↑
special

$$A(x+1)^2 + B(x)(x+1) + C(x)$$

$$A(x^2 + 2x + 1) + B(x^2 + x) + Cx$$

$$Ax^2 + 2Ax + A + Bx^2 + Bx + Cx$$

$$Ax^2 + Bx^2 + 2Ax + Bx + Cx + A$$

$$x^2(A+B) + x(2A+B+C) + A$$

given

$$\begin{aligned} A+B &= 5 \\ 2A+B+C &= 20 \\ A &= 6 \end{aligned} \quad \begin{aligned} \rightarrow (6) + B &= 5 \\ (B &= -1) \end{aligned} \quad \begin{aligned} 2(6) + (-1) + C &= 20 \\ 11 + C &= 20 \\ C &= 9 \end{aligned}$$

write

$$\frac{6}{x} - \frac{1}{x+1} + \frac{9}{(x+1)^2}$$

$$\frac{3x^2 + 4x + 5}{(x+2)(x-1)^2}$$

MLC

$$\rightarrow \frac{A}{x+2} + \frac{B}{x-1} + \frac{C}{(x-1)^2}$$

$$A(x-1)^2 + B(x+2)(x-1) + C(x+2)$$

$$A(x^2 - 2x + 1) + B(x^2 + x - 2) + C(x+2)$$

$$Ax^2 - 2Ax + A + Bx^2 + Bx - 2B + Cx + 2C$$

$$Ax^2 + Bx^2 - 2Ax + Bx + Cx + A - 2B + 2C$$

$$x^2(A+B) + x(-2A+B+C) + A - 2B + 2C$$

$$\begin{array}{rcl}
 A+B=3 & \cdot 2 & 2A+2B=6 \\
 -2A+B+C=4 & & \\
 + A-2B+2C=5 & \cdot 2 & 2A-4B+4C=10 \\
 & & \boxed{-3B+5C=14}
 \end{array}$$

rearrange

$$C = 2A - B + 4$$

$$\begin{aligned}
 A - 2B + 2(2A - B + 4) &= 5 \\
 A - 2B + 4A - 2B + 8 &= 5
 \end{aligned}$$

$$5A - 4B = -3$$

$$-5A - 5B = -15$$

$$-9B = -18$$

$$\boxed{B=2}$$

$$\begin{aligned}
 A + (2) &= 3 \\
 \boxed{A=1}
 \end{aligned}$$

$$-2(1) + (2) + C = 4$$

$$-2 + 2 + C = 4$$

$$0 + C = 4$$

$$\boxed{C=4}$$

$$6B = -9$$

$$\boxed{B = -1.5}$$

7.3 Multivariable Linear Systems

HW

11/14

p485

Get partial function

54 $\frac{1}{4x^2-9} \rightarrow \frac{1}{(2x-3)(2x+3)} = \frac{A}{2x-3} + \frac{B}{2x+3}$

LCD: $(2x+3)(2x-3)$

$$\frac{A(2x+3)(2x-3)}{2x-3} + \frac{B(2x+3)(2x-3)}{2x+3}$$

$$A(2x+3) + B(2x-3)$$

$$2Ax + 3A + 2Bx - 3B$$

$$2Ax + 2Bx + 3A - 3B$$

$$x(2A+2B) + 3A-3B$$

$$2A+2B=0 \xrightarrow{\text{why}} 3A+3B=0$$

$$3A-3B=1$$

numerator

$$6A = -1$$

$$A = -\frac{1}{6}$$

$$2\left(-\frac{1}{6}\right) + 2B = 0$$

$$-\frac{1}{3} + 2B = 0$$

$$2B = \frac{1}{3}$$

$$B = \frac{1}{6}$$

57 $\frac{1}{2x^2+x} \rightarrow \frac{1}{(x)(2x+1)} = \frac{A}{x} + \frac{B}{2x+1}$

$$A(2x+1) + B(x)$$

$$2Ax + A + Bx$$

$$2Ax + Bx + A$$

$$x(2A+B) + A$$

$$2A+B=0 \rightarrow 2(0)+B=0$$

$$A=0$$

$$B=-1$$

$$-2$$

$$\rightarrow \frac{0}{x} + \frac{-1}{2x+1}$$

Why - from numerator



numerator

$$2x^2+x$$

$$x(2x+1)$$

why

= numerator



59.

$$\frac{5-x}{2x^2+x+1}$$

$$\frac{5-x}{(x+1)(2x+1)}$$

$$2x^2-1=0$$

$$2x^2-1$$

$$x=\pm\frac{1}{2}$$

$$\frac{A}{(x+1)} + \frac{B}{(2x+1)}$$

$$A(2x+1) + B(x+1)$$

$$2Ax - A + Bx + B$$

$$2Ax + Bx - A + B$$

$$x(2A+B) - A + B$$

^

Fixed Systems

multiplication error

system multiplication

$$\begin{cases} 2A+B=-1 \\ -A+B=5 \\ -2A+2B=10 \end{cases}$$

$$3B=4$$

$$B=\frac{4}{3}$$

$$-A + \left(\frac{4}{3}\right) = -1$$

$$-A = -\frac{4}{3} - 1 = -\frac{7}{3}$$

$$A = \frac{7}{3}$$

$$\frac{4}{(x+1)} + \frac{3}{(2x+1)}$$

$$60. \frac{x^2+12x+12}{x^3-4x} \rightarrow \frac{x^2+12x+12}{(x)(x^2-4)} \rightarrow \frac{x^2+12x+12}{(x)(x+2)(x-2)}$$

$$\frac{A}{x} + \frac{B}{x+2} + \frac{C}{x-2}$$

$$LCD = (x)(x+2)(x-2)$$

$$A(x+2)(x-2) + B(x)(x-2) + C(x)(x+2)$$

$$A(x^2-4) + B(x^2-2x) + C(x^2+2x)$$

$$Ax^2 - 4A + Bx^2 - 2Bx + Cx^2 + 2Cx$$

$$x^2(A+B+C) + x(-2B+2C) - 4A$$

Fixed

carry

can use matrix to ✓

$$\begin{cases} A+B+C=1 \\ -2B+2C=12 \\ -4A=12 \end{cases}$$

$$\begin{cases} -3+B=1 \\ B+C=4 \end{cases}$$

$$\begin{cases} -2(4)+2C=12 \\ 4C=20 \\ C=5 \end{cases}$$

$$A=-3$$

$$\frac{-3}{x} + \frac{1}{(x+2)} + \frac{5}{(x-2)}$$

$$\begin{cases} (-3)+B+(5)=1 \\ 2+B=1 \\ B=-1 \end{cases}$$

63. $\frac{4x^2 + 2x - 1}{x^2(x+1)} \rightarrow \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x+1}$

$\uparrow$ LCD
 $\frac{A(x^2)(x+1)}{x^2(x+1)} + \frac{B(x^2)(x+1)}{x^2(x+1)} + \frac{C(x^2)(x+1)}{x^2(x+1)}$

carry error

$A(x^2+x) + B(x+1) + C(x^2)$
 $Ax^2 + Ax + Bx + B + Cx^2$
 $Ax^2 + (x^2 + Bx + A + B)$
 $x^2(A+C) + x(A+B) + A+B$

$A+C=4$

$A+(-1)=2$

$(3)+C=4$

$A+B=2$

$(A=3)$

$(C=1)$

$(B=-1)$

$\frac{-3}{x} + \frac{2}{x^2} + \frac{7}{x+1}$

65. $\frac{27-7x}{x(x-3)^2} \rightarrow \frac{A}{x} + \frac{B}{x-3} + \frac{C}{(x-3)^2}$

$\uparrow$ LCD

$\frac{A(x)(x-3)^2}{x(x-3)^2} + \frac{B(x)(x-3)}{(x-3)^2} + \frac{C(x)(x-3)}{(x-3)^2}$

$A(x^2-9) + B(x^2-3x) + Cx$

$Ax^2 - 9A + Bx^2 - 3Bx + Cx$

$Ax^2 + Bx^2 - 3Bx + Cx - 9A$

$x^2(A+B) + x(-3B+C) - 9A$

$A+B=0$

$-3+B=0$

$-3(3)+C=-7$

$-3B+C=-7$

$(B=3)$

$-9+C=-7$

$9A=27$

$(A=3)$

$(C=2)$

$\frac{-3}{x} + \frac{3}{x-3} + \frac{2}{(x-3)^2}$

66. $\frac{x^2 - x + 2}{x(x-1)^2} = \frac{A}{x} + \frac{B}{x-1} + \frac{C}{(x-1)^2}$

$$\frac{A(x)(x-1)^2}{x} + \frac{B(x)(x-1)}{(x-1)} + \frac{C(x)(x-1)}{(x-1)^2}$$

multiplication error
 $\rightarrow A(x^2 + 1) + B(x^2 - x) + C(x)$
 $A(x^2 + 1) + B(x^2 - x) + C(x)$
 $Ax^2 - A + Bx^2 - Bx + Cx$
 $Ax^2 + Bx^2 - Bx + Cx - A$
 $x^2(A+B) + x(-B+C) + A$

$$A + B = 1$$

$$-B + C = -1$$

$$A = 2$$

$$(2) + B = 1$$

$$B = -1$$

$$-(2) + C = -1$$

$$C = 1$$

write error

$$\frac{-2}{x} + \frac{1}{x-1} + \frac{1}{(x-1)^2}$$

$\uparrow$ Use Matrix on calc to $\checkmark$

7.3 Multivariable Linear Systems Day 2

11/15

ex1

$$\frac{4x^2 - 9x - 37}{x^3 + 2x^2 - 11x - 12}$$

do synthetic division to break into parts

$$\begin{array}{r|rrrr} -1 & 1 & 2 & -11 & -12 \\ & & -1 & -1 & 12 \\ \hline & 1 & 1 & -12 & 0 \end{array}$$

$$(x+1) \frac{x^2 + x - 12}{(x+4)(x-3)} \rightarrow \frac{4x^2 - 9x - 37}{(x+1)(x+4)(x-3)}$$

$$\frac{A}{(x+1)} + \frac{B}{(x+4)} + \frac{C}{(x-3)}$$

$$\begin{aligned} A(x+4)(x-3) + B(x+1)(x-3) + C(x+1)(x+4) \\ A(x^2 + x - 12) + B(x^2 + 2x - 3) + C(x^2 + 5x + 4) \\ Ax^2 + Ax - 12A + Bx^2 - 2Bx - 3B + Cx^2 + 5Cx + 4C \\ Ax^2 + Bx^2 + Cx^2 + Ax - 2Bx + 5Cx - 12A - 3B + 4C \\ x^2(A+B+C) + x(A-2B+5C) - 12A - 3B + 4C \end{aligned}$$

$$\begin{aligned} A+B+C &= 4 \quad \rightarrow 2A+2B+2C=8 \\ A-2B+5C &= -9 \quad \rightarrow 13A+7C=-17 \\ -12A-3B+4C &= -37 \quad \rightarrow 12A+28C=-4 \end{aligned}$$

or just

do calc matrix

$$A=2$$

$$B=3$$

$$C=-1$$

$$\frac{2}{(x+1)} + \frac{3}{(x+4)} - \frac{1}{(x-3)}$$

just do matrix

Degree of numerator > degree (denominator)

$$\frac{x^3 + 2x^2 - x + 1}{x^2 + 3x - 4}$$

$$\begin{array}{r} x-1 \\ x^2+3x-4 \overline{) x^3+2x^2-x+1} \\ \underline{-(x^3+3x^2-4x)} \\ -x^2+3x+1 \\ \underline{-(-x^2-3x+4)} \\ 6x-3 \end{array}$$

$$x-1 + \frac{6x-3}{x^2+3x-4} = \frac{6x-3}{(x+4)(x-1)}$$

$$\frac{A}{x+4} + \frac{B}{x-1}$$

$$\begin{aligned} A(x-1) + B(x+4) \\ Ax - A + Bx + 4B \\ Ax + Bx - A + 4B \\ x(A+B) - A + 4B \end{aligned}$$

$$\begin{aligned} A+B &= 6 \\ -A+4B &= -3 \end{aligned}$$

$$\begin{aligned} 5B &= 3 \\ B &= \frac{3}{5} \end{aligned}$$

$$\begin{aligned} A + \left(\frac{3}{5}\right) &= 6 \\ A &= 5\frac{2}{5} \text{ or } \frac{27}{5} \end{aligned}$$

$$x-1 + \frac{\frac{27}{5}}{x+4} + \frac{\frac{3}{5}}{x-1}$$

$$x-1 + \frac{27}{5} \cdot \frac{1}{x+4} + \frac{3}{5} \cdot \frac{1}{(x-1)}$$

Remove
complex
fractions ↓

$$x-1 + \frac{27}{5(x+4)} + \frac{3}{5(x-1)}$$

485
70.

$$\frac{4x^4}{(2x-1)^3} \rightarrow \frac{4x^4}{(2x-1)(2x-1)(2x-1)} \rightarrow \frac{4x^4}{(4x^2-4x+1)(2x-1)}$$

$$\frac{4x^4}{8x^3-12x^2+6x-1}$$

$$\begin{array}{r} 8x^3-12x^2+6x-1 \overline{) 4x^4+0x^3+0x^2+0x+0} \\ \underline{-(4x^4-6x^3+3x^2-\frac{1}{2}x)} \\ 6x^3-3x^2+\frac{1}{2}x+0 \\ \underline{-(6x^3-9x^2+\frac{9}{2}x-\frac{3}{4})} \\ 6x^2-4x+\frac{3}{4} \end{array}$$

$$\frac{6x^2-4x+\frac{3}{4}}{(2x-1)^3} \cdot 4 = \frac{24x^2-16x+3}{4(2x-1)^3} \quad \text{LCM}$$

degree numerator =
degree denominator
-1

$$24x^2-16x+3 = \frac{A}{(2x-1)} + \frac{B}{(2x-1)^2} + \frac{C}{(2x-1)^3} + \frac{0}{4} \quad \text{ex } 0,500$$

$$\frac{A(4)(2x-1)^2}{2x-1} + \frac{B(4)(2x-1)}{(2x-1)^2} + \frac{C(4)(2x-1)}{(2x-1)^3}$$

$$A(4)(2x-1)^2 + B(4)(2x-1) + 4C$$

$$4A(2x-1)(2x-1) + 4B(2x-1) + 4C$$

$$4A(2x^2-4x+1) + 8Bx-4B + 4C$$

$$8Ax^2-16Ax+4A+8Bx-4B+4C$$


$$8Ax^2-16Ax+8Bx+4A-4B+4C$$

$$x^2(8A) + x(-16A+8B) + 4A-4B+4C$$

$$8A=24$$

$$-16A+8B=16 \quad \rightarrow \text{matrix on calc}$$

$$4A-4B+4C=3$$



7.3 Multivariable Linear Systems HW2

11/15

p 485

53.

$$\frac{1}{x^2-1} \rightarrow \frac{A}{(x+1)} + \frac{B}{(x-1)}$$

$$\begin{aligned} A(x+1) + B(x-1) \\ Ax - A + Bx + B \\ Ax + Bx - A + B \\ x(A+B) - A + B \end{aligned}$$

write at

$$\begin{aligned} A+B &= 0 & 2B &= 1 & A &= -.5 \\ -A+B &= 1 & B &= .5 \end{aligned}$$

write as $\frac{1}{2} \left(\frac{1}{x+1} + \frac{1}{x-1} \right)$

55.

$$\frac{1}{x^2+x} \rightarrow \frac{1}{x(x+1)} \rightarrow \frac{A}{x} + \frac{B}{x+1}$$

$$\begin{aligned} A(x+1) + B(x) \\ Ax + A + Bx \\ x(A+B) + A \end{aligned}$$

First try

$$\begin{aligned} A+B &= 0 & (1) + B &= 0 \\ A &= -1 & B &= -1 \end{aligned}$$

$$\rightarrow \left(\frac{1}{x} - \frac{1}{x+1} \right)$$

56.

$$\frac{3}{x^2-3x} \rightarrow \frac{3}{x(x-3)} \rightarrow \frac{A}{x} + \frac{B}{x-3}$$

$$\begin{aligned} A(x-3) + B(x) \\ Ax - 3A + Bx \\ x(A+B) - 3A \end{aligned}$$

$$\begin{aligned} A+B &= 0 & (-1) + B &= 0 \\ -3A &= 3 & B &= 1 \\ A &= -1 \end{aligned}$$

$$\rightarrow \left(\frac{-1}{x} + \frac{1}{x-3} \right)$$

58. $\frac{5}{x^2+x-6} \rightarrow \frac{5}{(x-2)(x+3)} \rightarrow \frac{A}{(x-2)} + \frac{B}{(x+3)}$

$$\begin{aligned} A(x+3) + B(x-2) \\ Ax + 3A + Bx - 2B \\ x(A+B) + 3A - 2B \end{aligned}$$

✓
1st $A+B=0 \quad \cdot -3 \rightarrow -3A-3B=0$
 $3A-2B=5$

$$A+(-1)=0$$

$$(A=1)$$

$$-5B=5$$

$$(B=-1)$$

$$\left(\frac{1}{(x-2)} + \frac{-1}{(x+3)} \right)$$

60. $\frac{x-2}{x^2+4x+3} \rightarrow \frac{x-2}{(x+1)(x+3)} \rightarrow \frac{A}{(x+1)} + \frac{B}{(x+3)}$

✓
1st $A(x+3) + B(x+1)$
 $Ax + 3A + Bx + B$
 $x(A+B) + 3A + B$

$$\begin{aligned} A+B &= 1 \\ 3A+B &= -2 \quad \cdot -1 \rightarrow -3A-B=2 \end{aligned} \quad \begin{aligned} -2A &= 3 \\ (A &= -1.5 \end{aligned}$$

$$\begin{aligned} (-1.5) + B &= 1 \\ +1.5 \quad +1.5 \\ (B &= 2.5 \end{aligned}$$

$$\frac{-1.5}{(x+1)} + \frac{2.5}{(x+3)}$$

62. $\frac{x^2+12x-9}{x^3-9x} \rightarrow \frac{x^2+12x-9}{x(x^2-9)} \rightarrow \frac{x^2+12x-9}{(x)(x+3)(x-3)}$

$$\frac{A}{x} + \frac{B}{x+3} + \frac{C}{x-3} \quad \text{LCD} = (x)(x+3)(x-3)$$

$$A(x+3)(x-3) + B(x)(x-3) + C(x+3)(x)$$

$$A(x^2-9) + B(x^2-3x) + C(x^2+3x)$$

$$Ax^2 - 9A + Bx^2 - 3Bx + Cx^2 + 3Cx$$

$$Ax^2 + Bx^2 + Cx^2 - 3Bx + 3Cx - 9A$$

$$x^2(A+B+C) + x(-3B+3C) - 9A$$

$$A+B+C = 1 \quad A = 1$$

$$-3B+3C = 12 \quad B = -2$$

$$-9A = -9 \quad C = 2$$

calc
matrix

$$\frac{1}{x} - \frac{2}{x+3} + \frac{2}{x-3}$$

64. $\frac{2x-3}{(x-1)^2} \rightarrow \frac{2x-3}{(x-1)(x-1)} \rightarrow \frac{A}{x-1} + \frac{B}{(x-1)^2}$

↑ LCD

$$\frac{A(x-1)^2}{\cancel{x-1}} + \frac{B(\cancel{x-1})^2}{\cancel{(x-1)}^2}$$

$$Ax - A + B$$

$$(A = 2)$$

$$-A + B = -3$$

$$-(2) + B = -3$$

$$B = -1$$

$$\frac{2}{x-1} - \frac{1}{(x-1)^2}$$

$$67, \quad \frac{2x^3 - x^2 + x + 5}{x^2 + 3x + 2} \rightarrow \frac{2x^3 - x^2 + x + 5}{(x+1)(x+2)}$$

$$\begin{array}{r} 2x - 7 \\ x^2 + 3x + 2 \overline{) 2x^3 - x^2 + x + 5} \\ \underline{-(2x^3 + 6x^2 + 4x)} \\ -7x^2 - 3x + 5 \\ \underline{-(-7x^2 - 21x - 14)} \\ 18x + 19 \end{array}$$

$$\frac{18x + 19}{(x+1)(x+2)} = \frac{A}{x+1} + \frac{B}{x+2}$$

$$A(x+2) + B(x+1)$$

$$Ax + 2A + Bx + B$$

$$x(A+B) + 2A + B$$

$$A + B = 18 \quad \begin{cases} -2A - B = -36 \end{cases} \quad \begin{cases} -B = -17 \end{cases}$$

$$2A + B = 19$$

$$B = 17$$

$$A + 17 = 18$$

$$A = 1$$

$$2x - 7 + \frac{1}{x+1} + \frac{17}{x+2}$$

68, $\frac{x^3 + 2x^2 - x + 1}{x^2 + 3x - 4}$

$$\begin{array}{r} x-1 \\ x^2+3x-4 \overline{) x^3+2x^2-x+1} \\ \underline{-(x^3+3x^2-4x)} \\ -x^2+3x+1 \\ \underline{-(-x^2-3x+4)} \\ 6x-3 \end{array}$$

Small
removing
complex
fraction
error
only

$$\frac{6x-3}{(x-1)(x+4)} = \frac{A}{x-1} + \frac{B}{x+4}$$

$$\begin{aligned} A(x+4) + B(x-1) \\ Ax + 4A + Bx - B \\ x(A+B) + 4A - B \end{aligned}$$

$$\begin{aligned} A+B &= 6 \\ 4A-B &= -3 \end{aligned} \Rightarrow \begin{cases} 5A = 3 \\ A = \frac{3}{5} \end{cases}$$

$$\begin{aligned} \left(\frac{3}{5}\right) + B &= 6 \\ B &= \frac{27}{5} \end{aligned}$$

write
as

$$x-1 + \frac{\frac{3}{5}}{x-1} + \frac{\frac{27}{5}}{x+4}$$

$$\boxed{x-1 + \frac{3}{5(x-1)} + \frac{27}{5(x+4)}}$$



[Faint, illegible handwriting in the upper middle section of the page.]

[Faint, illegible handwriting in the lower middle section of the page.]

7.3 Multivariable Linear Systems Day 3

11/16

ex1

Partial Fractions with quadratic fractions in denominator

$$\frac{3x^2 + 7x + 4}{(x^2 + 2x + 2)(x + 2)} \rightarrow \frac{Ax + B}{x^2 + 2x + 2} + \frac{C}{x + 2}$$

quadratic factor, can't factor

* degree numerator = degree denominator - 1

$$LCD = (x^2 + 2x + 2)(x + 2)$$

$$(Ax + B) \frac{(x^2 + 2x + 2) + x + 2}{x^2 + 2x + 2} + \frac{C(x^2 + 2x + 2)(x + 2)}{x + 2}$$

$$(Ax + B)(x + 2) + C(x^2 + 2x + 2)$$

$$Ax^2 + 2Ax + Bx + 2B + Cx^2 + 2Cx + 2C$$

$$x^2(A + C) + x(2A + B + 2C) + 2B + 2C$$

$$\begin{aligned} A + C &= 3 & \text{calc} & A = 2 \\ 2A + B + 2C &= 7 & \Rightarrow & B = 1 \\ 2B + 2C &= 4 & \text{matrix} & C = 1 \end{aligned}$$

$$\frac{2x + 1}{x^2 + 2x + 2} + \frac{1}{x + 2}$$

ex2

$$\frac{1}{(x^2 + x + 1)(x^2 - x + 1)} = \frac{Ax + B}{x^2 + x + 1} + \frac{Cx + D}{x^2 - x + 1}$$

forgot
other
one

other one

other one

$$(Ax + B)(x^2 - x + 1) + (Cx + D)(x^2 + x + 1)$$

$$Ax^3 - Ax^2 + Ax + Bx^2 - Bx + B + Cx^3 + Cx^2 + Cx + Dx^2 + Dx + D$$

$$x^3(A + C) + x^2(-A + B + C + D) + x(A - B + C + D) + (B + D)$$

$$\begin{aligned} A + C &= 0 \\ -A + B + C + D &= 0 \\ A - B + C + D &= 0 \\ B + D &= 1 \end{aligned}$$

matrix
calc

$$\begin{aligned} A &= +\frac{1}{2} \\ B &= \frac{1}{2} \\ C &= -\frac{1}{2} \\ D &= \frac{1}{2} \end{aligned}$$

$$\frac{+\frac{1}{2}x + \frac{1}{2}}{x^2 + x + 1} + \frac{-\frac{1}{2}x + \frac{1}{2}}{x^2 - x + 1}$$

$$\frac{x + 1}{2x^2 + 2x + 2} - \frac{x + 1}{2x^2 - 2x + 2}$$

3 cases of systems

1. Same line

∞ # of solutions
consistent system

2. Parallel lines

no solutions
inconsistent system

3. One solution

consistent



485
84,

Standard equation for a circle

$$x^2 + y^2 + Dx + Ey + F = 0$$

Find D E F

Teacher
COPY
error

but I did it
w/ #s provided

$$\begin{pmatrix} -6, -1 \\ -4, -3 \\ 2, -5 \end{pmatrix} \quad \begin{aligned} &(-6)^2 + (-1)^2 + (-6)D + (-1)E + F \\ &[-6D - E + F = -37] \end{aligned}$$

$$\begin{aligned} &(-4)^2 + (-3)^2 + (-4)D + (-3)E + F \\ &[-4D - 3E + F = -25] \end{aligned}$$

$$\begin{aligned} &(2)^2 + (-5)^2 + (2)D + (-5)E + F \\ &[2D - 5E + F = -41] \end{aligned}$$

$$\begin{aligned} -7 &= 0 \\ -13 &= E \\ -92 &= F \end{aligned}$$

$$\rightarrow -2D + 5E - F = 41$$

$$-6D + 2E = 16 \quad \times 2 \rightarrow$$

$$-12D + 4E = 32$$

$$42 + 2E = 16$$

$$E = -13$$

$$\begin{aligned} 2(-7) - 5(-13) + F &= -41 \\ -14 + 65 + F &= -41 \\ 51 + F &= -41 \end{aligned}$$

$$F = -92$$

$$-8D + 4E = 4$$

$$12D - 4E = -32$$

$$4D = -28$$

$$D = -7$$

$$\hookrightarrow x^2 + y^2 - 7x - 13y - 92$$

486

86.

Borrowed \$1 million @ @ @
 Some borrowed at 8%, 10%, 12%
 Interest = \$97,200

$$x = 2y$$

$$x + y + z = \$1,000,000$$

$$.08x + .1y + .12z = \$97,200$$

$$x = 2y$$

$$\begin{aligned} &+1 \left\{ \begin{aligned} x - 2y &= 0 \\ -x + 2y &= 0 \end{aligned} \right. \end{aligned}$$

$$3y + z = \$1,000,000$$

$$-10 \left\{ \begin{aligned} -.08x - y - .12z &= -.972,000 \end{aligned} \right.$$

$$.12x - .12z = 28,000$$

$$.5 \left\{ \right.$$

$$x - z = 140,000$$

$$x + 3y = 1,140,000$$

$$-x + 2y = 0$$

$$5y = 1,140,000$$

$$y = \$228,000$$

$$x + 3(228,000) = 1,140,000$$

$$x = \$456,000$$

$$3(228,000) + z = 1,000,000$$

$$z = \$316,000$$



1.3 Multi variable Linear Systems HW Day 3

11/17

485

77.

$$\begin{pmatrix} 0, 0 \\ 2, -2 \\ 4, 0 \end{pmatrix}$$

Find equation that passes through $(y = Ax^2 + Bx + C)$

$$A(0)^2 + B(0) + C = 0$$

$$(C=0)$$

$$A(2)^2 + B(2) + C = -2$$

$$4A + 2B + C = -2 \rightarrow 4A + 2(-2) + (0) = -2$$

$$A(4)^2 + B(4) + C = 0$$

$$16A + 4B + C = 0$$

$$4A = 2$$

$$(A = \frac{1}{2})$$

$$-16A - 8B + (0) = 8$$

$$-4B = 8$$

$$(B = -2)$$

✓! w/ calc matrix

$$\frac{1}{2}x^2 - 2x + 0 = y$$

79.

$$\begin{pmatrix} 2, 0 \\ 3, -1 \\ 4, 0 \end{pmatrix}$$

$$A(2)^2 + B(2) + C = 0$$

$$4A + 2B + C = 0$$

$$A(3)^2 + B(3) + C = -1$$

$$9A + 3B + C = -1$$

$$A(4)^2 + B(4) + C = 0$$

$$16A + 4B + C = 0$$

$$8A + 4B + 4C = 0$$

$$\rightarrow 8A + C = 0$$

$$-4B - 3C = 0$$

$$-16A - 8B - 4C = 0$$

$$-8A - C = 0$$

$$A + 3B = -1$$

or just calc

$$A = 1$$

$$B = -6$$

$$C = 8$$

$$1x^2 - 6B + 8 = y$$

81,

$$x^2 + y^2 + Dx + Ey + F = 0$$

$$(0, 0)$$

$$(2, 2)$$

$$(4, 0)$$

$$(0)^2 + (0)^2 + D(0) + E(0) + F = 0$$

$$(F = 0)$$

$$(2)^2 + (2)^2 + D(2) + E(2) + F = 0$$

$$2D + 2E + F = -8$$

$$(4)^2 + (0)^2 + D(4) + E(0) + F = 0$$

$$4D + F = -16$$

$$2(-4) + 2E + (0) = -8$$

$$-8 + 2E = -8$$

$$2E = 0$$

$$(E = 0)$$

$$4D + 0 = -16$$

$$(D = -4)$$

$$x^2 + y^2 + (-4)x + (0)y + (0)$$

$$(x^2 + y^2 - 4x)$$

83,

$$(-3, -1)$$

$$(2, 4)$$

$$(-6, 8)$$

$$(-3)^2 + (-1)^2 + D(-3) + E(-1) + F = 0$$

$$-3D - E + F = -10$$

$$(2)^2 + (4)^2 + D(2) + E(4) + F = 0$$

$$2D + 4E + F = -20$$

$$(-6)^2 + (8)^2 + D(-6) + E(8) + F = 0$$

$$(-6D + 8E + F = -100)$$

(alc

$$D = 6$$

$$E = -8$$

$$F = 0$$

$$(x^2 + y^2 + 6x - 8y)$$

85.

borrowed 775,000

A = \$ invested 8%

B = " " 9%

C = " " 10%

Interest = 67,000

 $4C = A$

Wrong
but oh
well

$$A + B + C = 775,000$$

$$.08A + .09B + .1C = 67,000$$

$$4C = A$$

$$4C - A = 0$$

$$B + 5C = 775,000$$

$$-.08A - .09B - C = -67,000$$

$$.12A + .1B = 105,000$$

$$-2A - B = -105,000$$

$$B + 5C = 75,000$$

$$-2A + 5C = -30,000$$

$$-8C + 5C = -30,000$$

$$-3C = -30,000$$

$$C = 10,000$$

$$B + 5(10,000) = 775,000$$

$$B = 725,000$$

$$4(10,000) = A$$

$$A = 40,000$$

486
92.

Sports
34 points

11 plays

A = # 6 pt plays

B = # 1 pt plays

C = # 3 pt plays

A = C

$$A + B + C = 11$$

$$6A + B + 3C = 34$$

$$A = C$$

$$A - C = 0$$

$$2A + B = 11$$

$$-3 \quad -6A - 3B = -33$$

$$-2B + 3C = 6$$

↓

$$3A - 2B = 6$$

$$4A + 2B = 22$$

$$7A = 28$$

$$A = 4$$

$$C = 4$$

$$2(4) + B = 11$$

$$8 + B = 11$$

$$B = 3$$

4 touchdowns
3 extra pts
4 field goals

7.3 Multivariable Linear Systems Day 5

11/19

1. Factoring in getting started

$$\frac{1}{x^4 - x^3} \rightarrow \frac{1}{x^3(x-1)} = \frac{A}{x-1} + \frac{B}{x} + \frac{C}{x^2} + \frac{D}{x^3}$$

degree num = degree denom - 1

$$2. \frac{1}{x^4 - 1} \rightarrow \frac{1}{(x^2+1)(x^2-1)} \rightarrow \frac{A_{x+B}}{(x^2+1)} + \frac{\cancel{C}}{\cancel{(x^2-1)}} + \frac{C}{x+1} + \frac{D}{x-1}$$

more step

$$3. \frac{x^3 - 4x^2 + 2}{(x^2+1)(x^2+2)} \rightarrow \frac{A_{x+B}}{(x^2+1)} + \frac{C_{x+D}}{(x^2+2)}$$

$$4. \frac{x^4 + x^2 + 1}{x^2(x^2+4)^2} \rightarrow \frac{x^4 + x^2 + 1}{x^2(x^2+4)(x^2+4)}$$


$$\frac{A}{x} + \frac{B}{x^2} + \frac{C_{x+D}}{x^2+4} + \frac{E_{x+F}}{(x^2+4)^2}$$

Same as linear functions

$$5. \frac{x^3 - 3x^2 + 12x - 1}{x^3 + (x^2+x+1) + (x^2+2)^3} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x^3} + \frac{D_{x+E}}{x^2+x+1}$$

$$\frac{F_{x+G}}{x^2+2} + \frac{H_{x+I}}{(x^2+2)^2} + \frac{J_{x+K}}{(x^2+2)^3}$$

can't factor any more



7.3 Multivariable Systems

Review

11/19

1. $\frac{x}{8x^2 - 10x + 3}$ $\frac{A}{4x-3} + \frac{B}{2x-1}$

$$\frac{A}{4x-3} + \frac{B}{2x-1}$$

$$\frac{A(4x-3)(2x-1)}{4x-3} + \frac{B(4x-3)(2x-1)}{(2x-1)}$$

$$A(2x-1) + B(4x-3)$$

$$2Ax - A + 4Bx - 3B$$

$$x(2A + 4B) - A - 3B$$

$$2A + 4B = 1$$

$$-A - 3B = 0$$

$$-2A - 6B = 0$$

$$-2B = 1$$

$$B = -\frac{1}{2}$$

$$2A + 4\left(-\frac{1}{2}\right) = 1$$

$$2A = 3$$

$$A = \frac{3}{2}$$

2. $\frac{9x^2 - 4x + 6}{2x^3 - x^2 - 8x + 4}$ $\frac{A}{x+2} + \frac{B}{x-2} + \frac{C}{2x-1}$

$$x^2(2x-1) - 4(2x-1)$$

$$(x^2 - 4)(2x-1)$$

$$(x+2)(x-2)(2x-1)$$

factor wrong

$$A(x-2)(2x-1) + B(x+2)(2x-1) + C(x+2)(x-2)$$

$$A(2x^2 - x - 4x + 2) + B(2x^2 - x + 4x - 2) + C(x^2 - 4)$$

$$2Ax^2 - 5Ax + 2A + 2Bx^2 + 3Bx - 2B + Cx^2 - 4C$$

$$2Ax^2 + 2Bx^2 + Cx^2 - 5Ax + 3Bx + 2A - 2B - 4C$$

$$x^2(2A + 2B + C) + x(-5A + 3B) + 2A - 2B - 4C$$

$$2A + 2B + C = 9$$

$$\rightarrow A = 3$$

$$\frac{3}{x+2} + \frac{2}{x-2} - \frac{1}{2x-1}$$

$$-5A + 3B = -9$$

$$\rightarrow B = 2$$

$$2A - 2B - 4C = 6$$

$$\rightarrow C = -1$$

$$2 - 2 - 4$$

$$3. \frac{x-3}{x^3+3x} \rightarrow \frac{x-3}{x(x^2+3)} \rightarrow \frac{A}{x} + \frac{Bx+C}{x^2+3}$$

$$A(x^2+3) + (Bx+C)(x) \\ Ax^2 + 3A + Bx^2 + Cx \\ x^2(A+B) + x(C) + 3A$$

$$A+B=0$$

$$-1+B=0$$

$$C=1$$

$$B=1$$

$$3A=-3$$

$$A=-1$$

$$\frac{-1}{x} + \frac{1x+1}{x^2+3}$$

$$4. \frac{x^5 - 2x^4 + x^3 + x + 5}{x^3 - 2x^2 + x - 2}$$

$$\begin{array}{r} x^3 - 2x^2 + x - 2 \overline{) x^5 - 2x^4 + x^3 + 0x^2 + x + 5} \\ \underline{-(x^5 - 2x^4 + x^3 - 2x^2)} \\ 4x^2 + x + 5 \\ \underline{-(4x^2 - 7x^3 - x + 2x)} \\ 7x^3 + 8x + 3 \\ \underline{-(7x^3 + 3x^2 - 1x + 5)} \\ -2x^2 + 4x - 2 \\ \underline{-(-2x^2 + 4x - 2)} \\ 0 \end{array}$$

bad subtraction

remainder

4
error 11

$$\frac{-x^2 + 3x + 9}{x^3 - 2x^2 + x - 2} = x^2(x-2) + 1(x-2) \\ (x^2+1)(x-2)$$

$$\frac{Ax+B}{x^2+1} + \frac{C}{x-2} = \frac{(Ax+B)(x-2) + C(x^2+1)}{(x^2+1)(x-2)} \\ Ax^2 - 2Ax + Bx - 2B + Cx^2 + C \\ x^2(A+C) + x(-2A+B) + C - 2B$$

copy error

carry error

copy error

$$\downarrow A + C = -1 \quad 2$$

$$-2A + B = 1$$

$$+C = 5$$

$$C = 5$$

$$A + \overset{5}{(+1)} = -1$$


$$A = -6$$

$$-(-6) + B = 3$$

$$B = -3$$

$$x^2 + \frac{-6x-3}{x^2+1} + \frac{5}{x-2}$$

still wrong



8.1 Sequences + Series

Notes

11/27

A sequence has order
Has a first number (position)
A second position, etc

Is a function whose domain is the set of positive integers

ex: a_1, a_2, a_3, a_4, a_n
 $1, 4, 7, 10, \dots$ (infinite sequence)
 $\downarrow \quad \downarrow \quad \downarrow \quad \downarrow$
 $+3 \quad +3 \quad +3 \quad +3$

ex: $2, 4, 2, 4, \dots$ (oscillating sequence)

ex: $1, \frac{1}{4}, \frac{1}{9}, \frac{1}{16}, \frac{1}{25}, \dots$

Finite sequence - has a specific number of terms

consists of the first n positive integers

ex The first 4 terms of the sequence given by

$$a_n = 3n - 2$$

$$a_1 = n = 1 \rightarrow 1$$

$$a_2 = n = 2 \rightarrow 4$$

$$a_3 = n = 3 \rightarrow 7$$

$$a_4 = n = 4 \rightarrow 10$$

P1

$$a_n = 3 + (-1)^{n-1}$$

$$a_1 = 2$$

$$a_2 = 4$$

$$a_3 = 2$$

$$a_4 = 4$$

563
18.

$$(-1)^n \left(\frac{n}{n+1} \right)$$

$$\begin{aligned} a_1 &= -.5 & -\frac{1}{2} \\ a_2 &= .667 & \Rightarrow \frac{2}{3} \\ a_3 &= -.75 & -\frac{3}{4} \\ a_4 &= .8 & \frac{4}{5} \\ a_5 &= -.833 & -\frac{5}{6} \end{aligned}$$


30.

$$a_n = \frac{3n^2}{n^2+1}$$

→

Same degree

$$asy = \frac{3}{1} \Rightarrow 3$$



To generate sequence - need more than 2 terms

ex

$$\begin{aligned} \text{ex: } & \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16} \dots \frac{1}{2^n} \\ (n) = & 1 \quad 2 \quad 3 \quad 4 \quad \rightarrow \end{aligned}$$

ex

$$1, 3, 5, 7 \dots 2n-1 \leftarrow y\text{-intercept}$$

↑ slope

ex

$$2, 5, 10, 17 \dots n^2 + 1$$

ex

$$1, \frac{1}{4}, \frac{1}{9}, \frac{1}{16}, \frac{1}{25} \dots \frac{1}{n^2}$$

564

44.

$$\frac{2}{1}, \frac{3}{2}, \frac{4}{3}, \frac{5}{4}, \frac{6}{5} \dots \frac{n+1}{2n-1}$$

46.

$$\frac{1}{3}, \frac{-2}{9}, \frac{4}{27}, \frac{-8}{81} \dots \frac{-2^{(n-1)}}{3^n}$$

48.

$$1 + \frac{1}{2}, 1 + \frac{3}{4}, -1 + \frac{7}{8}, 1 + \frac{15}{16}, 1 + \frac{31}{32} \dots 1 + \frac{2^{n-1}}{2^n}$$

← look at numerator + denominator separately

564
52, 1, -1, 1, -1, 1 ... $(-1)^{n+1}$

Some sequences are defined recursively
- uses a previous term to predict future term

ex

Fibonacci Sequence

1, 1, 2, 3, 5, 8, 13, 21, 34
 $a_0, a_1, a_2, a_3, a_4, \dots$

add the previous two to get to the next term

$$a_k = a_{k-2} + a_{k-1}$$

$$a_4 = a_{4-2} + a_{4-1}$$

$$a_2 + a_3$$

$$2 + 3$$

$$a_4 = 5$$

54, $a_{k+1} = a_k + 3$

↳ recursive sequence

$$a_1 = 15$$

$$a_2 = 18 \quad \downarrow +3$$

$$a_3 = 21$$

$$a_4 = 24$$

$$a_5 = 27$$

58, $a_{k+1} = a_k - 5$

$$a_1 = 25$$

$$a_2 = 20$$

$$a_3 = 15$$

$$a_4 = 10$$

$$a_5 = 5$$

Normal rule

$$a_n = 30 - 5n$$

$$a_n = 5(6-n) \quad \text{or}$$

60. $a_{k+1} = -2a_k$

$$a_1 = 14$$

$$a_2 = -2(14) = -28$$

$$a_3 = -2(-28) = 56$$

$$a_4 = -2(56) = -112$$

$$a_5 = -2(-112) = 224$$

Normal Rule(s)

$$\frac{-7(-2^n)}{(-1)^n(n+1)(-7)}$$

$$14(-2^{(n-1)})$$

8.1 Sequences + Series HW

11/27

563 5, $a_n = \left(-\frac{1}{2}\right)^n$

$$\begin{aligned}a_1 &= -\frac{1}{2} \\a_2 &= \frac{1}{4} \\a_3 &= -\frac{1}{8} \\a_4 &= \frac{1}{16} \\a_5 &= -\frac{1}{32}\end{aligned}$$

11, $a_n = \frac{1 + (-1)^n}{n}$

$$\begin{aligned}a_1 &= 0 \\a_2 &= 1 \\a_3 &= 0 \\a_4 &= \frac{1}{2} \\a_5 &= 0\end{aligned}$$

15, $a_n = \frac{1}{n^{3/2}}$

$$\begin{aligned}a_1 &= 1 \\a_2 &= .35 \\a_3 &= .19 \\a_4 &= \frac{1}{8} \\a_5 &= .091\end{aligned}$$

19, $a_n = (2n-1)(2n+1)$


$$\begin{aligned}a_1 &= 3 \\a_2 &= 15 \\a_3 &= 35 \\a_4 &= 63 \\a_5 &= 99\end{aligned}$$

21, $a_n = (-1)^n (3n-2)$ $a_{25} = -73$

23, $a_n = \frac{n^2}{n^2+1}$ $a_{10} = .990$

Graph

27. $a_n = 16(-.5)^{n-1}$




33. $a_n = 1 + \frac{n+1}{n}$

n	a_n
1	3
2	2.5
3	$2\frac{1}{3}$
4	$2\frac{1}{4}$
5	$2\frac{1}{5}$
6	$2\frac{1}{6}$
7	$2\frac{1}{7}$
8	$2\frac{1}{8}$
9	$2\frac{1}{9}$
10	$2\frac{1}{10}$

564

37. $a_n = 4(.5)^{n-1}$



Write sequence

39. 1, 4, 7, 10, 13

$a_n = 3n - 2$

41. 0, 3, 8, 15, 24

$a_n = n^2 - 1$

43. $\frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \frac{5}{6}, \frac{6}{7}$

$a_n = \frac{n+2}{n+3}$

try to make an $a_0 = -2$
 $\nearrow$ y-int piece

45, $\frac{1}{2}, -\frac{1}{4}, \frac{1}{8}, -\frac{1}{16}$ $a_n = \frac{(-1)^{n+1}}{2^n}$

47, $1 + \frac{1}{1}, 1 + \frac{1}{2}, 1 + \frac{1}{3}, 1 + \frac{1}{4}, 1 + \frac{1}{5}$ $a_n = 1 + \frac{1}{n}$

49, $1, \frac{1}{2}, \frac{1}{6}, \frac{1}{24}, \frac{1}{120}$ $a_n = \frac{1}{n!}$

51, $1, 3, 1, 3, 1$ $a_n = \frac{(-1)^n + 2(1)^n}{(-1)^n + 2}$

57, $a_{k+1} = a_k + 2$

Normal

$a_1 = 6$


$a_2 = 8$

$a_3 = 10$

$a_4 = 12$

$a_5 = 14$

$4 + 2n$



8.1 Sequences + Series

Day 2

11/28

Factorials

ex

I have 4 ice cream flavors. How many unique combos are there?

e

Flavors = C, V, S, B

CVSB	VCSB		
CVBS	VCBS		
CBVS	VBCS	→ ...	→ ...
CBSV	VBSC		
CSBV	VSCB		
CSVB	VSBC		
6	6	6	6 = 24

$$4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24$$

✓ = ⊙

ex2

$$8! = 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 40320$$

↑ can type in calc [8] [MATH] [=] [4] [Enter]

ex3

$2n!$ n can start at 0

$$0! = 1$$

$$a_0 = 2(0)! = 2 \quad \text{~~can't calc~~ } 2n! = 2 \underbrace{(1 \cdot 2 \cdot 3 \cdot 4 \cdot n)}_{n!}$$

$$a_1 = 2(1)! = 2$$

$$a_2 = 2(2 \cdot 1) = 4$$

$$a_3 = 2(3 \cdot 2 \cdot 1) = 12$$

ex4

$$a_n = \frac{2^n}{n!}$$

$$a_0 = 1$$

$$a_1 = 2$$

$$a_2 = \frac{2}{2}$$

$$a_3 = \frac{4}{3}$$

$$a_4 = \frac{2}{3}$$

~~can't calc but is 1 when n=0~~

564

$$62, a_n = \frac{1}{(n+1)!}$$

$$\begin{aligned} a_0 &= 1 \\ a_1 &= \frac{1}{2} \\ a_2 &= \frac{1}{6} \\ a_3 &= \frac{1}{24} \\ a_4 &= \frac{1}{120} \end{aligned}$$

$$64, a_n = \frac{(-1)^{2n+1}}{(2n+1)!}$$

$$\begin{aligned} a_0 &= -1 \\ a_1 &= -\frac{1}{2} \\ a_2 &= -\frac{1}{120} \\ a_3 &= -\frac{1}{5040} \\ a_4 &= -\frac{1}{362880} \end{aligned} \quad \left. \begin{array}{l} \\ \\ \\ \\ \end{array} \right\} \text{ same as } \frac{1}{(2n+1)!}$$

$$68, \frac{5!}{7!} = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = \frac{1}{7 \cdot 6} = \frac{1}{42}$$

$$70, \frac{10! \cdot 3!}{4! \cdot 6!} = \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 3 \cdot 2 \cdot 1}{4 \cdot 3 \cdot 2 \cdot 1 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = \frac{10 \cdot 9 \cdot 8 \cdot 7}{1} = 1260$$

$$72, \frac{(n+2)!}{n!} = \frac{(n+2)(n+1)\overbrace{n!}^4}{n!} = (n+2)(n+1) \quad \uparrow \text{ solution}$$

$$74, \frac{(2n+2)!}{2n!} = \frac{(2n+2)(2n+1)\overbrace{2n!}^{2n!}}{2n!} = (2n+2)(2n+1)$$

Summation (Sigma)

Adds (sums) the terms of a sequence

Σ $\in$ sigma - greek letter

n $\in$ upper boundary (limit)

$\sum a_i$ $\in$ index (what you are doing)

$i=1$ $\in$ lower boundary (start)
 $\in$ do index

$a_1 + a_2 + a_3 + a_4 + a_n \dots$
 $\nearrow$ start i here $\nearrow$ to upper limit

ex

$$\sum_{i=1}^5 3i = 3(1) + 3(2) + 3(3) + 3(4) + 3(5)$$

$3 + 6 + 9 + 12 + 15$
 45

$\nearrow$ stop

ex2

$$\sum_{k=3}^6 (1+k^2) = (1+(3)^2) + (1+(4)^2) + (1+(5)^2) + (1+(6)^2)$$

$10 + 17 + 26 + 37$
 90

564

76.

$$\sum_{i=1}^6 (3i-1) = (3(1)-1) + (3(2)-1) + (3(3)-1) + (3(4)-1) + (3(5)-1) + (3(6)-1)$$

$2 + 5 + 8 + 11 + 14 + 17$

78.

$$\sum_{k=1}^5 6 = 6 + 6 + 6 + 6 + 6$$

$6 \cdot 5$
 30

57

ex $\sum_{i=1}^{10} \frac{3}{i+1}$ Calc: $\left[\begin{array}{c} 2nd \\ " \end{array} \right] \left[\begin{array}{c} List \\ " \end{array} \right] \left[\begin{array}{c} \leftarrow \\ \rightarrow \end{array} \right] \left[\begin{array}{c} 5 \\ 5 \end{array} \right] \text{sum(}$
 $\left. \begin{array}{c} seq(\\ 3/(n+1) \\ , \\ n \\ , \\ 10 \\) \end{array} \right) \left[\begin{array}{c}) \\ Enter \end{array} \right]$
 6.06

94. $\left[1 - \left(\frac{1}{6} \right)^2 \right] + \left[1 - \left(\frac{2}{6} \right)^2 \right] + \dots + \left[1 - \left(\frac{6}{6} \right)^2 \right]$ Write as $\sum$
 $\sum_{i=1}^6 \left[1 - \left(\frac{i}{6} \right)^2 \right] = 3.4722$
 realc

98. $\frac{1}{1 \cdot 3} + \frac{1}{2 \cdot 4} + \frac{1}{3 \cdot 5} + \dots + \frac{1}{10 \cdot 12} \sum_{i=1}^{10} \frac{1}{i \cdot (i+2)}$

Properties of Sums

1. $\sum_{i=1}^n c = n \cdot c \rightarrow \sum_{i=1}^3 5 = 5 + 5 + 5$
 $3 \cdot 5 = 15$

2. $\sum_{i=1}^n c \cdot a_i = c \sum_{i=1}^n a_i \rightarrow \sum_{i=1}^3 3 \cdot i^2 = 3(1^2 + 2^2 + 3^2)$
 $3(14) = 42$

$$3. \sum_{i=1}^n (a_i + b_i) = \sum_{i=1}^n a_i + \sum_{i=1}^n b_i$$

$$4. \sum_{i=1}^n (a_i - b_i) = \sum_{i=1}^n a_i - \sum_{i=1}^n b_i$$

Sum the terms of a finite or infinite sequence, it's called a series (not sequence)

Partial sums = $\sum_{i=1}^n a_i$

Adding all terms of an infinite sequence

$$\sum_{i=1}^{\infty} a_i$$

Ex $\sum_{i=1}^{\infty} \frac{3}{10^i} \in \text{infinite series}$

a) Find the 3rd partial sum

$$\sum_{i=1}^3 \frac{3}{10^i} = \frac{3}{10^1} + \frac{3}{10^2} + \frac{3}{10^3} = .333$$

b) Find the sum of the series

.3 + .03 + .003 + .0003 ...

$$\left(\frac{1}{3}\right) \in \text{limit} \swarrow .33333333$$

565
106.

$$\sum_{i=1}^{\infty} 4\left(\frac{1}{10}\right)^i = 4\left(\frac{1}{10}\right)^1 + 4\left(\frac{1}{10}\right)^2 + 4\left(\frac{1}{10}\right)^3 + 4\left(\frac{1}{10}\right)^4 + \dots$$

$.4 + .04 + .004 + .0004 + \dots$
 $.4444$
 $\frac{4}{9}$

Sequences + Series

Day 2 HW

11/28

564

57. $a_{k+1} = a_k + 2$

$a_1 = 6$

$a_2 = 8$

$4 + 2n$

$a_3 = 10$

$a_4 = 12$

$a_5 = 14$

59. $a_{k+1} = \frac{1}{3} a_k$

$a_1 = 81$

$a_2 = 27$

$a_3 = 9$

$a_4 = 3$

$a_5 = 1$

65. $a_n = \frac{(-1)^n}{(2n)!}$

$a_0 = 1$

$a_1 = \frac{1}{2}$

$a_2 = \frac{1}{24}$

$a_3 = \frac{1}{720}$

$a_4 = \frac{1}{40320}$

Simplify,

69. $\frac{12!}{4 \cdot 8!}$

$12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$
 $4 \cdot 3 \cdot 2 \cdot 1 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$

$\frac{11880}{24} = 495$

71. $\frac{(n+1)!}{n!}$

$\frac{(n+1)(n!)}{n!}$

$n+1$

77.

$\sum_{k=1}^4 10 = 4 \cdot 10 = 40$

$$83. \sum_{i=1}^4 [(i-1)^2 + (i+1)^3] = [(1-1)^2 + (1+1)^3] + [(2-1)^2 + (2+1)^3] \\ + [(3-1)^2 + (3+1)^3] + [(4-1)^2 + (4+1)^3] = 238$$

$$89. \sum_{k=0}^4 \frac{(-1)^k}{k+1} = \frac{47}{60}$$

$$93. \left[2\left(\frac{1}{8}\right) + 3 \right] + \left[2\left(\frac{2}{8}\right) + 3 \right] + \dots + \left[2\left(\frac{8}{8}\right) + 3 \right] \\ \sum_{i=1}^8 \left[2\left(\frac{i}{8}\right) + 3 \right] = 33$$

$$95. 3 - 9 + 27 - 81 + 243 - 729 \sum_{i=0}^3 (-1)^{i+1} 3^i = -546$$

↑ looked up
↓ makes odd ⊖

$$97. \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots - \frac{1}{20^2} = \sum_{i=1}^{20} \frac{(-1)^{i+1}}{n^2} \approx 2.821$$

$$99. \frac{1}{4} + \frac{9}{8} + \frac{7}{16} + \frac{15}{32} + \frac{31}{64} = \sum_{i=1}^5 \frac{n-1}{2} \frac{2^i-1}{2^{i+1}} = 2.0156$$

$$101. \sum_{i=1}^{\infty} 5\left(\frac{1}{2}\right)^i \quad \text{4th partial sum} \\ = 5\left(\frac{1}{2}\right)^1 + 5\left(\frac{1}{2}\right)^2 + 5\left(\frac{1}{2}\right)^3 + 5\left(\frac{1}{2}\right)^4 \\ = \frac{5}{2} + \frac{5}{4} + \frac{5}{8} + \frac{5}{16} = \frac{75}{16}$$

$$103. \sum_{n=1}^{\infty} 4\left(-\frac{1}{2}\right)^n \quad \text{3rd partial sum} \\ = 4\left(-\frac{1}{2}\right)^1 + 4\left(-\frac{1}{2}\right)^2 + 4\left(-\frac{1}{2}\right)^3 \\ = -2 + 1 - \frac{1}{2} = -\frac{3}{2}$$

∞ Find sum of infinite series

105. $\sum_{i=1}^{\infty} 6\left(\frac{1}{10}\right)^i$ $6\left(\frac{1}{10} + \frac{1}{100} + \frac{1}{1000} \dots\right) = 6(0,111) = \frac{6}{9}$

107. $\sum_{k=1}^{\infty} \left(\frac{1}{10}\right)^k$ $\frac{1}{10} + \frac{1}{100} + \frac{1}{1000} \dots = 0,111 = \frac{1}{9}$

109. \$5,000 interest
3% interest compounded quarterly

$A_n = 5000 \left(1 + \frac{0.03}{4}\right)^n$ $n = \text{number of quarters}$
balance

$a_1 = 5037,5$

$a_2 = 5075,28$

$a_3 = 5113,35$

$a_4 = 5151,69$

$a_5 = 5190,33$


$a_6 = 5229,26$

$a_7 = 5268,48$

$a_8 = 5308$

$a_{40} = 6741,74$

III, $a_n = 1,37n^2 + 3,1n + 698$ $n = 3 = 1093$



Enrollment is rising

566

113,

Net Profit

$$a_n = 2.15n^2 + 235.9$$

$$n = 4 = 1994$$

\$ 3,491,000,000 same as bar graph

8.2 Arithmetic Sequences + Partial Sums Notes

11/29

Arithmetic sequences - difference b/w consecutive terms
which is the same (constant rate of change)

common difference = d

ex

$$7, 11, 15, 19 \quad \text{so } d = 4$$

$\swarrow \quad \swarrow \quad \swarrow$
 $+4 \quad +4 \quad +4$

get $a_0 = 3$ $\leftarrow$ y-intercept

$$\text{so } a_n = mx + b \rightarrow \boxed{4n + 3}$$

ex2

$$2, -3, -8, -13 \quad d = -5$$

$\swarrow \quad \swarrow \quad \swarrow$
 $-5 \quad -5 \quad -5$

$$\rightarrow a_0 = 7$$

$$\boxed{a_n = -5n + 7}$$

P1

$$1, \frac{5}{4}, \frac{3}{2}, \frac{7}{4}$$

$\swarrow \quad \swarrow \quad \swarrow$
 $\frac{1}{4} \quad \frac{1}{4} \quad \frac{1}{4}$

$$d = \frac{1}{4}$$

$$\boxed{y = \frac{1}{4}x + \frac{3}{4}}$$

P2

$$1, 4, 9, 16$$

not arithmetic

The n th term of an arithmetic sequence is the
form of $mx + b$ or $dn + c = a_n$

d = common difference

c = y-intercept value ($c = a_1 - d$)

ex3

$$2, 6, 10, 14, 18 \quad \boxed{a_n = 4n - 2}$$

$a_0 = -2$

573

12. $a_n = \frac{1 + (n-1)4}{1 + 4(n-1)}$ recursive
 $4n-3$ not recursive

22. 10, 5, 0, -5, -10
 $a_n = -5n + 15$

24. $a_1 = -4$ $20 \overset{5-1}{/} 4 = 5$
 $a_5 = 16$
 $a_n = 5n - 9$

26. $a_5 = 190$ $115-180$
 $a_{10} = 115$ $-75/5 = -15$
 $a_n = -15n + 265$

30. $a_4 = 16$ $30/6 = 5 = d$
 $a_{10} = 46$
 $a_0 = a_4 - 4d$ $a_n = 5n - 4$
 $a_0 = 16 - 20$
 $a_0 = -4$

$$a_n = dn + c \quad c = a_1 - d$$

$$a_n = dn + (a_1 - d)$$

$$a_n = dn - d + a_1$$

$$a_n = d(n-1) + a_1$$

Proof

36,

$$a_1 = 200$$

$$a_{k+1} = a_k - 10$$

$$a_1 = 200$$

$$a_2 = 190$$

$$a_3 = 180$$

$$a_4 = 170$$

$$a_5 = 160$$

$$-10n + 210 = a_n$$

40.

$$a_1 = 3$$

$$a_7 = 13$$

$$10n - 7 = a_n$$

$$a_9 = 83$$

Sum of a Finite Arithmetic Sequence

$$S_n = \frac{n}{2} (a_1 + a_n)$$

↑ # of
terms

ex

$$1 + 3 + 5 + 7 + 9 + 11 + 13 + 15 + 17 + 19$$

↑ $n = 10$ terms

↑ a_{10}

$$S_{10} = \frac{10}{2} (1 + 19)$$


100

Sum of integers from 1 to 100

$$S_n = \frac{100}{2} (1 + 100)$$

$$50 \cdot 101$$

$$5050$$



P3

$$a_1 = 15$$

$$a_{100} = 307$$

Probably infinite
Find the partial sum

$$S_{100} = \frac{100}{2} (15 + 307)$$

$$50 \times 322$$

$$16100$$

P4

Find the sum of the integers from -10 to 50

$$S_{60} = \frac{60}{2} (-10 + 50)$$

$$30 \times 40$$

$$1200$$

P5

Find the 150th partial sum of this sequence

$$5, 16, 27, 38, 49 \quad d = 11 \quad a_n = 11n - 6$$

$$5 - 11 = -6 = a_0$$

$$11(150) - 6 = 1644$$

$$S_{150} = \frac{150}{2} (5 + 1644)$$

$$123675$$

P574

56.

$$4.2, 3.7, 3.2, 2.7$$

$$-1.5n + 4.7$$

Find sum of first 12 terms

$$a_{12} = -1.5(12) + 4.7$$

$$-6 + 4.7$$

$$-1.3$$

$$S_{12} = \frac{12}{2} (4.2 - 1.3)$$

$$17.4$$

64, $\sum_{n=51}^{100} 7n$ Find sums $S_{50} = \frac{50}{2} (357 + 700)$ $\downarrow$ 7.51

52850

66. $\sum_{n=51}^{100} n - \sum_{n=1}^{50}$ $\rightarrow \frac{50}{2} (1+50)$

1275

$\downarrow$

$S_{50} = \frac{50}{2} (51+100)$

3775

$3775 - 1275$

2500

8.2 Arithmetic Sequences & Partial Sums

11/29

573/7

Arithmetic?
3, 7, 4, 3, 4, 9, 5, 5, 6, 1 - Arithmetic
 $D = .6$

11, $\frac{1}{n+1}$ $a_1 = \frac{1}{2}$
 $a_2 = \frac{1}{3}$ Not arithmetic
 $a_3 = \frac{1}{4}$
 $a_4 = \frac{1}{5}$
 $a_5 = \frac{1}{6}$

13, $a_n = 158 - 7n$ $a_1 = 143$
 $a_2 = 136$ $d = 7$
 $a_3 = 129$
 $a_4 = 122$
 $a_5 = 115$

19, $a_1 = 180$ $108 - 8n$
 $d = -8$

21, $4\frac{3}{2}, -1, -\frac{7}{2}$ $6.5 - 2.5n$
 $d = -2.5$


25, $a_3 = 94$ $-9/3 = -3$
 $a_6 = 85$ $105 - 3n$
 $a_0 = 94 + 3(-3)$
 105

35, $a_{k+1} = a_k + 4$ $a_1 = 15$
 $a_2 = 19$ $11 + 4n$
 $a_3 = 23$
 $a_4 = 27$
 $a_5 = 31$

37. $a_{k+1} = a_k - \frac{1}{4}$ $a_1 = \frac{7}{2}$
 $a_2 = 3\frac{1}{4}$
 $a_3 = 3$ $3.75 - .25 =$
 $a_4 = 2\frac{3}{4}$
 $a_5 = 2\frac{1}{2}$

41. $a_1 = 4.2$ $a_7 = 18.6$
 $a_2 = 6.6$ $1.8 + 2.4n$

45. $a_n = .5n + 4$



574

55. $1.5, 1.3, 2.1, 2.9 \dots n=10$
 $+1.8$

Find Partial sum

$a_n = -.3 + .8n$
 $a_{10} = -.3 + .8(10)$
 7.7

(✓)

$S_{10} = \frac{10}{2} (.5 + 7.7)$
 $S_{10} = 41$

57. $a_1 = 100$ $n = 25$ $S_{25} = \frac{25}{2} (100 + 220)$
 $a_{25} = 220$ 4000

59. First 100 odd positive integers
 $1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21$

$a_1 = 1$
 $a_{100} = 199$

$S_{100} = \frac{100}{2} (1 + 199)$
 $10,000$

$a_n = a_1 + (n-1)d$
 $1 + (n-1)2$
 $1 + 2n - 2$
 $2n - 1$

65. $\sum_{n=1}^{30} n - \sum_{n=1}^{10} n \rightarrow S_{10} = \frac{10}{2} (1+10)$
 $\downarrow$
 $S_{20} = \frac{20}{2} (11+30)$
 $410 - 55$
 355

67. $\sum_{n=1}^{500} (n+8)$ $S_{500} = \frac{500}{2} (9+508)$
 129250

71. $\sum_{n=1}^{100} \frac{n+4}{2}$ $S_{100} = \frac{100}{2} (2.5+52)$
 2725

75. Salary = \$32,500 $a_n = 31000 + 1500n$
 Annual raise = \$1,500 $a_6 = 40,000$

$S_6 = \frac{6}{2} (32500 + 40000)$
 $217,500$

79. Seating Capacity

15 rows

First row has 25 seats

Each row has 2 more seats

$23+2n$

$S_{15} = \frac{15}{2} (25 + 53)$
 585 seats

575
81

Sales

\$20,000 1st year

15,000 + 5,000

\$5,000 more each year for 4 years

Profit for first 5 years

$$S_5 = \frac{5}{2} (20,000 + 40,000)$$

\$150,000

8.3 Geometric Sequences + Series Notes

11/30

Consecutive terms of a geometric sequence have a common ratio
multiplier

$r = \text{common ratio, } r \neq 0$

ex, $2, 4, 8, 16$ $r = 2$ $= 2^n$
 $\frac{4}{2} = 2$ $\frac{16}{8} = 2$

ex2 $-\frac{1}{3}, \frac{1}{9}, -\frac{1}{27}, \frac{1}{81}$

$\frac{1/9}{-1/3} = -\frac{1}{3}$

582

8 $9, -6, 4, -\frac{8}{3}$ $r = -\frac{2}{3}$
 $\frac{-6}{9} = -\frac{2}{3}$

10. $\frac{1}{5}, \frac{2}{7}, \frac{3}{9}, \frac{4}{11}$

$\frac{2/7}{1/5} = \frac{2}{7} \cdot \frac{5}{1} = \frac{10}{7} = r$ for that - but not geometric

N^{th} term of a geometric sequence

$a_n = a_1 \cdot r^{n-1}$

$r^1 \left\{ \begin{array}{l} a_1 = a_1 \cdot r^{1-1} = a_1 \cdot r^0 \\ a_2 = a_1 \cdot r^{2-1} = a_1 \cdot r^1 \\ a_3 = a_1 \cdot r^2 \\ a_4 = \\ a_5 = \\ a_n = \end{array} \right.$

$$14. \quad r = \frac{1}{3}$$

$$a_1 = 2$$

$$a_2 = 2 \cdot \left(\frac{1}{3}\right)^{2-1} = \frac{2}{3}$$

$$a_3 = 2 \cdot \left(\frac{1}{3}\right)^2 = \frac{2}{9}$$

$$a_4 = 2 \cdot \left(\frac{1}{3}\right)^3 = \frac{2}{27}$$

$$a_5 = 2 \cdot \left(\frac{1}{3}\right)^4 = \frac{2}{81}$$

$$18. \quad r = \sqrt{3}$$

$$a_1 = 4$$

$$a_2 = 4 \cdot \sqrt{3} = 4\sqrt{3}$$

$$a_3 = 4 \cdot (\sqrt{3})^{3-1} = 12$$

$$a_4 = 4 \cdot (\sqrt{3})^3 = 4 \cdot 3 \cdot \sqrt{3} = 12\sqrt{3}$$

$$a_5 = 4 \cdot (\sqrt{3})^4 = 36$$

$$20. \quad a_{k+1} = \frac{1}{3} a_k$$

$$a_1 = 81$$

$$a_2 = 27$$

$$a_3 = 9$$

$$a_4 = 3$$

$$a_5 = 1$$

$$26. \quad r = \frac{3}{2}$$

$$a_1 = 5$$

$$a_8 = 5 \cdot \left(\frac{3}{2}\right)^{8-1} = 5 \cdot \left(\frac{3}{2}\right)^7 = 85.43$$

$$a_5 = a_3 \cdot r^2$$

$$\frac{64}{27} = \frac{16}{3} \cdot r^2$$

$$\frac{16}{3} \cdot \frac{27}{64} = r^2 = \frac{9}{8} \Rightarrow \sqrt[2]{\frac{9}{8}} = \pm \frac{3}{2\sqrt{2}}$$

32. Geometric

$$a_3 = \frac{16}{3}$$

$$a_5 = \frac{64}{27}$$

$$a_7 = ?$$

$$a_5 = a_3 \cdot r^2$$

$$\frac{64}{27} = \frac{16}{3} \cdot \left(\frac{3}{2}\right)^2 \rightarrow \frac{256}{243}$$

ex

$$a_4 = 125$$

$$a_{10} = 125/64$$

$$a_{14} = ?$$

$$a_{10} = a_4 \cdot r^6$$

$$\frac{125}{64} = 125 \cdot r^6$$

$$\frac{1}{64} = r^6$$

$$\frac{1}{2} = r$$

$$a_{14} = a_{10} \cdot r^4$$

$$a_{14} = 125/64 \cdot \left(\frac{1}{2}\right)^4$$

$$a_{14} = \frac{125}{1024}$$

34.

$$a_1 = 3 \quad \downarrow \quad \frac{36}{3} = 12$$

$$a_2 = 36$$

$$a_3 = 432$$

$$a_7 = 3 \cdot (12)^6$$

$$8,957,952$$

Sum of a finite geometric sequence

$$S_n = \sum_{i=1}^n a_i \cdot r^{i-1} = a_1 \left(\frac{1-r^n}{1-r} \right)$$

ex

$$\sum_{n=1}^{12} 4(.3)^n = a_1 + a_2 + a_3 + \dots + a_{12}$$

$$4(.3)^1 + 4(.3)^2 + 4(.3)^3 + \dots + 4(.3)^{12}$$

$$a_1 = 4(.3) = 1.2$$

$$r = .3$$

$$n = 12$$

$$a_1 \left(\frac{1-r^n}{1-r} \right)$$

$$1.2 \left(\frac{1-.3^{12}}{1-.3} \right)$$

$$\approx 1.714$$

583

50.

$$\sum_{h=0}^{15} 2 \left(\frac{4}{3} \right)^h$$

$$2 \left(\frac{1 - \frac{4}{3}^{16}}{1 - \frac{4}{3}} \right) = 592.65$$

$\nearrow$ 1st term

$$2 \left(\frac{4}{3} \right)^0$$

$$58. \quad 15 - 3 + \frac{3}{5} - \dots - \frac{3}{625}$$

$$\frac{-3}{15} = -\frac{1}{5} = r$$

$\rightarrow$ Write sequence in summation

$$\sum_{n=1} 15 \left(-\frac{1}{5} \right)^{n-1} = \frac{-3}{625} \cdot \frac{1}{5} = \frac{-1}{3125} \rightarrow 3125 = -5^{n-1}$$

see next problem for better steps

Note Exponent is negative and odd so through -1, n will be even

So Guess + ✓ from there

$$-5^{n-1} = -3125$$

$$n = 6$$

$$\sum_{n=1}^6 15 \left(-\frac{1}{5}\right)^{n-1}$$

56. $7 + 14 + 28 + \dots + 896$

$$\frac{14}{7} = 2 = r$$

$$a_n = a_1 \cdot r^{n-1}$$

$$896 = 7 \cdot 2^{n-1}$$

$$128 = 2^{n-1}$$

$$\log 128 = \log 2^{n-1}$$

$$\log 128 = n-1 \log 2$$

$$\frac{\log 128}{\log 2} = n-1$$

$$7 = n-1$$

$$8 = n$$

Sum of Infinite Geometric Sequence

$$r^{\infty}$$

$$r^1 + r^2 + r^3 + r^4 + \dots + r^{100} + \dots + r^{\infty}$$

$$r = \frac{1}{2} + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^4 + \dots + \left(\frac{1}{2}\right)^{100} + \dots + \left(\frac{1}{2}\right)^{\infty}$$

$$r^n \rightarrow 0$$

$$r < 1$$

$$a_1 \frac{(1-r^n)}{(1-r)} = a_1 \frac{(1-0)}{(1-r)} = \boxed{\frac{a_1}{1-r}}^{\text{infinite}}$$

If $|r| < 1$ and $n \rightarrow \infty$, then you have a definite limit - use $\frac{a_1}{1-r}$

ex) $1, .5, .25, .125, .0625, .03125$

$$r = \frac{1}{2} (< 1)$$

$$1 + .5 + .25 + .125 + .0625 + .03125$$

approaches limit of $2 = \frac{a_1}{1-r}$

If $|r| > 1$

ex) $r = 2 \leftarrow 1^2 + 2^2 + 3^2 + 4^2$
 $1 + 4 + 16 + 64 + 256$
 no limit

62. $\sum_{n=0}^{\infty} 2\left(-\frac{2}{3}\right)^n$ $\left|-\frac{2}{3}\right| < 1$

limit approaches $\frac{a_1}{1-r} = \frac{2}{1-(-\frac{2}{3})} = \frac{6}{5}$

64. $\sum_{n=1}^{\infty} \frac{1}{2}(4)^n$ $4 > 1$

no limit

2000

2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

2000 2000 2000 2000

8.3/Geometric Sequences + Series

HW

12/1

582

Geometric?

3. 6, 18, 30, 42, ...

$$\begin{array}{r} \swarrow \quad \swarrow \quad \swarrow \\ 18 \quad 30 \quad 42 \\ \underline{6} \quad \underline{18} \quad \underline{30} \\ 3 \quad \frac{5}{3} \quad \frac{7}{3} \end{array}$$

(X) Not Geometric

7. $\frac{1}{8}, \frac{1}{4}, \frac{1}{2}, 1$

$$\begin{array}{r} \swarrow \quad \swarrow \quad \swarrow \\ \frac{125}{125} \quad \frac{5}{25} \quad \frac{1}{5} \\ 2 \quad 2 \quad 2 \end{array}$$

(✓) Geometric

15. $r = -\frac{1}{18}$ $a_1 = 5$

$$a_2 = -\frac{1}{2}$$

$$a_3 = \frac{1}{28}$$

$$a_4 = -\frac{1}{280}$$

$$a_5 = \frac{1}{2800}$$

(✓)

21. $a_{n+1} = 2a_n$ $a_1 = 4$

$$a_2 = 18$$

$$a_3 = 36$$

$$a_4 = 72$$

$$a_5 = 144$$

23. $a_{n+1} = -\frac{3}{2}a_n$ $a_1 = 6$

$$a_2 = -9$$

$$a_3 = 13.5$$

$$a_4 = -20.25$$

$$a_5 = 30.375$$

27. $r = -\frac{1}{3}$

$a_1 = 6$

$a_{12} = ? = 6 \cdot \left(-\frac{1}{3}\right)^{12-1} = -0.00003387$

31.

~~Wrong~~
Variables

$a_2 = -18$

~~$a_3 = a_1 \cdot r$~~
 ~~$= 18 \cdot \left(-\frac{1}{3}\right)$~~

~~$= 27 = a_1 \cdot r^2$~~
 ~~$a_6 = -27 \cdot \left(\frac{2}{3}\right)^{5-1}$~~

$-\frac{2}{3^{10}}$

$a_5 = \frac{2}{3}$

$= a_2 \cdot r^{5-2}$

$\frac{2}{3} = -18 \cdot r^3$

$r = -\frac{1}{3}$

$a_6 = a_5 \cdot r$

$a_6 = \frac{2}{3} \cdot \frac{1}{3}$

$a_6 = -\frac{2}{9}$

35.

$a_1 = 5$
 $a_2 = 30 \rightarrow \frac{30}{5} = 6$

$a_3 = 180$

$a_{10} = ? = 5(6)^{10-1} = 50388480$

39.

$a_n = 2(1.3)^{n-1}$

n	1	2	3	4	5	6	7	8	9
$u(n)$	2	2.6	3.38	4.394	5.712	7.43	9.6	12.55	16.3

10 - 21.21

41.

$8, -4, 2, -1, \frac{1}{2}$

$s_1 = 8$

$s_2 = 8 + -4 = 4$

$s_3 = 8 - 4 + 2 = 6$

$s_4 = 8 - 4 + 2 - 1 = 5$

583

47.

$\sum_{i=1}^7 64\left(-\frac{1}{2}\right)^{i-1} = 64\left(\frac{1-(-\frac{1}{2})^7}{1-(-\frac{1}{2})}\right) = 64\left(\frac{128}{3/2}\right) = 43$

49. $\sum_{n=0}^{20} 3\left(\frac{3}{2}\right)^n = 3\left(\frac{1-\frac{3^{21}}{2}}{1-\frac{3}{2}}\right) = 3(9973.77) = 29921.31$

55. $5 + 15 + 45 + \dots + 3645$ Find sum

$3 = r$

$a_n = a_1 \cdot r^{n-1}$

$3645 = 5 \cdot 3^{n-1}$

$729 = 3^{n-1}$

$\log 729 = \log 3^{n-1}$

$\log 729 = (n-1) \log 3$

$\frac{\log 729}{\log 3} = n-1$

$6 = n-1$

$7 = n$

$5\left(\frac{1-3^7}{1-3}\right) = 5\left(\frac{-2186}{-2}\right) = 5(1093) = 5465$

57. $2^{-\frac{1}{2}} + \frac{1}{8} + \dots + \frac{1}{2048}$

$\frac{1}{2048} = 2 \cdot \left(-\frac{1}{4}\right)^{n-1}$

$\frac{1}{4096} = -\frac{1}{4}^{n-1}$

$\log(1/4096) = (n-1) \log(-1/4)$

$\frac{\log(1/4096)}{\log(-1/4)} = n-1$

$6 = n-1$

$n = 7$

$2\left(\frac{1-(-\frac{1}{4})^7}{1-(-\frac{1}{4})}\right) = 1.6$

65. $\sum_{n=0}^{\infty} (.4)^n = \frac{.4^0}{.4 < 1}$ bottom of sign

(✓)

$a_1 = (.4)^0 = 1$
 $\frac{a_1}{1-r} = \frac{1}{1-.4} = \frac{1}{.6} = \frac{5}{3}$

67. $\sum_{n=0}^{\infty} -3(.9)^n = \frac{-3}{1-.9} = \frac{-3}{.1} = -30$

(✓)

71. $3 - 1 + \frac{1}{3} - \frac{1}{9} \dots$
 $\downarrow$
 $-\frac{1}{3}$

(✓)

$\sum_{n=1}^{\infty} 3\left(-\frac{1}{3}\right)^n = \frac{3}{1-(-\frac{1}{3})} = \frac{3}{\frac{4}{3}} = 2.25 = \frac{9}{4}$

8.3 Geometric Sequences + Series Day 2

12/3

Annuitys

A deposit of \$50 is made at the start of each month to a savings account which pays 6% interest compounded monthly.

What is the balance at the end of 2 years?

- 24 months = 24 deposits -

$$\frac{6}{12} = .01 \quad \frac{.06}{12} = .005 \quad = \left[A = P \left(1 + \frac{r}{n} \right)^{nt} \right]$$

$$A_1 + A_2 + A_3 + \dots + A_{24}$$

↑ biggest deposit

$$A_{24} = 50 \left(1 + \frac{.06}{12} \right)^{24} = \$56.35$$

$$A_{23} = 50 \left(1 + \frac{.06}{12} \right)^{23} = \$56.077$$

Finite, geometric series

$$\text{multiplier} = 1 + \frac{.06}{12} = 1.005 = r$$

$$S_{24} = a_1 \cdot \left(\frac{1 - r^n}{1 - r} \right)$$

$$50 \left(1 + \frac{.06}{12} \right) = \frac{1 - 1.005^{24}}{1 - 1.005}$$

↑ last deposit - since only earns interest for 1 month

$$\$1277.96$$

583

80.

\$ 50 deposited in beginning of each month
 8% compounded monthly
 5 years
 $\rightarrow 5 \cdot 12 = 60$ months

$$A_1 = 50 \left(1 + \frac{.08}{12} \right)^1 = 50.33$$

$$A_{60} = 50 \left(1 + \frac{.08}{12} \right)^{60} = 74.49$$

$$S_{60} = 50.33 \left(\frac{1 + \frac{.08}{12}}{1 - \frac{.08}{12}} \right)^{60}$$

$r = 1 + \frac{.08}{12}$
 don't forget


$$\$ 3698.34 \quad \checkmark$$

Limits of Sequences

$$a_n = \frac{1}{n}$$

$$\frac{1}{1}, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5} \dots$$

Series -
 numbers
 you are
 adding



As n increases (w/o boundry) a_n gets closer
 closer to 0

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0 \quad \text{converges to } 0$$

ex2 $a_n = \frac{n+1}{n}$

n	1	2	3	4	...
a_n	2	$3/2$	$4/3$	$5/4$	
	2	1.5	1.33	1.25	

$$\lim_{n \rightarrow \infty} \frac{n+1}{n} = 1$$

← note that when degree(num) = degree(denom) → horiz asy, $\frac{+}{+} = 1$

ex3 $a_n = b_n = \frac{3n+2}{n+1}$

degrees =, so limit = $\frac{3}{1} = 3$

ex4 $b_n = \frac{n^2+5}{3n^2}$

$$\lim_{n \rightarrow \infty} = \frac{1}{3}$$

ex5 $a_n = 2^n$

$2^1; 2^2, 2^3, 2^4$
 $\lim_{n \rightarrow \infty} = \infty \rightarrow 2, 4, 9, 16, \dots$
diverges

ex6 $a_n = \sqrt{n}$

$\sqrt{1}, \sqrt{2}, \sqrt{3}, \sqrt{4}, \sqrt{5}$
 $1, 1.41, 1.73, 2, 2.2$
 $\lim_{n \rightarrow \infty} = \infty$
divergent

ex7

$$\frac{n^3+1}{2n^2-n}$$

n	1	2	3	4	5	...
a _n	2	1.5	1.8	2.3	2.8	...

$$\lim_{n \rightarrow \infty} = \infty \text{ divergent}$$

ex8 $b_n = (-1)^n$

n	1	2	3	4
b _n	-1	1	-1	1

No limit = divergent (even though bandy)

ex9 $a_n = \left(-\frac{1}{2}\right)^n$

n	1	2	3	4	5	...
a _n	-1/2	1/4	-1/8	1/16	-1/32	...

$$\lim_{n \rightarrow \infty} = 0 = \text{convergent to } 0$$

ex10

$$\sqrt{4 - \frac{1}{n}}$$

n	1	2	3	3	...
a _n	2.82	2.91	2.94	2.95	...

$$\lim_{n \rightarrow \infty} = 3$$

ex 11 $6 + \frac{4n-2}{n+8}$

n	1	2	3	4	5
a _n	6.22	6.6	6.9	7.1	7.3

$$\lim_{n \rightarrow \infty} \frac{4n-2}{n+8} = 4$$


$$6 + 4 = 10 = \lim_{n \rightarrow \infty} 6 + \frac{4n-2}{n+8}$$

ex 12 $\frac{5+\sqrt{2}}{n+1}$

$$\lim_{n \rightarrow \infty} = 0$$

ex 13 $b_n = 7$





$$\lim_{n \rightarrow \infty} 7 = 7$$



How are the notions of "getting larger and larger" and "getting closer and closer" measured? The notion of *arbitrarily close* provides a more precise definition of limit.

The limit of a sequence a_n is L if as n increases without bound, the terms of a_n get arbitrarily close to L . By "arbitrarily close" is meant simply "as close as you please." In other words, no matter how small a positive number ϵ (Greek epsilon) is chosen, for large enough N , all the terms after a_N are within ϵ of L . Stated even more precisely, a sequence a_n has a limit L if for any choice of $\epsilon > 0$, there exists a number N such that for all $n > N$, $|a_n - L| < \epsilon$.

For example, the limit of the harmonic sequence $\frac{1}{n}$ is 0. To illustrate this, let ϵ be a very small number, say $\epsilon = 0.000001$. For n large enough, in this case, if $n > N = 1,000,000$, then $|a_n - 0| = \left| \frac{1}{n} - 0 \right| = \frac{1}{n} < 0.000001$. No matter how small a number ϵ is chosen, n can always be chosen large enough to guarantee that $\left| \frac{1}{n} - 0 \right| < \epsilon$. As a further illustration of these ideas, the three sequences of Example 5 are graphed below.



Class Exercises

Find the limit of the indicated sequence as n increases without bound.

1. $a_n = \frac{1}{n^2}$ $\lim_{n \rightarrow \infty} = 0$

2. $b_n = \frac{3n+1}{n^2}$ $\lim_{n \rightarrow \infty} = 0$

3. $c_n = \sqrt{2} + \left(\frac{1}{2}\right)^n$ $\sqrt{2}$

Determine whether each sequence converges. If it converges, give the limit.

4. $d_n = \left(\frac{1}{4}\right)^n$

5. $t_n = 4^n$ Divergent

6. $m_k = (-3)^k$

7. $c_n = \frac{2n^2+4}{n^2+2}$ 2 Converges

8. $\frac{2n^3+2n+4}{n^2+1}$

9. $\frac{2}{n^2+1}$ 0

10. $a_n = \cos n\pi$

11. $b_n = (-0.1)^n$ Divergent

12. $c_n = 10^n$

13. **Thinking Critically** A certain infinite sequence has the property that terms with odd subscripts are positive and those with even subscripts are negative. The sequence converges. What must its limit be? Why?

14. Does the sequence of prime numbers seem to be convergent? Why?

15. Find the first ten terms of the sequence of the reciprocals of the prime numbers. Does this sequence seem to be convergent or divergent? Why?

Practice Exercises



Use appropriate technology.

Find the first six terms of each sequence. Tell whether the sequence is convergent or divergent. Justify your response using a graph.

1. $a_n = \frac{3}{n^3}$

2. $b_n = \sqrt{n}$

3. $c_n = -1$

Determine whether each sequence converges. If it converges, give the limit.

4. $d_n = 5^n$

5. $t_n = \frac{n-3}{n+3}$

6. $r_n = \frac{n^2 + n - 2}{n + 1}$

7. $a_n = (-1)^{n+1}\pi$

8. $b_k = \frac{2n+1}{3n+1}$

9. $c_t = \frac{2n^2}{n^2+1}$

10. $r_n = \sin \frac{1}{2}n\pi$

11. $d_n = \cos \frac{1}{2}n\pi$

12. $m_k = \left(-\frac{1}{4}\right)^k$

13. $a_n = \frac{3n^5 + n}{4n^5 + 8}$

14. $b_n = \frac{3n^5 + n}{4n^5 + 5}$

15. $c_n = \frac{3n^5 + n}{4n^5 + 8}$

16. Is the sequence $-1, 1, -1, \frac{1}{2}, -1, \frac{1}{4}, -1, \frac{1}{8}, \dots$ convergent or divergent? Why?

17. Graph $f(x) = \sin 2x$, when $0 \leq x < 2\pi$.

18. Is the Fibonacci sequence convergent or divergent? Why?

19. Find the sum of the first 15 terms of the sequence $c_n = 3n + 2$.

20. Let a_n be a constant sequence. Is this sequence convergent or divergent? Why?

21. Find the 56th term of the sequence $-31, -27, -23, -19, -15, \dots$

Determine whether each geometric sequence is convergent or divergent. If it is convergent, give the limit.

22. $a_n = 4(3^n)$

23. $b_n = 4(0.2)^n$

24. $c_n = 10\left(\frac{1}{2}\right)^n$

25. $r_k = \frac{1}{2}(10)^k$

26. $t_n = 6(-1)^n$

27. $d_n = 6\left(-\frac{1}{2}\right)^n$

28. Thinking Critically Let ar^n be a geometric sequence. What conditions on the common ratio r will make the sequence convergent? Divergent? Why?

29. Given triangle ABC with $c = 6$ ft, $b = 9$ ft, and $\angle C = 35^\circ$, solve the triangle.

Determine whether each arithmetic sequence is convergent or divergent.

30. $a_n = 2n + 1$

31. $b_n = 4 - 2n$

32. $c_n = -2.8$

33. Landscaping Describe the rectangular plot of maximum area that can be enclosed using 180 ft of fencing given that one side of the plot is along a building and needs no fencing.

34. Thinking Critically Let $a_n = a + (n - 1)d$ be an arithmetic sequence. What conditions on the common difference d will make the sequence convergent? Divergent? Why?

35. Find the first six terms of the sequence $a_n = e^n$. Is it convergent? Why?

36. Find the first six terms of the sequence $a_n = e^{-n}$. Is it convergent? Why?

8.3 Geometric Sequences + Series

12/3

583

77. \$1000 principal
3% interest
10 years
compounded:

annually (1) $\downarrow$ # per year

$$1000 \left(1 + \frac{.03}{1} \right)^{1 \cdot 10} \text{ years}$$

$$\$ 1343.92$$

semi-annually (2)

$$1000 \left(1 + \frac{.03}{2} \right)^{2 \cdot 10}$$

$$\$ 1346.86$$

Quarterly (4)

$$\$ 1348.35$$

Monthly (12)

$$\$ 1349.35$$

Daily (365)

$$\$ 1338.79$$

79. \$100/month
6% interest
5 years

$$A = 100 \left(1 + \frac{.06}{12} \right)^1 + \dots + 100 \left(1 + \frac{.06}{12} \right)^{60}$$

$$100 \left(1 + \frac{.06}{12} \right) \left(\frac{1 - \left(1 + \frac{.06}{12} \right)^{60}}{1 - 1 + \frac{.06}{12}} \right)$$

$$\$ 7011.88$$

584
89.

$$a_n = 3343(1.013)^n$$

$$n = 5, 6, \dots, 11$$

$$n = 1995$$

a) $r = 1.013$
 1.3%

b) $a_{20} = a_1 \cdot r^d$
 $a_{20} = 3343(1.013)^{20}$

4828.374 thousand
 $4,828,374$

c) $4100 = 3343(1.013)^n$

$$\begin{aligned} 1.126 &= 1.013^n \\ \log(1.126) &= n \log(1.013) \\ \frac{\log(1.126)}{\log(1.013)} &= n \end{aligned}$$

$15.8 = n$
 $\frac{8}{10}$ through 2005 August

91. Doubling Daily Wage $a_1 = .01$
 $r = 2$ (doubles)

a) $s_{20} = .01 \left(\frac{1-2^{20}}{1-2} \right)$
 $\$5,368,709.11$

b) $s_{30} = .01 \left(\frac{1-2^{30}}{1-2} \right)$
 $\$10,737,418.23$

c) $s_{31} = \$21,474,836.47$

8 extra Sequence + Series

Limits Day 2

12/4

Limits of Infinite Series

Find the sum of an infinite series

sum of an infinite series of numbers = limit of its associate sequence of partial sums so the number of terms increases without bounds

ex

Find the sum (if one exists) of the infinite series:

$$\sum_{k=1}^{\infty} \frac{1}{k(k+1)} \quad \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} + \dots$$

$\frac{4}{5}$

Partial Sums

$$\begin{aligned} s_1 &= \frac{1}{2} \\ s_2 &= \frac{1}{2} + \frac{1}{6} = \frac{2}{3} \\ s_3 &= \frac{1}{2} + \frac{1}{6} + \frac{1}{12} = \frac{3}{4} \\ s_4 &= \frac{1}{2} + \frac{1}{6} + \frac{1}{12} + \frac{1}{20} = \frac{4}{5} \end{aligned} \quad \left. \begin{array}{l} \\ \\ \\ \end{array} \right\} \frac{n}{n+1}$$

$$\lim_{n \rightarrow \infty} \frac{n}{n+1} = \frac{1}{1} = 1$$

ex2

$$\sum_{k=1}^{\infty} \frac{1}{k(k+1)(k+2)}$$

$$\begin{aligned} s_1 &= \frac{1}{6} \\ s_2 &= \frac{1}{6} + \frac{1}{24} = \frac{5}{24} \\ s_3 &= \frac{5}{24} + \frac{1}{60} = \frac{9}{40} \\ s_4 &= \frac{9}{40} + \frac{1}{120} = \frac{7}{30} \\ s_5 &= \frac{7}{30} + \frac{1}{210} = \frac{5}{21} \end{aligned} \quad \left. \begin{array}{l} \\ \\ \\ \\ \end{array} \right\} \frac{n(n+3)}{4(n+1)(n+2)}$$
$$\lim_{n \rightarrow \infty} = \frac{1}{4}$$

If the limit exists = convergent series

If the limit doesn't exist = divergent series

ex3

$$\sum_{k=1}^{\infty} 2k-1 \quad \text{arithmetic}$$

$$S_n = 1 + 3 + 5 + 7 + \dots$$

$$s_1 = 1$$

$$s_2 = 1 + 3 = 4$$

$$s_3 = 1 + 3 + 5 = 9$$

$$s_4 = 1 + 3 + 5 + 7 = 16$$

n^2

$\lim_{n \rightarrow \infty} n^2 = \text{divergent (no limit)}$

p4

$$\sum_{k=1}^{\infty} (-1)^k 10 = -10 + 20 - 40 + 80$$

divergent = no limit
(oscillates the limit)

$$r = -1$$

$$\|r\| = 1$$

$$\|r\| \geq 1$$

✓ divergent

p5

$$6 + 2 + \frac{2}{3} + \frac{2}{9} + \dots$$

$$\frac{1}{3} \quad \frac{1}{3}$$

$$r = 1/3$$

$$r < 1$$

$$\frac{a_1}{1-r} = \frac{6}{1-\frac{1}{3}} = 9$$

p6

$$1 + 2 + 4 + 8 + \dots$$

$$2 \quad 2 \quad 2$$

$$r = 2$$

$$r > 1$$

no limit

p7 $5 + .2 + .04 + .008 + .0016 + \dots$

look here

$$\frac{.04}{.02} = \frac{1}{5} = r$$

$$\frac{1}{5} < 1$$

Convergent by

$$5 \frac{.2}{1 - \frac{1}{5}} = 5 \frac{1}{4}$$

p8

$$10 + \frac{1}{9} + \frac{1}{9^2} + \frac{1}{9^3} + \dots$$

do separately $\rightarrow r = \frac{1}{9} = \frac{1}{9}$

$$\frac{1}{9} < 1$$

has limit

$$10 + \frac{1}{8} = 10 \frac{1}{8}$$

$$\frac{1}{8} = \frac{\frac{1}{9}}{1 - \frac{1}{9}}$$

ex9

A ball is dropped from a height of 24 m
Rebounds to a height of 75 %
Find the total distance it travels

← travels up & down

$$d = 24 + 2(.75)24 + 2(.75^2)24 + 2(.75^3)24 + \dots$$

↑ geometric series

↓ factor out 2

$$d = 24 + 2 \left(\frac{a_1}{1-r} \right)$$

$$24 + 2 \left(\frac{.75 \cdot 24}{1 - .75} \right) = 168 \text{ m}$$

P10

Dropped from 8ft

Bounces 70%

Find total distance it travels

$$d = 8 + 2(.7^1)8 + 2(.7^2)8 + 2(.7^3)8$$

$$8 + 2\left(\frac{.7 \cdot 8}{1 - .7}\right) = 45\frac{1}{3} \text{ ft}$$

Determine whether each series converges or diverges. If it converges, find the sum. If it diverges, tell why it diverges.

5. $\frac{1}{4} + \left(\frac{1}{4}\right)^2 + \left(\frac{1}{4}\right)^3 + \left(\frac{1}{4}\right)^4 + \dots$

6. $1 + 5 + 5^2 + 5^3 + \dots$

7. $\sum_{k=1}^{\infty} (0.15)^k$

8. $\sum_{i=1}^{\infty} 3\left(\frac{8}{9}\right)^i$

9. $\sum_{k=1}^{\infty} \left(-\frac{1}{4}\right)^k$

10. $\sum_{j=1}^{\infty} 3 \cdot \pi^j$

Express each repeating decimal as a ratio of two integers.

11. 0.2323232323 ... 12. 2.040404040 ... 13. 12.899999999 ... 14. -36.7777777 ...

Practice Exercises Use appropriate technology.

Find the first five terms of the sequence of partial sums for each series.

1. $3 + 6 + 9 + 12 + \dots$ *3/9/18/30/45*

2. $0.6 + (0.6)^2 + (0.6)^3 + \dots$ *7/10, 49/100, 343/1000, 216/625, 1296/3125*

3. $\sum_{k=1}^{\infty} \frac{1}{k^2}$ *1/1, 1/4, 1/9, 1/16, 1/25*

4. $\sum_{i=1}^{\infty} 5(0.9)^i$ *4.5, 4.05, 3.645, 3.2805, 2.95245*

5. Determine whether each series of Exercises 1 and 2 converges or diverges. If the series converges, find the sum. If it diverges, tell why it diverges.

Determine whether each series converges or diverges. If the series converges, find the sum. If it diverges, tell why it diverges.

6. $18 + 18(0.2) + 18(0.2)^2 + \dots$ *converges 22.5*

7. $1 + 0.1 + 0.001 + 0.0001 + \dots$ *converges 1.111...*

8. $\sum_{k=1}^{\infty} 8\left(\frac{2}{3}\right)^k$ *converges 5 1/3*

9. $\sum_{j=1}^{\infty} \left(\frac{-3}{11}\right)^j$ *converges -3/11*

10. $\sum_{k=1}^{\infty} -9(0.7)^k$ *converges -9*

11. $\sum_{i=1}^{\infty} (1.1)^i$ *divergent*

12. $\sum_{k=1}^{\infty} -3^k$ *divergent*

13. $\sum_{n=1}^{\infty} (-3)^n$ *divergent*

14. **Thinking Critically** Suppose that the terms of a sequence $a_1, a_2, a_3, \dots$ increase without bound. What conclusion can you draw about the related series $\sum_{n=1}^{\infty} a_n$? *divergent*

15. Solve the system of linear equations using an augmented matrix.

$$\begin{cases} x + 3y - z = 4 \\ 2x + 4y - 5z = 6 \\ 2y + 3z = 7 \end{cases}$$

Express each decimal as a ratio of two integers.

16. 0.45454545 ... 17. 12.999999 ... 18. 0.6758585858 ... 19. -8.9312424

20. **Thinking Critically** In the lesson, it is pointed out that $\lim_{n \rightarrow \infty} \frac{a(1-r^n)}{1-r} = \frac{a}{1-r}$ because r^n gets closer and closer to 0 as n increases without bound, providing $|r| < 1$. Discuss why the restriction $|r| < 1$ is necessary.

21. In the class exercises of Lesson 13.4 it was proved that $1 + 2 + 3 + \dots + n = \frac{1}{2}n(n+1)$. Use this fact to show that the series $\sum_{k=1}^{\infty} k$ is divergent.

22. Find the first five partial sums S_1, S_2, S_3, S_4 , and S_5 of the infinite series $\frac{1}{3} + \frac{1}{6} + \frac{1}{10} + \frac{1}{15} + \frac{1}{21} + \frac{1}{28} + \dots$. From the pattern, find a possible formula for S_n . Then use the definition of "infinite sum" to evaluate the sum of the infinite series. *Hint: Express S_2 and S_4 as unreduced fractions.*
- Handwritten notes: $S_1 = \frac{1}{3}$, $S_2 = \frac{2}{5}$, $S_3 = \frac{3}{7}$, $S_4 = \frac{4}{9}$, $S_5 = \frac{5}{11}$*

Find the first five partial sums S_1, S_2, S_3, S_4 , and S_5 of the following series.

$$\frac{3}{1 \cdot 4} + \frac{5}{4 \cdot 9} + \frac{7}{9 \cdot 16} + \frac{9}{16 \cdot 25} + \dots + \frac{2n+1}{n^2(n+1)^2} + \dots$$

Handwritten notes: $S_n = \frac{n}{(n+1)^2}$

23. From the pattern that you observe, what seems to be a formula for S_n ?
24. Use your formula for S_n and the definition of "infinite sum" to find the sum of the infinite series.

25. Use the principle of mathematical induction to prove that

$$\frac{1}{2 \cdot 5} + \frac{1}{5 \cdot 8} + \frac{1}{8 \cdot 11} + \dots + \frac{1}{(3n-1)(3n+2)} = \frac{n}{2(3n+2)}$$

Handwritten note: When degree numerator = degree denominator

26. Use the result of Exercise 25 to find the sum of the series $\sum_{k=1}^{\infty} \frac{1}{(3k-1)(3k+2)}$

27. Use the principle of mathematical induction to prove that

$$\frac{1}{1 \cdot 3} + \frac{1}{3 \cdot 5} + \frac{1}{5 \cdot 7} + \dots + \frac{1}{(2n-1)(2n+1)} = \frac{n}{2n+1}$$

28. Use the result of Exercise 27 to find the sum of the series $\sum_{k=1}^{\infty} \frac{1}{(2k-1)(2k+1)}$

29. **Physics** A ball is dropped from a height of 36 m. Each time it strikes the ground it rebounds to a height of 70% of the distance it fell. Find the total distance the ball travels after four bounces.
- Handwritten notes: $36 + 2(12.6 + 4.41 + 1.53 + 0.54) = 70.14$*

30. **Physics** A ball is dropped from a height of 40 m. Each time it strikes the ground it rebounds to a height of 45% of the distance it fell. Find the total distance the ball travels.
- Handwritten notes: $40 + 2(18 + 8.1 + 3.645 + 1.6455) = 75.45$*

31. **Economics** A federal tax cut adds \$10 billion to the income of taxpayers. Assume that each taxpayer as a result of this tax cut spends 90% of the money and the recipients of this money in turn spend 90% of the money. Find the total amount of spending that results from the initial tax cut.
- Handwritten notes: $10 + 9 + 8.1 + 7.29 + \dots = 100$ billion*

32. **Ecology** A radioactive substance is leaking from an underground storage tank. It spread 1000 m from the source during the first year, 800 m the second year, and 640 m the third year. If this pattern continues, how far will it spread in 6 years? Will the radioactive substance ever reach a river 6000 m from the source? Why or why not?
- Handwritten notes: $1000(1 - 0.8^6) = 3689.2$ m*

33. **Writing in Mathematics** Devise several situations that, like the wall-painting problem, can be described mathematically by the infinite series $\frac{1}{2} + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \dots$. Show how one of them is described by the series.

34. Identify an infinite geometric series that has a sum of 6 and a first term of 2.

35. Identify an infinite geometric series that has a sum of $\frac{8}{3}$ and a common ratio of $\frac{1}{4}$.

36. **Thinking Critically** Find an example of a geometric series with a negative sum that has a positive common ratio. Can an infinite geometric series with a negative sum have a positive first term? Why or why not?

Handwritten notes:

$6000 = 1000(1.9)^n$

$6 = 1.8^n$

$\frac{\log 6}{\log 1.8} = n$

$n \approx 384$ years

or $\frac{1000}{1-0.8} = \text{limit} = 5000$

8.5 Expanding Binomials

Notes

12/5

Binomial - polynomial w/ 2 terms

$$(x+y)^0 = 1$$


$$(x+y)^1 = x+y$$

$$(x+y)^2 = (x+y)(x+y) = x^2 + 2xy + y^2$$

$$(x+y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x+y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

$$(x+y)^5 = x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5$$



↑ terms must add to the degree

$$(x+y)^4 = \dots 6x^2y^2 \dots$$

$$\begin{matrix} 2+2 \\ 4 \end{matrix}$$

Binomial Theorem

$$(x+y)^n = x^n + nx^{n-1}y + \dots + nC_r x^{n-r}y^r + \dots + nx^n + y^n$$

Combinations

$$nC_r = \frac{n!}{(n-r)!r!}$$

used to get binomial coefficients

ex

$$8C_2 = \text{taking 8 objects - 2 at a time} = \binom{8}{2}$$

A B C D E F G H

AB BC ...

AC BD ...

AD BE ...

A... ..

$$= \frac{8!}{(8-2)!2!} = 28$$

599

$$6. \quad {}_{12}C_3 = \frac{12!}{(12-3)!(3)!} = \frac{12!}{9! \cdot 3!} = 220$$

14. ${}_{52}C_{47}$ on Calc:

$$\boxed{5} \boxed{2} \boxed{\text{Math}} \boxed{=}, \boxed{3} \boxed{4} \boxed{7} \boxed{\text{Enter}} = 2598960$$

prb nCr

$$12. \quad {}_{17}C_4 = \frac{17!}{13! \cdot 4!} = 2380$$

Pascal's Triangle

$(x+y)^0$	1	row 0
$(x+y)^1$	1 1	row 1
$(x+y)^2$	1 2 1	row 2
$(x+y)^3$	1 3 3 1	:
$(x+y)^4$	1 4 6 4 1	:
$(x+y)^5$	1 5 10 10 5 1	
$(x+y)^6$	1 6 15 20 15 6 1	

Add up the 2 numbers above it

$\begin{array}{cc} 1 & 2 \\ \swarrow & \searrow \\ & 3 \end{array}$

ex $(x+y)^7 = x^7 + 7x^6y + 21x^5y^2 + 35x^4y^3 + 35x^3y^4 + 21x^2y^5 + 7xy^6 + y^7$

$\downarrow C_1 \quad \downarrow C_2 \quad \downarrow C_3 \quad \downarrow C_4$

26. $(y-2)^5 \quad n=5 \rightarrow {}_5C_r$

$$1y^5 + 5y^4(-2) + 10y^3(-2)^2 + 10y^2(-2)^3 + 5y(-2)^4 + (-2)^5$$

$$y^5 - 10y^4 + 40y^3 - 80y^2 + 80y - 32$$

$$(4x)^6 = 4^6 x^6$$

$$4096 x^6$$


$$30. (4x + 3y)^4 = {}_4C_0 (4x)^4 + {}_4C_1 (4x)^3 (3y) + {}_4C_2 (4x)^2 (3y)^2 + {}_4C_3 (4x) (3y)^3 + {}_4C_4 (3y)^4$$

$$4 \cdot 4^4 x^4 + 4 \cdot 4^3 \cdot x^3 \cdot 3 \cdot y + 6 \cdot 4^2 \cdot x^2 \cdot 3^2 \cdot y^2 + 4 \cdot 4 \cdot x \cdot 3^3 \cdot y^3 + 1 \cdot 3^4 \cdot y^4$$

$$256x^4 + 4 \cdot 64x^3 \cdot 3y + 6 \cdot 16x^2 \cdot 9y^2 + 4 \cdot 4x \cdot 27y^3 + 81y^4$$

$$256x^4 + 768x^3y + 864x^2y^2 + 432xy^3 + 81y^4$$

$$256x^4 + 768x^3y + 864x^2y^2 + 432xy^3 + 81y^4$$



8.5 Binomial Theorem

Day 2 Notes

12/8

Review

$$x^n + n x^{n-1} y + \dots + n C_r x^{n-r} y^r + \dots + n \cdot x y^{n-1} + y^n$$

599

ex: $(x+y)^5 = x^5 + 5x^4y + 5C_2 x^3y^2 + 5C_3 x^2y^3 + 5C_4 xy^4 + y^5$

32. $(2x-y)^5 = (2x)^5 + 5(2x)^4(-y) + 10(2x)^3(-y)^2 + 10(2x)^2(-y)^3 + 5(2x)(-y)^4 + (-y)^5$

$$32x^5 + 5 \cdot 16x^4(-y) + 10 \cdot 8x^3y^2 + 10 \cdot 4x^2(-y)^3 + 5 \cdot 2xy^4 + (-y)^5$$

$$32x^5 - 80x^4y + 80x^3y^2 - 40x^2y^3 + 10xy^4 - y^5$$

38. $\left(\frac{1}{x} + 2y\right)^6 = \left(\frac{1}{x}\right)^6 + 6\left(\frac{1}{x}\right)^5(2y) + 15\left(\frac{1}{x}\right)^4(2y)^2 + 20\left(\frac{1}{x}\right)^3(2y)^3 + 15\left(\frac{1}{x}\right)^2(2y)^4 + 6\left(\frac{1}{x}\right)(2y)^5 + (2y)^6$

$$\frac{1}{x^6} + 6 \frac{1}{x^5} 2y + 15 \frac{1}{x^4} 4y^2 + 20 \frac{1}{x^3} 8y^3 +$$

$$15 \frac{1}{x^2} 16y^4 + 6 \frac{1}{x} 32y^5 + 64y^6$$

$$\frac{1}{x^6} + \frac{12y}{x^5} + \frac{60y^2}{x^4} + \frac{160y^3}{x^3} +$$

$$\frac{240y^4}{x^2} + \frac{192y^5}{x} + 64y^6$$

can put
in numerator too

$$\frac{12y}{x^5}$$

ex2 Find the 6th term of $(a+2b)^8$
 $\leftarrow$ 1 minus what you are asked for
 ${}^8C_5 a^3 (2b)^5$
 $56 a^3 32 b^5$
 $1792 a^3 b^5$

52. $(5a+6b)^5$
 $n=5$
 ${}^5C_4 (5a)^1 (6b)^4$
 $5 \cdot 5a \cdot 1296 b^4$
 $32400 a b^4$

ex3 Find the coefficient of the term $a^6 b^5$
in the expression $(2a-5b)^{11}$
degree = 11
 $r = 5$
 ${}^{11}C_5 (2a)^6 (-5b)^5$
 $462 \cdot 64 a^6 - 3125 b^5$
 $-92,400,000 a^6 b^5$

56. Binomial $(x+4)^{12}$ Find coefficient of
Term = $a x^5$
 $\leftarrow$ 1st term
 ${}^{12}C_5 x^5 \cdot 4^7$
 $792 \cdot x^5 \cdot 16384$
 $12,976,128 x^5$
 $\hookrightarrow 12,976,128$

58. $(4x-y)^{10}$
 $a x^2 y^8$
coefficient
 ${}^{10}C_2 (4x)^2 (-y)^8$
 $45 \cdot 16 \cdot x^2 y^8$
 $720 x^2 y^8$
 $\hookrightarrow 720$

$$62. (z^2 - 1)^{12}$$

$$a_{2^6}$$


Find a

what shows up

$${}^{12}C_9 \left((z^2)^3 (-1)^9 \right)$$

$$220 z^6 - 1$$

$$-220 z^6$$



8.5 Binomial Theorem

Day 2 thw

12/7

599

Find the binomial coefficient

$$3. \binom{12}{0} = 12C_0 = 1$$

$$9. \binom{100}{48} = 100C_{48} = 4950$$

$$13. 22C_9 = 497420$$

$$17. 7C_4 = 35$$

$$27. (x+y)^5 = 1x^5 + 5C_1 x^4 y + 5C_2 x^3 y^2 + 5C_3 x^2 y^3 + 5C_4 x y^4 + y^5$$

$$x^5 + 5x^4 y + 10x^3 y^2 + 10x^2 y^3 + 5x y^4 + y^5$$

④

$$34. (5-2y)^3 = 5^3 + 3C_1 5^2(-2y) + 3C_2 5(-2y)^2 + (-2y)^3$$

$$125 + 3 \cdot 25 \cdot -2y + 3 \cdot 5 \cdot 4y^2 + -8y^3$$

$$125 - 150y + 60y^2 - 8y^3$$

X

$$35. (x^2+y^2)^4 = (x^2)^4 + 4(x^2)^3 y^2 + 4C_2 (x^2)^2 (y^2)^2 + 4C_3 (x^2)(y^2)^3 + 1(y^2)^4$$

$$x^8 + 4x^6 y^2 + 6x^4 y^4 + 4x^2 y^6 + y^8$$

$$41. -3(x-2)^3 - 4(x+1)^6$$

$$-3(x^3 + 3x^2(-2) + 3C_2 x + (-2)^2 + 1(-2)^3)$$

$$-3(x^3 - 6x^2 + 12x - 8)$$

$$-3x^3 + 18x^2 - 36x + 24$$

$$-4(x+1)^6$$

$$-4(x^6 + 6x^5 + 15x^4 + 20x^3 + 15x^2 + 6x + 1)$$

$$-4x^6 - 24x^5 - 60x^4 - 80x^3 - 60x^2 - 24x - 4$$

$$-4x^6 - 24x^5 - 60x^4 - 83x^3 - 42x^2 - 60x + 20$$

④

$$43. (3t - s)^3 = (3t)^3 + 3(3t)^2(-s) + 3(3t)(-s)^2 + (-s)^3$$

$$9t^3 - 9t^2s + 9ts - s^3$$

$$45. (x + 2y)^5 = x^5 + 5x^4(2y) + 10x^3(2y)^2 + 10x^2(2y)^3 + 5x(2y)^4 + (2y)^5$$

$$x^5 + 10x^4y + 40x^3y^2 + 80x^2y^3 + 80xy^4 + 32y^5$$

$$49. (x - 6y)^5$$

$$n=3 \quad {}^5C_0 x^5 (-6y)^0$$

$$10x^4 \cdot 36y^2$$

$$360x^3y^2$$

$$53. (10x - 3y)^{12}$$

$$n=9 \quad {}^{12}C_8 (10x)^4 (-3y)^8$$

$$495 \cdot 10000 \cdot x^4 \cdot 6561 y^8$$

$$3,24 \cdot 10^{10} \cdot x^4 y^8$$

$$59. (3x - 2y)^9$$

$$a x^6 y^3 \quad {}^{10}C_6 (3x)^6 (-2y)^3$$

$$210 \cdot 729 x^6 \cdot -8 y^3$$

$$-1224720 x^6 y^3$$

$$62. (z^2 - 1)^{12}$$

$$a z^6 \quad {}^{12}C_6 (z^2)^6 (-1)^6$$

could be wrong

$$63. (\sqrt{x} + 5)^4 = (\sqrt{x})^4 + 4(\sqrt{x})^3 5 + 6(\sqrt{x})^2 5^2 + 4(\sqrt{x}) 5^3 + 5^4$$

$$x^2 + 4x^{3/2} 5 + 150x + 500x^{1/2} + 625$$

$$64. (u^{3/5} + v^{1/5})^5 = (u^{3/5})^5 + 5(u^{3/5})^4 (v^{1/5}) + 10(u^{3/5})^3 (v^{1/5})^2 + 10(u^{3/5})^2 (v^{1/5})^3 + 5(u^{3/5}) (v^{1/5})^4 + (v^{1/5})^5$$

$$u^3 + 5u^{12/5} + 10u^{9/5} v^{2/5} + 10u^{6/5} v^{3/5} + 5u^{3/5} v^{4/5} + v$$

$$(u^{3/5})^2 = u^{(3/5) \cdot 2} = u^{6/5}$$

multiply

67. $f(x) = x^3$ $\frac{f(x+h) - f(x)}{h}$

$\frac{(x+h)^3 - x^3}{h}$

$\frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h}$

$\frac{3x^2h + 3xh^2 + h^3}{h}$

$(3x^2 + 3h + h^2)$

71. $(1+i)^4 = 1^4 + 4 \cdot 1^3 \cdot i + 6 \cdot 1^2 \cdot i^2 + 4 \cdot 1 \cdot i^3 + i^4$
 $1 + 4i + 6(-1) + 4(-i) + 1$
 $1 + 4i - 6 - 4i + 1$
 -4

85. Coin Toss

7 times

4 heads $\approx {}^7C_4 \left(\frac{1}{2}\right)^4 \left(\frac{1}{2}\right)^3 \rightarrow \left(\frac{1}{2} + \frac{1}{2}\right)^7$
 $35 \left(\frac{1}{16}\right) \left(\frac{1}{8}\right)$
 $1 \ 2 \ 7 \ 3 \ 4 \ 3 \ 7 \ 5$

27.34%

$$f(x) = \ln(x)$$

$$f'(x) = \frac{1}{x}$$

$$f''(x) = -\frac{1}{x^2}$$

$$f'''(x) = \frac{2}{x^3}$$

$$f^{(4)}(x) = -\frac{6}{x^4}$$

$$f^{(5)}(x) = \frac{24}{x^5}$$

$$f^{(6)}(x) = -\frac{120}{x^6}$$

$$f^{(7)}(x) = \frac{720}{x^7}$$

$$f^{(8)}(x) = -\frac{5040}{x^8}$$

$$f^{(9)}(x) = \frac{40320}{x^9}$$

$$f^{(10)}(x) = -\frac{362880}{x^{10}}$$

$$f^{(11)}(x) = \frac{3981312}{x^{11}}$$

$$f^{(12)}(x) = -\frac{4778240}{x^{12}}$$

$$f^{(13)}(x) = \frac{62918400}{x^{13}}$$

8/ Sequences Formula Sheet

12/10

Arithmetic Sequence

$$a_n = \underset{\substack{\uparrow \\ \text{slope}}}{d}n + c$$

$d = \text{common diff}$
 slope

$$c = a_1 - d$$

$$a_n = a_1 + (n-1)d \quad \leftarrow \text{recursive version}$$

Sum of Finite Arithmetic Series

$$S_n = \frac{n}{2}(a_1 + a_n) \quad \text{partial sum}$$

next term =
 $a_{n+1} = a_n + d$

Geometric Sequences

$$\frac{a_2}{a_1} = \frac{a_3}{a_2} = \frac{a_4}{a_3} = r = \text{common ratio}$$

$$a_n = a_1 r^{n-1} \quad \leftarrow \text{recursive} \quad \text{) same thing}$$

$$a_n = \left(a_1, r\right) \left(\frac{1}{r}\right)^n$$

Sum of a Finite Geometric Series

$$S_n = a_1 \left(\frac{1-r^n}{1-r} \right)$$

Sum of an Infinite Geometric Series

$$S = \frac{a_1}{1-r} \quad \begin{array}{l} |r| < 1 \rightarrow \text{convergent} \\ |r| \geq 1 \rightarrow \text{divergent} \end{array}$$

13/11

1. The first part of the question is to find the value of x such that $f(x) = 0$.

2. The second part is to find the value of x such that $f(x) = 1$.

3. The third part is to find the value of x such that $f(x) = 2$.

4. The fourth part is to find the value of x such that $f(x) = 3$.

5. The fifth part is to find the value of x such that $f(x) = 4$.

6. The sixth part is to find the value of x such that $f(x) = 5$.

7. The seventh part is to find the value of x such that $f(x) = 6$.

8. The eighth part is to find the value of x such that $f(x) = 7$.

9. The ninth part is to find the value of x such that $f(x) = 8$.

10. The tenth part is to find the value of x such that $f(x) = 9$.

11. The eleventh part is to find the value of x such that $f(x) = 10$.

12. The twelfth part is to find the value of x such that $f(x) = 11$.

13. The thirteenth part is to find the value of x such that $f(x) = 12$.

14. The fourteenth part is to find the value of x such that $f(x) = 13$.

15. The fifteenth part is to find the value of x such that $f(x) = 14$.

16. The sixteenth part is to find the value of x such that $f(x) = 15$.

17. The seventeenth part is to find the value of x such that $f(x) = 16$.

18. The eighteenth part is to find the value of x such that $f(x) = 17$.

19. The nineteenth part is to find the value of x such that $f(x) = 18$.

20. The twentieth part is to find the value of x such that $f(x) = 19$.

21. The twenty-first part is to find the value of x such that $f(x) = 20$.

22. The twenty-second part is to find the value of x such that $f(x) = 21$.

23. The twenty-third part is to find the value of x such that $f(x) = 22$.

24. The twenty-fourth part is to find the value of x such that $f(x) = 23$.

25. The twenty-fifth part is to find the value of x such that $f(x) = 24$.

26. The twenty-sixth part is to find the value of x such that $f(x) = 25$.

27. The twenty-seventh part is to find the value of x such that $f(x) = 26$.

28. The twenty-eighth part is to find the value of x such that $f(x) = 27$.

29. The twenty-ninth part is to find the value of x such that $f(x) = 28$.

30. The thirtieth part is to find the value of x such that $f(x) = 29$.

Sequences + Series

Review Hw

12/10

626

3. Write the first 5 terms

$$a_n = \frac{72}{n!}$$

n	1	2	3	4	5
a_n	72	36	12	3	$\frac{1}{6}$

$$11. \frac{2! \cdot 5!}{6!} = \frac{2 \cdot 1 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = \frac{2 \cdot 1}{6} = \frac{2}{6} = \frac{1}{3}$$

$$15. \sum_{j=1}^4 6/j^2 = 8.54 \quad \leftarrow \text{calc } \text{sum}(\text{seq}(\text{thing}, x, \overset{\text{start}}{1}, \text{limit}))$$

$$17. \sum_{k=1}^{10} 2k^3 = 770$$

$$23. \frac{1}{2} + \frac{2}{3} + \frac{3}{4} + \dots + \frac{9}{10} \quad \sum_{x=1}^9 \frac{x}{x+1} = 7.07$$

$$27. \sum_{k=1}^{\infty} 2\left(\frac{1}{100}\right)^k \quad \begin{array}{l} \text{a) Find 4th term: } 2\left(\frac{1}{100}\right)^4 = 2 \cdot 10^{-8} \\ \text{b) Sum} = r = \frac{1}{100} < 1 \text{ (convergent)} \end{array}$$

$$\frac{\frac{2}{100}}{1 - \left(\frac{1}{100}\right)} = .0202 = \frac{2}{99}$$

29. Compound Interest

$n = \text{quarters}$

$$a_n = 2500 \left(1 + \frac{.08}{4}\right)^n$$

quarter	1	2	3	4	5	6
total	2550	2601	2653	2706.1	2760.2	2815.4
	7	8		40		
	2871.7	2929.1		5520.10		

37. $a_4 = 10$
 $a_{10} = 28$ $\frac{28-10}{10-4} = \frac{18}{6} = 3$ $a_n = 3n + c$

n	a_n
1	1
2	4
3	7
4	10
5	13

$a_1 = a_4 - 3d$
 $10 - 3 \cdot 3$
 $a_1 = 1$

39. $a_{k+1} = a_k - 3$
 $a_1 = 35$
 $a_2 = 32$
 $a_3 = 29$
 $a_4 = 26$
 $a_5 = 23$

Q27

43. Sum of first 20 terms

$a_1 = 100$
 $d = -3$

$s_n = \frac{n}{2} (a_1 + a_{100})$ $a_{100} = 100 + 99(-3)$

$s_{20} = \frac{20}{2} (100 + -197)$

$s_{20} = -970$

47. $\sum_{k=1}^{11} \left(\frac{2}{3}k + 4 \right) =$ $a_1 = \frac{2}{3} + 4 = 4\frac{2}{3}$
 $a_{11} = \frac{2}{3} \cdot 11 + 4 = 11\frac{1}{3}$

$s_{11} = \frac{11}{2} \left(4\frac{2}{3} + 11\frac{1}{3} \right)$
 88

49. First 100 positive multiples of 5

(1)

$\sum_{k=1}^{100} 5k$ $a_1 = 5$
 $a_{100} = 500$

$s_{100} = \frac{100}{2} (5 + 500)$
 25250

50.

$$\sum_{k=20}^{80} k$$

$$a_{20} = 20$$

$$a_{80} = 80$$

$$S_{61} = \frac{61}{2} (20 + 80)$$

$$3,000$$

$$3,058$$

51.

Salary = \$34,000 (year 1)

Raise = \$2250/year for 5 years

$$\text{Year 5} = 34,000 + 2250 \cdot 4$$

$$\$43,000$$

$$5 \text{ year sum} = \frac{5}{2} (34,000 + 43,000)$$

$$\$192,500$$

52.

1st trip = 123 bales

Each trip = 11 fewer bales

total after 8 trips

trip + trip + 6 more trips

2 additional

$$a_1 = 123$$

$$a_8 = 123 - 11 \cdot 7$$

$$46$$

$$S_8 = \frac{8}{2} (123 + 46)$$

$$46$$

$$S_8 = 810$$

$$676$$

Geometric

59.

$$a_1 = 9$$

$$a_3 = 4$$

$$a_3 = a_1 \cdot r^{3-1}$$

$$4 = 9 \cdot r^2$$

$$\frac{4}{9} = r^2$$

$$r = \frac{2}{3}$$

$$a_1 = 9$$

$$a_2 = 6$$

$$a_3 = 4$$

$$a_4 = 2\frac{2}{3}$$

$$a_5 = 1\frac{6}{9}$$

63. $a_1 = 25$
 $a_{k+1} = -\frac{3}{5} a_k$

$a_1 = 25$
 $a_2 = -15$
 $a_3 = 9$
 $a_4 = -5.4$
 $a_5 = 3.24$

65. Find n th + sum of first 20 terms

$a_1 = 16$
 $a_2 = -8$

$r = \frac{-8}{16} = -\frac{1}{2}$

①

$a_n = 16 \cdot \left(-\frac{1}{2}\right)^{n-1}$

← memorize $a_1 \cdot r^{n-1}$

~~$|r| < 1$ convergent~~
 not infinite

$\frac{16}{1 - (-\frac{1}{2})} = 10\frac{2}{3}$

$\uparrow 16 \frac{(1 - (-\frac{1}{2})^{20})}{1 - (-\frac{1}{2})}$

75. $\sum_{i=1}^{10} 10 \left(\frac{3}{5}\right)^{i-1}$

$a_1 = 10$
 $a_{10} = 1.10077696$

$S_n = \frac{9}{2} (10 + 1.10077696)$
 45.45

79.

$\sum_{k=1}^{\infty} 4 \left(\frac{2}{3}\right)^{k-1}$

$a_1 = 4$
 $\frac{2}{3} < 1$ convergent

$\frac{4}{1 - \frac{2}{3}} = 12$

81. Depreciation
\$120,000
30% year depreciation

a) $\$120,000 (.7)^t$

b) $t=5 \rightarrow 120,000 (.7)^5$
\$26,168.14

82. Annuity
\$75/month
4% interest

finite, geometric sequence

$$A = 75 \left(1 + \frac{.04}{12}\right)^1 + \dots + 75 \left(1 + \frac{.04}{12}\right)^{48}$$

$a_{48} = 75.25$ $a_1 = 75.50$ $= \$87.99$

It was already reversed

$$r = \frac{75.50}{75.25} = 1.003333$$

$$a_1 = 75.25 \left(\frac{1 - 1.00333^{48}}{1 - 1.00333} \right)$$

geometric

~~\$4571.93~~ **\$3909.96**

628

93. $a_n = a_{n-1} - 1$
 $a_1 = 16$

$a_1 = 16$
 $a_2 = 15$
 $a_3 = 14$
 $a_4 = 13$
 $a_5 = 12$

linear

+ study worksheets

$$97. \binom{9}{4} = {}^9C_4 = 126$$

$$101. \binom{8}{4} = {}^8C_4 = 70$$

$$105. (a-4b)^5 = a^5 + 5a^4(-4b) + 10a^3(-4b)^2 + 10a^2(-4b)^3 + 5a(-4b)^4 + (-4b)^5$$

$$a^5 + 5a^4 \cdot -4 \cdot b + 10a^3 \cdot 16 \cdot b^2 + 10a^2 \cdot -64b^3 + 5a \cdot 256b^4 - 1024b^5$$

$$a^5 - 20a^4b + 160a^3b^2 - 640a^2b^3 + 1280ab^4 - 1024b^5$$

$$107. (7+2i)^4 = 7^4 + 4 \cdot 7^3(2i) + 6 \cdot 7^2(2i)^2 + 4 \cdot 7 \cdot (2i)^3 + (2i)^4$$

$$2401 + 2744i + 1176i^2 + 224i^3 + 16i^4$$

$$2401 + 2744i - 1176 - 224i + 16$$

$$1241 + 2520i$$

(✓)

Sequences + Series

Review 2

12/11

1. Find formula
arithmetic
 $a_5 = 24$
 $a_9 = 40$

$$\frac{40 - 24}{9 - 5} = \frac{16}{4} = 4$$

$$a_0 = 24 - 4 \cdot 5$$
$$a_0 = 4$$

$$a_n = 4n + 4$$

2. Find missing 3 terms
geometric
 $a_1 = 4$
 $a_5 = 324$

$$324 = 4 \cdot r^4$$
$$81 = r^4$$
$$\sqrt[4]{81} = \sqrt[4]{4^4}$$
$$r = 3$$

$$a_1 = 4$$
$$a_2 = 12$$
$$a_3 = 36$$
$$a_4 = 108$$
$$a_5 = 324$$

3. A firm offers \$26,000 salary
\$900/year raise
What is salary in 9th year?

$$a_9 = 26,000 + 900(9-1)$$
$$\$33,200$$

4. Show w/ sigma: $3 - 6 + 9 - 12 + 15$

$$\sum_{x=1}^5 3n(-1)^{n+1}$$

memorize rotating negative

5. Find sum of first 25 terms

$$3\text{rd term} = 24$$

$$\text{diff} = 3.5$$

$$a_1 = 24 - 2(3.5)$$

$$a_1 = 17$$

$$a_{25} = 17 + \overset{24}{\cancel{25}}(3.5)$$

$$\cancel{104.5} \quad 101$$

differences

$$\frac{25}{2} (17 + \overset{101}{\cancel{104.5}})$$

$$\cancel{1518.75} \quad 1475$$

6. Find a_5

$$r = 1.75$$

$$S_5 = 20,775$$

$$S_5 = a_1 \left(\frac{1 - 1.75^5}{1 - 1.75} \right)$$

$$20775 = a_1 \cdot 20.55$$

$$a_1 = 1010.91$$

$$a_5 = a_1 r^4$$

$$a_5 = 1010.91 \cdot 1.75^4$$

$$9481.23$$

7. Bouncing Ball

dropped 6 ft

rebounds $\frac{1}{3}$ of previous bounce

total distance?

$$6 + 2(.33)6 + 2(.33^2)6 + 2(.33^3)6 + 2(.33^4)6 \dots$$

$$6 + 2 \left(\frac{a_1}{1-r} \right) \rightarrow 6 + 2 \left(\frac{6 \cdot .33}{1-.33} \right) \rightarrow 11.94 \text{ ft}$$

8. A architect makes a theater with 15 seats in the first row
18 seats = 2nd
21 seats = 3rd
seating capacity = 870
How many rows 870 is the.

~~$870 = 12 + 3d$~~ $\rightarrow$ $870 = \frac{n}{2}(a_1 + a_n)$ ^{Sum}

~~$858 = 3d$~~

~~$d = 286$ rows~~

$870 = \frac{n}{2}(15 + a_n)$

$870 = \frac{n}{2}(15 + [12 + 3n])$

$870 = \frac{n}{2}(27 + 3n)$

$1740 = n(27 + 3n)$

$n^2 + 9n - 580$

$\rightarrow$

$1740 = 27n + 3n^2$

$3n^2 + 27n - 1740 = 0$

rooten

$n = 20$

$(n - 20)(n + 29)$

$$(1.0 \times 10^{-3}) / 10 = 1.0 \times 10^{-4}$$

$$(1.0 \times 10^{-4}) / 10 = 1.0 \times 10^{-5}$$

$$(1.0 \times 10^{-5}) / 10 = 1.0 \times 10^{-6}$$

$$(1.0 \times 10^{-6}) / 10 = 1.0 \times 10^{-7}$$

$$(1.0 \times 10^{-7}) / 10 = 1.0 \times 10^{-8}$$

$$(1.0 \times 10^{-8}) / 10 = 1.0 \times 10^{-9}$$

$$(1.0 \times 10^{-9}) / 10 = 1.0 \times 10^{-10}$$

$$(1.0 \times 10^{-10}) / 10 = 1.0 \times 10^{-11}$$

$$(1.0 \times 10^{-11}) / 10 = 1.0 \times 10^{-12}$$

$$(1.0 \times 10^{-12}) / 10 = 1.0 \times 10^{-13}$$

$$(1.0 \times 10^{-13}) / 10 = 1.0 \times 10^{-14}$$

8 Sequences + Series

Test Objectives

12/11

Write first 5 terms

Use n/r

Find sums of infinite series

Factorial notation

Finding n th term

Find formula for n th term


Use sigma to write the sum

Binomial Theorem

- expanding

- getting specific term

- limit of sequences + series (worksheet)



$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$1 + \cot^2 x = \csc^2 x$$

Cofunction Identities

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x \quad \cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$\csc\left(\frac{\pi}{2} - x\right) = \sec x \quad \tan\left(\frac{\pi}{2} - x\right) = \cot x$$

$$\sec\left(\frac{\pi}{2} - x\right) = \csc x \quad \cot\left(\frac{\pi}{2} - x\right) = \tan x$$

Negative Angle Identities

$$\sin(-x) = -\sin x \quad \cos(-x) = \cos x$$

$$\csc(-x) = -\csc x \quad \tan(-x) = -\tan x$$

$$\sec(-x) = \sec x \quad \cot(-x) = -\cot x$$

Sum and Difference Formulas

$$\sin(u \pm v) = \sin u \cos v \pm \cos u \sin v$$

$$\cos(u \pm v) = \cos u \cos v \mp \sin u \sin v$$

$$\tan(u \pm v) = \frac{\tan u \pm \tan v}{1 \mp \tan u \tan v}$$

Double Angle Formulas

$$\sin 2x = 2 \sin x \cos x$$

$$\cos 2x = \cos^2 x - \sin^2 x = 2 \cos^2 x - 1 = 1 - 2 \sin^2 x$$

$$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$$

Additional Trigonometric Formulas/Identities

Half - Angle Identities:

$$\sin^2 u = \frac{1 - \cos 2u}{2}$$

$$\cos^2 u = \frac{1 + \cos 2u}{2}$$

$$\tan^2 u = \frac{1 - \cos 2u}{1 + \cos 2u}$$

Half - Angle Formulas:

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$$

Alternative Half - Angle Formulas for Tangent:

$$\tan \frac{\theta}{2} = \frac{1 - \cos \theta}{\sin \theta}$$

$$\tan \frac{\theta}{2} = \frac{\sin \theta}{1 + \cos \theta}$$

Product - to - Sum Formulas:

$$\sin u \cos v = \frac{1}{2} [\sin(u + v) + \sin(u - v)]$$

$$\cos u \sin v = \frac{1}{2} [\sin(u + v) - \sin(u - v)]$$

$$\cos u \cos v = \frac{1}{2} [\cos(u + v) + \cos(u - v)]$$

$$\sin u \sin v = \frac{1}{2} [\cos(u - v) - \cos(u + v)]$$

Sum - to - Product Formulas:

$$\sin u + \sin v = 2 \sin \left(\frac{u + v}{2} \right) \cos \left(\frac{u - v}{2} \right)$$

$$\sin u - \sin v = 2 \cos \left(\frac{u + v}{2} \right) \sin \left(\frac{u - v}{2} \right)$$

$$\cos u + \cos v = 2 \cos \left(\frac{u + v}{2} \right) \cos \left(\frac{u - v}{2} \right)$$


$$\cos u - \cos v = -2 \sin \left(\frac{u + v}{2} \right) \sin \left(\frac{u - v}{2} \right)$$

Name Michael Plasker

Unit Circle in terms of Degrees & Radians

(in multiples of 30° , 45° , 60°)

Directions: Convert the radian measure to degrees and
Convert the given degree measure to radians.



5/ISS3

More Practice + Review


12/13

1. $\csc(-x) = -5$

$-\csc x = -5$

$\csc x = 5 = \frac{1}{\sin x} = 5$

$= \frac{1}{\sin x} = 5$



$5^2 = 1^2 + x^2$

$x = \sqrt{24}$

$\sin = \frac{y}{r} = \left(\frac{1}{5}\right)$

$\csc = (5)$

$\cos = \frac{x}{r} = \left(\frac{\sqrt{24}}{5}\right)$

$\sec = \frac{r}{x} = \frac{5}{\sqrt{24}} \rightarrow \frac{5\sqrt{24}}{24} = \frac{5 \cdot 2\sqrt{6}}{24}$

$\tan = \frac{y}{x} = \frac{1}{\sqrt{24}}$

$\cot = \frac{x}{y} = \frac{\sqrt{24}}{1} \rightarrow \left(\sqrt{24}\right) \left(\frac{1}{5\sqrt{6}}\right)$

$\rightarrow \frac{\sqrt{24}}{24} \rightarrow \frac{\sqrt{4}\sqrt{6}}{24} = \frac{2\sqrt{6}}{24} = \left(\frac{\sqrt{6}}{12}\right)$

2. $\frac{\sin \theta + \cos \theta}{\sin \theta} - \frac{\cos \theta - \sin \theta}{\cos \theta} = \sec \theta \csc \theta$

$\frac{\cos \theta (\sin \theta + \cos \theta) - \sin \theta (\cos \theta - \sin \theta)}{\sin \theta \cos \theta}$

$\frac{\sin \theta \cos \theta + \cos^2 \theta - \sin \theta \cos \theta + \sin^2 \theta}{\sin \theta \cos \theta}$

$\frac{\cos^2 \theta + \sin^2 \theta}{\sin \theta \cos \theta}$

$\rightarrow \frac{1}{\sin \theta} \cdot \frac{1}{\cos \theta} = \csc \theta \sec \theta = \sec \theta \csc \theta$

$$3. \frac{\tan x + \cot x}{\tan x \cot x} = \tan x + \cot x$$

$$\frac{\tan x}{\tan x \cot x} + \frac{\cot x}{\tan x \cot x} = \tan x + \cot x$$

$$\cot x + \tan x = \tan x + \cot x$$

$$4. \sec x \csc x = 2 \csc x$$

$$\frac{1}{\cos x} \frac{1}{\sin x} = \frac{2}{\sin x}$$

$$\sec x \csc x - 2 \csc x = 0$$

$$\csc x (\sec x - 2) = 0$$

$$\csc x = 0$$

$$\frac{1}{\sin x} = 0$$

no solution

$$\sec x - 2 = 0$$

$$\sec x = 2$$

$$\cos x = \frac{1}{2}$$

$$1st: 60^\circ \frac{\pi}{3}$$

$$4th: 300^\circ \frac{5\pi}{3}$$

$$5. \tan^2 3x = 3 \quad - \text{Solve for } x - \text{no range}$$

$$\tan 3x = \sqrt{3}$$

$$3x = \pm \frac{60^\circ}{3}$$

$$x = \pm 20^\circ$$

$$1st:$$

$$2nd:$$

$$3rd:$$

$$4th:$$

$$3x = \frac{\pi}{3} + n\pi$$

$$\frac{2\pi}{3} + n\pi$$

$$3 \text{ then } 2$$

$$x = \frac{\pi}{9} + n\pi/3$$

$$x = \frac{2\pi}{9} + \frac{n\pi}{3}$$

$$\frac{5\pi}{9} + \frac{n\pi}{3}$$

$$\frac{7\pi}{9} + \frac{n\pi}{3}$$

Equations Praticce

12/17

$$1. \quad 5\sqrt{3}x - 3 = 8\sqrt{3}x$$

$$\begin{array}{r} -5\sqrt{3}x \\ -3 = 3\sqrt{3}x \\ \hline 3\sqrt{3} \quad 3\sqrt{3} \end{array}$$

$$-\frac{1}{\sqrt{3}} = x \rightarrow -\frac{\sqrt{3}}{3}$$

$$2. \quad 3x^2 + x - 1 = 0$$

Rooten w/ quad formula $\rightarrow$ by hand

$$x = 1.434 \text{ or } \frac{1 \pm \sqrt{13}}{6} \quad \checkmark \text{ to get}$$

$$x = -1.767$$

$$3. \quad 4x^2 - 8x = -3$$

$$4x^2 - 8x + 3 = 0$$

Rooten

$$x = 1.5$$

$$x = .5$$

$$(2x-3)(2x-1)$$

$$\begin{array}{r} 12 \quad 1x \\ -2 \quad 6 \quad 1x \\ \hline -8 \quad \checkmark \end{array} \quad \begin{array}{r} 2x-1 \\ 2x \quad 4x^2 - 2x \\ \hline 3 \quad -6x \quad 3 \end{array}$$

$$4. \quad 2x^2 - 3x - 2 = 0$$

rooten

$$x = 2$$

$$x = -.5$$

$$(2x+1)(x-2)$$

$$5. \quad x^2 - \frac{3}{2}x$$

$$x^2 - \frac{3}{2}x = 0$$


Rooten

$$x = 0$$

$$x = 1.5$$

$$\rightarrow 2x^2 - 3x = 0$$

$$\leftarrow x(2x-3)$$



Test Corrections

12/17

9. $\cos x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

$\cos x = \frac{-1 \pm \sqrt{13}}{6}$

$\cos x = .434$

$\cos^{-1} \cos^{-1}$

1.12r

$\cos x = -.767$

$\cos^{-1} \cos^{-1}$

2.44r

⊕ 1st 1.12r

⊕ 4th 5.16r

⊖ 2nd 2.44r

⊖ 3rd 3.84r

10. $4 \sin^2 x - 8 \sin x = -3$

$4 \sin^2 x - 8 \sin x + 3 = 0$

$(2 \sin x - 3)(2 \sin x - 1)$

↓

$\sin x = \frac{3}{2}$

↓

out of domain

$\sin x = \frac{1}{2}$

⊕ 30° $\frac{\pi}{6}$
⊕ 2nd $\frac{5\pi}{6}$

11. $2 \sec^2 \frac{x}{2} - 3 \sec \frac{x}{2} - 2 = 0$

$(2 \sec \frac{x}{2} + 1)(\sec \frac{x}{2} - 2) = 0$

↓

$\sec \frac{x}{2} = -\frac{1}{2}$

↓

$\cos \frac{x}{2} = -2$

out of

Range

↓

$\sec \frac{x}{2} = 2$

↓

$\cos \frac{x}{2} = \frac{1}{2}$

↓

$\frac{x}{2} = 60^\circ = \frac{\pi}{3}$

$\frac{x}{2} = \frac{\pi}{3}$

→ $\frac{2\pi}{3}$

12.

$$\tan^2 \theta = \frac{3}{2} \sec \theta$$

$$\sec^2 \theta - 1 = \frac{3}{2} \sec \theta$$

$$\sec^2 \theta - \frac{3}{2} \sec \theta - 1 = 0$$

$$2\sec^2 \theta - 3\sec \theta - 2 = 0$$

$$(\sec \theta - 2)(2\sec \theta + 1)$$

 $\downarrow$

$$\sec \theta = 2$$

 $\downarrow$

$$\sec \theta = -\frac{1}{2}$$

 $\downarrow$

$$\cos \theta = \frac{1}{2}$$

 $\downarrow$

$$\cos \theta = -2$$

$$\theta = \frac{\pi}{3}, \frac{5\pi}{3}$$

5 Trig Review Before Requiz

12/17

346

36. Simplify

$$\cot\left(\frac{\pi}{2} - x\right) \cos x$$

$$\tan x \cos x$$

42.
$$\frac{1 + \csc \theta}{\cot \theta + \cos \theta} = \sec \theta$$

$$\frac{1 + \csc \theta}{\frac{\cos \theta}{\sin \theta} + \cos \theta}$$

$$\frac{1 + \csc \theta}{\cos \theta \left(\frac{1}{\sin \theta} + 1 \right)}$$

$$\frac{1}{\cos \theta} = \sec \theta$$

48.
$$\frac{\tan \theta}{1 + \sec \theta} \cdot \frac{1 + \sec \theta}{\tan \theta} = 2 \csc \theta$$

$$\frac{\tan \theta (\tan \theta)}{1 + \sec \theta (\tan \theta)} \cdot \frac{1 + \sec \theta (1 + \sec \theta)}{\tan \theta (1 + \sec \theta)}$$

$$\frac{\tan^2 \theta + 1 + 2 \sec \theta + \sec^2 \theta}{\tan \theta (1 + \sec \theta)}$$

$$\frac{\tan^2 \theta + 1 + \sec^2 \theta + 2 \sec \theta}{\tan \theta (1 + \sec \theta)}$$

$$\frac{2 \sec^2 \theta + 2 \sec \theta + 1}{\tan \theta (1 + \sec \theta)} \rightarrow \frac{2 \sec \theta (\sec \theta + 1)}{\tan \theta (\sec \theta + 1)}$$

factor 2 sec θ

$$\frac{2 \sec \theta}{\tan \theta}$$

$$\frac{2 \sec \theta \cdot \cos \theta}{\frac{\cos \theta}{\sin \theta}} = \frac{2 \cdot \frac{1}{\cos \theta} \cdot \cos \theta}{\frac{\cos \theta}{\sin \theta}}$$

$$2 \frac{1}{\sin \theta}$$

$$2 \csc \theta$$

66.
$$\frac{1}{\sec x + 1} - \frac{1}{\sec x - 1}$$

$$= \frac{1(\sec x - 1)}{\sec x + 1(\sec x - 1)} - \frac{1(\sec x + 1)}{\sec x - 1(\sec x + 1)}$$

$$\frac{\sec x - 1}{\sec^2 x + \sec x - \sec x - 1} - \frac{\sec x + 1}{\sec^2 x + \sec x - \sec x - 1}$$

$$\frac{\sec x - 1 - (\sec x + 1)}{\sec^2 x - 1}$$

$$\frac{\sec x - 1 - \sec x - 1}{\sec^2 x - 1}$$

$$\frac{-2}{\sec^2 x - 1}$$

$$-2 \cot^2 x$$

p 353
12

$$\frac{\csc x - 1}{1 - \sin x} = \csc x \cdot \frac{1 - \sin x}{1 - \sin x} = \csc x \cdot \frac{1 - \sin x}{1 - \sin x}$$

$$\frac{\csc x - 1}{1 - \sin x} \cdot \frac{1 + \sin x}{1 + \sin x}$$

$$\frac{\csc x + \csc x \cdot \sin x - 1 - \sin x}{1 + \sin x - \sin x - \sin^2 x}$$

$$\frac{\csc x + \sin x \cdot \sin x - 1 - \sin x}{-\sin^2 x + 1}$$

$$\frac{1 - 1 - \sin x + \csc x}{-\sin^2 x + 1}$$

$$-\sin x + \frac{1}{\sin x}$$

$$-\sin x + \frac{1}{\sin x}$$

$$\frac{1}{\sin x} - 1$$

$$\frac{1 - \sin x}{\sin x}$$

$$\frac{1 - \sin x}{\sin x} \cdot \frac{1}{1 - \sin x}$$

$$\frac{1}{\sin x}$$

convert

$$14, \sec x - \cos x = \sin x \tan x$$

$$\frac{1}{\cos x} - \cos x = \frac{\sin x}{1} \cdot \frac{\sin x}{\cos x}$$

common denominator $\cos x$

$$\frac{1}{\cos x} - \frac{\cos x \cdot \cos x}{\cos x}$$

$$\frac{1 - \cos^2 x}{\cos x} = \frac{\sin^2 x}{\cos x} = \frac{\sin x \sin x}{\cos x}$$

$$\sin x \tan x$$

$$22, \sec^6 x (\sec x \tan x) - \sec^4 x (\sec x \tan x) = \sec^5 x \tan^3 x$$

$$\begin{aligned} &\sec^5 x \tan x (\sec^2 x - 1) \\ &\sec^5 x \tan x (\tan^2 x) \\ &\sec x \tan^3 x \end{aligned}$$

$$30, \frac{\tan x + \tan y}{1 - \tan x \tan y} = \frac{\cot x + \cot y}{\cot x \cot y - 1}$$

expand

$$\frac{1}{\cot x} + \frac{1}{\cot y} = \frac{1}{1 - \frac{1}{\cot x} \cdot \frac{1}{\cot y}}$$

common denom

$$\frac{1(\cot y)}{\cot x(\cot y)} + \frac{1(\cot x)}{\cot y(\cot x)}$$

$$\frac{\cot x \cot y}{\cot x \cot y} = \frac{1}{\cot x \cot y}$$

reciprocal

$$\frac{\cot y + \cot x}{\cot x \cot y - 1} = \frac{\cot x \cot y}{\cot x \cot y - 1}$$

$$\frac{\cot y \cot x}{\cot x \cot y - 1}$$

$$52. \frac{\cos x}{1 - \tan x} + \frac{\sin x \cos x}{\sin x - \cos x} \rightarrow \frac{\cos x}{1 - \frac{\sin x}{\cos x}} + \frac{\sin x \cos x}{\sin x - \cos x}$$

Common denominator

$$\frac{\cos x (1 + \tan x)}{(1 - \tan x)(1 + \tan x)}$$

$$\frac{\cos x}{\cos x - \sin x} \rightarrow \frac{\cos x}{\cos x - \sin x} + \frac{\cos x}{\cos x - \sin x}$$

$$\frac{\cos^2 x}{\cos x - \sin x} + \frac{\sin x \cos x}{\sin x - \cos x}$$

$$\frac{\cos^2 x}{\cos x - \sin x} - \frac{\sin x \cos x}{\sin x - \cos x} \rightarrow \frac{\cos^2 x - \sin x \cos x}{\cos x - \sin x}$$

355

$$64. \sec^4 x \tan^2 x = (\tan^2 x + \tan^4 x) \sec^2 x$$

$$\sec^2 x \sec^2 x \tan^2 x$$

$$\sec^2 x (1 + \tan^2 x) \tan^2 x$$

$$\sec^2 x (\tan^2 x + \tan^4 x)$$

✓

$$364. 12. 3 \cot^2 x - 1 = 0$$

$$3 \cot^2 x = 1$$

$$\cot^2 x = \frac{1}{3}$$

$$\cot x = \pm \sqrt{\frac{1}{3}}$$

$$\tan x = \pm \sqrt{3}$$

$$60^\circ + 2n\pi$$

$$\frac{\pi}{3} + 2n\pi$$

$$\frac{2\pi}{3} + 2n\pi$$

$$\frac{4\pi}{3} + 2n\pi$$

$$\frac{5\pi}{3} + 2n\pi$$

✓
correct

$$\frac{\pi}{3} + n\pi$$

$$\frac{2\pi}{3} + n\pi$$

$$16. (3\tan^2 x - 1)(\tan^2 x - 3) = 0 \quad \tan x = 3$$

$$3\tan^2 x = 1$$

$$\tan^2 x = \frac{1}{3}$$

$$\tan x = \pm \sqrt{1/3}$$

$$30^\circ$$

$$\frac{\pi}{6} + 2n\pi$$

$$\frac{5\pi}{6} + 2n\pi$$

$$\frac{7\pi}{6} + 2n\pi$$

$$\frac{11\pi}{6} + 2n\pi$$

Forgot

$$\frac{\pi}{3} + 2n\pi$$

$$\frac{2\pi}{3} + 2n\pi$$

$$\frac{4\pi}{3} + 2n\pi$$

$$\frac{5\pi}{3} + 2n\pi$$

$$\frac{\pi}{3} + n\pi$$

$$\frac{2\pi}{3} + n\pi$$

$$\frac{4\pi}{3} + n\pi$$

$$22. 2\sin^2 x = 2 + \cos x$$

$$2(1 - \cos^2 x) = 2 + \cos x$$

$$2 - 2\cos^2 x = 2 + \cos x$$

$$-2 + 2\cos^2 x = \cos x$$

$$\cos x + 2\cos^2 x = 0$$

$$\text{factor } \cos x (2\cos x + 1) = 0$$

$$x = 0$$

$$x = -1/2$$

$$\cos^{-1}(0) = 90^\circ$$

$$\cos^{-1}(-1/2) = 120^\circ$$

$$240^\circ$$

Unit circle error

$$\frac{\pi}{2}, \frac{3\pi}{2}$$

$$\frac{2\pi}{3}, \frac{4\pi}{3}$$

$$2\cos x = -1$$

$$\cos x = -1/2$$

$$24. \sec x \csc x = 2 \csc x$$

$$\sec x \csc x - 2 \csc x = 0$$

$$\csc x (\sec x - 2) = 0$$

$$\csc x = 0$$

$$\frac{1}{\sin x} = 0$$

no solution

$$\sin x = \frac{1}{0}$$

$$\sec x = 2$$

$$\cos x = \frac{1}{2}$$

$$60^\circ$$

$$\text{1st: } \frac{\pi}{3}$$

$$\text{4th: } \frac{5\pi}{3}$$

don't divide

split out

365

40. $\sec 4x = 2$
 $\cos 4x = \frac{1}{2}$
 $\cos^{-1} \cos^{-1}$

$\frac{60^\circ + 2\pi n}{4}$

$\frac{\pi}{12} + \frac{1}{2}\pi n$ write
 $\frac{\pi}{12} + \frac{n\pi}{2}$

34 $\csc^2 x = 3 \csc x + 4$
 $\csc^2 x - 3 \csc x - 4 = 0$
 $(\csc x - 4)(\csc x + 1) = 0$
 $\csc x = 4$ $\csc x = -1$
 $\sin x = \frac{1}{4}$ $\sin x = -1$
 $\sin^{-1} \sin^{-1}$

$\frac{300^\circ + 2\pi n}{4}$

$\frac{5\pi}{12} + \frac{1}{2}\pi n$
 $\frac{5\pi}{12} + \frac{n\pi}{2}$

3rd 270° 3π
 4th 270° 2π

46. $\tan \frac{x}{3} = 1$
 $\tan^{-1} \tan^{-1}$ start w/ 1 for tan on diagonal

$\frac{45^\circ + n\pi}{x3}$

$\frac{225^\circ + n\pi}{x3}$

$\frac{3\pi}{4} + 3n\pi$

$\frac{15\pi}{4} + 3n\pi$

50. $y = \sec^4\left(\frac{\pi x}{8}\right) - 4 \rightarrow 4 = \sec^4\left(\frac{\pi x}{8}\right)$

$(-2, 0)$
 $(2, 0)$

$\frac{1}{4} = \cos^4\left(\frac{\pi x}{8}\right)$

$\pm \sqrt[4]{\frac{1}{4}} = \cos\left(\frac{\pi x}{8}\right)$

$\sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} \cdot \sqrt{2} = 4 \rightarrow \pm \sqrt[4]{1} = \cos\left(\frac{\pi x}{8}\right)$
 $\rightarrow \pm \sqrt[4]{4} = \cos\left(\frac{\pi x}{8}\right)$

$\pm \frac{1}{\sqrt{2}} = \cos\left(\frac{\pi x}{8}\right)$

$\cos^{-1} \cos^{-1}$

$\pm \frac{\pi}{4} = \frac{\pi x}{8}$

$\pm \frac{2\pi}{\pi} = \frac{\pi x}{\pi}$

$x = 2$

$x = -2$

60. $\cos^2 x - 2 \cos x - 1 = 0$ $[0, \pi]$

Roots
 $\frac{-(-2) \pm \sqrt{(-2)^2 - 4(1)(-1)}}{2(1)}$

$2 \pm \sqrt{8}$


2.414 $\in (2.414)$
 -1.414

correct

9.1 Conics: Parabolas

1/2/08

Chart p632



(can define conics:

- intersection of planes & cones
- algebraically in terms of 2nd degree equation


$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$
- in terms of a locus (collection of points) satisfying a certain geometric property)

Circle - collection of all points that are equidistant from a fixed point (center $[h, k]$)


$$(x-h)^2 + (y-k)^2 = r^2$$

$$x^2 + y^2 = r^2$$

Parabola




Distance from Vertex to Focus =
Distance from Vertex to Directrix




$|P| = \text{Distance}$

center
Parabola

Directrix




Upright $y = x^2$
 $p > 0$




Directrix

Upside Down $y = -x^2$
 $p < 0$



To the left $x = -y^2$
 $p < 0$




To the right $x = y^2$
 $p > 0$

Standard Equation

$$\begin{array}{lcl}
 x^2 = 4py & \leftarrow h & (x-h)^2 = 4p(y-k) \\
 y^2 = 4px & \leftarrow k & (y-k)^2 = 4p(x-h)
 \end{array}$$

flipped

ex 1 Find the standard form of the parabola with the vertex at the origin and focus is at $(0, 4)$




$$p = 4 \oplus \downarrow$$

$$\begin{aligned} x^2 &= 4py \\ x^2 &= 4(4)y \\ x^2 &= 16y \\ \frac{1}{16}x^2 &= y \end{aligned}$$

p 637

12. Find vertex, directrix, focus + sketch

$$\begin{aligned} x + y^2 &= 0 \\ y^2 &= -x \quad \rightarrow p = -\frac{1}{4} \\ &\text{Opens } \leftarrow \end{aligned}$$



$$\begin{aligned} 14. \quad (x-5) + (y+4)^2 &= 0 \\ (y+4)^2 &= -(x-5) \\ (y-k)^2 &= 4p(x-h) \end{aligned}$$

$$\text{vertex} = (h, k) = (5, -4)$$


$$p = -\frac{1}{4}$$

$$\text{focus } (-\frac{1}{4}, 0)$$

← Opens

$$\text{directrix } \rightarrow x = \frac{1}{4} + 5$$

$$x = 5\frac{1}{4} \quad \text{original}$$



16.

$$y^2 - 4y - 4x = 0 \quad \downarrow \text{Complete the square}$$

$$y^2 - 4y = 4x$$


$$y^2 - 4y + 4 = 4x + 4 \quad \left(\frac{-4}{2}\right)^2$$

$$(y-2)^2 = 4(x+1)$$


$$\text{vertex} = (-1, 2)$$

$$4p = 4$$

$$p = 1$$



26.



Get equation

$$y^2 = 4px \quad \rightarrow \quad 6^2 = 4p(-2)$$

$$36 = -8p$$

$$p = -4.5$$

sideways

$$y^2 = 4(-4.5)x$$

$$y^2 = -18x$$

32.

$$\text{Directrix} \rightarrow y = 3$$


$$\text{Vertex } (0, 0)$$

$$\text{Focus} = (-3, 0)$$

$$p = -3$$

$$x^2 = 4(-3)y$$

$$x^2 = -12y$$



ex2

Find focus

$$y = -\frac{1}{2}x^2 - x + \frac{1}{2}$$

$$-2y = x^2 + 2x - 1$$

$$-2y + 1 = x^2 + 2x$$

$$x^2 + 2x + 1 = -2y + 1 + 1 \quad \downarrow \text{Complete the sq.}$$

$$(x+1)^2 = -2(y-1)$$


$$\text{vertex} = (-1, 1)$$

$$4p = -2$$

$$p = -\frac{1}{2}$$

$$\text{focus} = (-1, \frac{1}{2})$$

$$\text{directrix} \rightarrow y = \frac{3}{2}$$



20.

$$x = \frac{1}{4}(y^2 + 2y + 33)$$

$$4x = y^2 + 2y + 33$$

$$4x - 33 = y^2 + 2y$$

$$y^2 + 2y + 1 = 4x - 33 + 1$$


$$(y+1)^2 = 4x - 32$$

$$(y+1)^2 = 4(x-8)$$

vertex = (8, -1)

$4p = 4$

$p = 1$



Focus = (h+p, k)

Directrix $x = h-p$

sideways

22.

$$x^2 - 2x + 8y + 9 = 0$$

$$x^2 - 2x = -8y - 9$$


$$x^2 - 2x + 1 = -8y - 9 + 1$$

$$(x-1)^2 = -8(y+1)$$

vertex = (-1, 1)

$4p = -8$

$p = -2$



sideways

ex3

Find standard form

vertex (1, 0)

focus (2, 0)

$p = 1$



$$(y-0)^2 = 4(1)(y-1)$$

$$y^2 = 4(y-1)$$

638

40.

vertex (3, -3)

focus (3, - $\frac{9}{4}$)

$p = -3 - \frac{9}{4} = -\frac{21}{4}$

y changing

$$(x-h)^2 = 4p(y-k)$$

$$(x-3)^2 = 4\left(-\frac{21}{4}\right)\left(y+3\right)$$

$$(x-3)^2 = -21(y+3)$$



46.

Focus $(0,0)$ Directrix $\rightarrow y=4$ 


$$p = -2 \quad \text{vertex} = (h,k)$$

$$(x-h)^2 = 4p(y-k)$$

$$(x-0)^2 = 4(-2)(y-2)$$

$$x^2 = -8(y-2)$$

38.



$$(y-k)^2 = 4p(x-h)$$

$$(4-3)^2 = 4p(4.5-5)$$

$$1^2 = 4p(-.5)$$

$$1 = -2p$$

$$p = -\frac{1}{2}$$


$$(y-3)^2 = 4(-\frac{1}{2})(x-5)$$


$$(y-3)^2 = -2(x-5)$$


9.1 Conics: Parabolas


HW


1/2/08


3. $x^2 = -8y$  (d)

5. $(y-1)^2 = 4(x-3)$
vertex = $(3, 1)$ (a) 

7. $y = \frac{1}{2}x^2$ vertex = $(0, 0)$
 $4p = 2$
 $p = \frac{1}{2}$
focus $(0, \frac{1}{2})$
directrix $x = -\frac{1}{2}$ 

10. $y^2 = 3x$
vertex = $(0, 0)$
 $4p = 3$
 $p = \frac{3}{4}$
focus $(\frac{3}{4}, 0)$
directrix $y = -\frac{3}{4}$ 

14. $(x-5) + (y+4)^2 = 0$ vertex = $(5, -4)$
 $(y+4)^2 + (x-5) = 0$
 $4p = 1$
 $p = \frac{1}{4}$
directrix $x = y = -4\frac{3}{4}$
focus $(5\frac{1}{4}, -4)$ 

15. $y^2 + 6y + 8x + 25 = 0$ vertex = $(2, -3)$
 $y^2 + 6y = -8x - 25$
 $y^2 + 6y + 9 = -8x - 25 + 9$
 $(y+3)^2 = -8x - 16$
 $(y+3)^2 = -8(x+2)$
 $4p = -8$
 $p = -2$
focus = $(0, -3)$
directrix $y = 4$ 

21

$$(x^4)_{h=2} \left(\frac{x}{2} + x \right)$$

vertex = $(2, -3)$ $y = 4$

$$h = d$$

focus

focus = (7, 2.5)

Directly $y = -5.5$

22.

$$0 = 6 + 8x - 2x^2$$

$$6 - 8 = -2$$

$$1 + 2x - 8 = 1 + 2x - 8$$

$$(1+x) \rightarrow (1-x)$$


$$\begin{pmatrix} 1 & 1 & -1 \end{pmatrix} = x^2 + 1 \quad \checkmark$$

$$g_1 \sim dh$$

$$z = d$$

$$(1 - \frac{1}{2}) = \frac{1}{2}$$

Ex. 2. Lexical



29.


$$\begin{aligned} \text{Vertex} &= (0, 0) \\ \text{Focus} &= (-2, 0) \end{aligned}$$

Vertex = $(0,0)$

$$p = -2$$

$$x(2-4) = -2$$

$x_{p-1} = h$



33.

$$\text{Directrix: } x = 2$$

$$V_{r+x} = 0,0$$

$$\begin{array}{r} x^2 - 8x + 12 \\ x^2 - 4x - 2 \\ \hline -4x + 14 \end{array}$$

$$x(z-1)h = 4$$

$$x^2 - 8x + 16 = 0$$

35.

point (y, g)

Verf. $(0,0)$

max, w

~~$$d^2 h^2 = 27$$~~
~~$$d^2 h = 27$$~~
~~$$d^2 h = 27$$~~

~~$36 \times 6 = 24 \times 16$~~

~~$36 \times 6 = 24 \times 16$~~

$$x^2 = 4 \left(\frac{x}{3} \right)$$

✓

✓

39

$$\text{Vertex } (-2, 0) \quad \text{Focus } (-3/2, 0)$$

Focus $(-\frac{3}{2}, 0)$

$$\begin{aligned} (z+x) \frac{dz}{dx} &= z^2 \\ (z+x) \left(\frac{z}{h} \right) dh &= z^2 dh \\ (z+x) dh &= z(1-h) \\ (y-x) dh &= z(1-h) \end{aligned}$$

$$x^2 - 2(x+2)$$

$$y^2 = 4\left(\frac{z}{2}\right)\left(\frac{z}{2}\right) \quad y^2 = z^2$$

$$y^2 = 4\left(\frac{z}{2}\right)\left(\frac{z}{2}\right) \quad y^2 = z^2$$

$$y^2 = 4\left(\frac{z}{2}\right)\left(\frac{z}{2}\right) \quad y^2 = z^2$$

638

41.

Vertex (5, 2)

Focus (3, 2)

$$(y - k)^2 = 4p(x - h)$$

$$(y - 2)^2 = 4p(x - 5)$$

$$(y - 2)^2 = 4(-2)(x - 5)$$

$$(y - 2)^2 = -8(x - 5)$$

①

44.

Vertex (-2, 1)

Directrix $x = 1$

$$(y - k)^2 = 4p(x - h)$$

$$(y - 1)^2 = 4(-3)(x + 2)$$

$$(y - 1)^2 = -12(x + 2)$$

Correct ✓

56.

Road Design

32 ft wide

4 ft higher in center

(0, 0) vertex

(6, -4)

 $x^2 = 4py$

$$x^2 = 4(p)y$$

$$16^2 = 4(p)(-2/5)$$

$$p = -160$$


$$x^2 = -640y$$

$$-160 - 1$$

$$x^2 = 4(1/4)(1/3)$$

$$x^2 = 1/3$$

$$x = 1/69 \text{ ft} = 8 \text{ feet}$$




b) (1, 1/3)

639

57.

Highway Design



$$y^2 = 4px$$

$$800^2 = 4p(1000)$$

$$640,000 = 4000p$$

$$160 = p$$

$$y^2 = 4(160)x$$

$$y^2 = 640x$$

$$\begin{aligned} (1-2)x &= 4(1-x-1) \\ (1-2)x &= 4(-x) \\ (1-2)x &= -4x \\ (1-2)x + 4x &= 0 \\ 3x &= 0 \\ x &= 0 \end{aligned}$$

$$\begin{aligned} (1-2)x &= 4(1-x-1) \\ (1-2)x &= 4(-x) \\ (1-2)x &= -4x \\ (1-2)x + 4x &= 0 \\ 3x &= 0 \\ x &= 0 \end{aligned}$$


$$\begin{aligned} x &= 4(1-x-1) \\ x &= 4(-x) \\ x &= -4x \\ x + 4x &= 0 \\ 5x &= 0 \\ x &= 0 \end{aligned}$$

$$\begin{aligned} x &= 4(1-x-1) \\ x &= 4(-x) \\ x &= -4x \\ x + 4x &= 0 \\ 5x &= 0 \\ x &= 0 \end{aligned}$$

$$x + 4(1-x-1) = 0$$

$$\begin{aligned} x &= 4(1-x-1) \\ x &= 4(-x) \\ x &= -4x \\ x + 4x &= 0 \\ 5x &= 0 \\ x &= 0 \end{aligned}$$

$$\begin{aligned} x &= 4(1-x-1) \\ x &= 4(-x) \\ x &= -4x \\ x + 4x &= 0 \\ 5x &= 0 \\ x &= 0 \end{aligned}$$



9.1 Conics: Parabolas

Day 2


1/3

Tangent Lines to Parabolas


- have reflective properties related to the use of parabolas in constructing relative surfaces

The tangent line to a parabola at point "p" makes equal angles with the following 2 lines:

- the line passing thru the point of tangency in the focus
- the axis of the parabola



ex1 Find equation of tangent line to parabola $(4-y x^2)$ and point of tangency $(1,1)$



$$d_1 = \sqrt{(0-1)^2 + (1-1)^2} = \sqrt{1} = 1$$

$$d_2 = 1 - 0 = 1$$

$$d_1 = d_2 = 1$$

$$\frac{y_2 - y_1}{x_2 - x_1} = \frac{1 - 0}{1 - 0} \text{ slope} = 1$$

$$\text{equation} = 1x + 0$$

638


50.

$$x^2 = 2y \rightarrow 4p = 2$$

$$p = \frac{1}{2}$$

$(-3, 9/2)$ tangent
 $(0, 0)$ vertex
 $(0, 1/2)$ focus

$$(-3, 9/2)$$



$$d_1 = \sqrt{(-3-0)^2 + (9/2 - 1/2)^2}$$

$$d_1 = 5$$

$$d_2 = 5$$

$$b / \frac{1}{2} - 5 = -4\frac{1}{2}$$

$$(0, -4\frac{1}{2})$$

$$\frac{y_2 - y_1}{x_2 - x_1} = \frac{9/2 - (-9/2)}{-3 - 0} = \frac{9}{-3} = -3$$

$$y = -3x - 4\frac{1}{2}$$

9.1 Conics: Parabolas

HW Day 2

1/3

638


49.

Find equation of tangent line d_1

$$x^2 = 2y \rightarrow y = \frac{1}{2}x^2$$

$(4, 8)$ tangent

$(0, 0)$ vertex



$$d_1 = \sqrt{(4-0)^2 + (8-\frac{1}{2})^2}$$

$$\sqrt{72.25}$$

$$8.5$$

$$d_2 = 8.5$$

$$b/\frac{1}{2} - 8.5 = -8$$

$$\frac{y_2 - y_1}{x_2 - x_1} = \frac{8 - -8}{4 - 0} = \frac{16}{4} = 4$$


$$y = 4x - 8$$

51.

$$y = -2x^2 \rightarrow x^2 = -\frac{1}{2}y$$

$(-1, -2)$ tangent

$(0, 0)$ vertex



$$d_1 = \sqrt{(-1-0)^2 + (-2-\frac{1}{2})^2}$$

$$\sqrt{1 + 3.51}$$

$$2.125$$

$$b/\frac{1}{2} + 2.125 = -2.125$$

add

-4

$$\frac{y_2 - y_1}{x_2 - x_1} = \frac{-2 - \frac{1}{2}}{-1 - 0} = \frac{-2.5}{-1} = 2.5$$

$$y = 2.5x - 2.25$$

$$y = 4x + 2$$

pressed up w/ b

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3

1/3



1/3

1/3



1/3

9.2 Ellipses

Notes



An ellipse is the set of all points in a plane, the sum of whose distances from fixed points (foci) is constant



hypotenuse

$$d_1 = \sqrt{b^2 + c^2} = 2a$$

$$a = \sqrt{b^2 + c^2}$$

$$a^2 = b^2 + c^2$$

$$a^2 - b^2 = c^2$$


Standard Equations (0,0) center

biggest $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$
 x axis = major axis



$$\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$$

$\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$
 y axis = major axis



$$\frac{(x-h)^2}{b^2} + \frac{(y-k)^2}{a^2} = 1$$

eccentricity = measures the oval-ness

$$\frac{c}{a} = e$$

$0 < e < 1$
 $\nearrow$ circlely $\nearrow$ very long

ex

$$4x^2 + y^2 = 36$$

$$\frac{4x^2}{36} + \frac{y^2}{36} = 1$$

$$\frac{x^2}{9} + \frac{y^2}{36} = 1$$

$\nearrow$ bigger # = a^2

$$a^2 = 36$$

$$a = \pm 6$$

$$b^2 = 9$$

$$b = \pm 3$$


$$\text{vertex} = (0, \pm a) = (0, 6), (0, -6)$$

$$c^2 = a^2 - b^2$$

$$c^2 = 36 - 9$$

$$c^2 = 27$$

$$c = \pm \sqrt{27} = \pm 3\sqrt{3} = \pm 5.19$$



eccentricity

$$\frac{c}{a} = \frac{3\sqrt{3}}{6} = .577$$

646

8.

$$\frac{x^2}{64} + \frac{y^2}{28} = 1$$

$\nearrow a^2$ $\nearrow b^2$

$$a^2 = 64$$

$$a = 8$$

$$b^2 = 28$$

$$b = 5.29$$

$$\text{vertex} = (8, 0)$$

$$(-8, 0)$$

$$\text{minor vertex} = (0, 5.29)$$

$$(0, -5.29)$$

$$\text{eccentricity } \frac{c}{a} = \frac{6}{8} = .75$$

$$c^2 = a^2 - b^2$$


$$64 - 28$$

$$c = \sqrt{36}$$

$$c = 6$$

$$\text{Focus} = (6, 0)$$

$$(-6, 0)$$



ex 2

$$4x^2 + y^2 - 8x + 4y - 8 = 0$$

$$4x^2 - 8x + y^2 + 4y = 8 \quad \text{complete square twice}$$

$$4(x^2 - 2x + 1) + y^2 + 4y + 4 = 8 + 4 + 4$$

$$4(x-1)^2 + (y+2)^2 = 16$$

$$\frac{4(x-1)^2}{16} + \frac{(y+2)^2}{16} = 1$$

$$\frac{(x-1)^2}{4} + \frac{(y+2)^2}{16} = 1$$

major $\begin{pmatrix} 1, -2+4 \\ 1, -2-4 \end{pmatrix} \begin{pmatrix} 1, 2 \\ 1, -6 \end{pmatrix}$

minor $\begin{pmatrix} 1+2-2 \\ 1-2, -2 \end{pmatrix} \begin{pmatrix} 3, -2 \\ -1, -2 \end{pmatrix}$

$$c^2 = a^2 - b^2$$

$$c^2 = 16 - 4$$

$$c = 2\sqrt{3} = \pm 3.46$$

$$a^2 = 16$$

$$b^2 = 4$$

$$a = \pm 4$$

$$b = \pm 2$$

Center at $(1, -2)$ Foci $\begin{pmatrix} 1, -2+2\sqrt{3} \\ 1, -2-2\sqrt{3} \end{pmatrix}$

6.46

18.

$$x^2 + 4y^2 - 6x + 20y - 2 = 0$$

$$x^2 - 6x + 4y^2 + 20y = 2$$

$$(x^2 - 6x + 9) + (4y^2 + 20y + 25) = 2 + 9 + 25$$

$$(x-3)^2 + 4(y+2.5)^2 = 36$$

$$(x-3)^2 + 4(y+2.5)^2 = 36$$

$$\frac{(x-3)^2}{36} + \frac{4(y+2.5)^2}{36} = 1$$

$$\frac{(x-3)^2}{36} + \frac{(y+2.5)^2}{9} = 1$$

$$a^2 = 36$$

$$b^2 = 9$$

$$a = \pm 6$$

$$b = \pm 3$$

center $(3, -2.5)$

major $\begin{pmatrix} 3+6, -2.5 \\ 3-6, -2.5 \end{pmatrix} \begin{pmatrix} 9, -2.5 \\ -3, -2.5 \end{pmatrix}$


minor $\begin{pmatrix} 3, -2.5+3 \\ 3, -2.5-3 \end{pmatrix} \begin{pmatrix} 3, 0.5 \\ 3, -5.5 \end{pmatrix}$

focus $\begin{pmatrix} 3+5.196, -2.5 \\ 3-5.196, -2.5 \end{pmatrix} \begin{pmatrix} 8.196, -2.5 \\ -2.196, -2.5 \end{pmatrix}$

$$c^2 = a^2 - b^2$$

$$c^2 = 36 - 9$$

$$c = \sqrt{27} = 5.196$$





9.2 Ellipses

HW

1/3

646 2. $\frac{x^2}{9} + \frac{y^2}{4} = 1$ $a^2 = 9$ $a = 3$ major $(3, 0)$ $(-3, 0)$ (c)
 $\uparrow$ center $(0, 0)$


5. $\frac{(x-2)^2}{16} + (y+1)^2 = 1$ (a)
 $\uparrow$ center $(2, -1)$


6. $\frac{(x+2)^2}{9} + \frac{(y+2)^2}{4} = 1$ (e)
 $\uparrow$ center $(-2, -2)$


8. $\frac{x^2}{64} + \frac{y^2}{28} = 1$ Did in school

~~9.~~ $\frac{(x-4)^2}{16} + \frac{(y+1)^2}{25} = 1$ major $(4+4, -1)$ $(8, -1)$ $c^2 = a^2 - b^2$
 $\uparrow$ $a^2 = 16$ $a = 4$ $\uparrow$ $b^2 = 25$ $b = 5$ $(4-4, -1)$ $(0, -1)$ $12 - 25$
 flipped $\uparrow$ and $\downarrow$ minor $(4, -1+5)$ $(4, 4)$ $c = \sqrt{-13}$
 $(4, -1-5)$ $(4, -6)$ ± 3.60
 $(4+3.6, -1)$ $(7.6, -1)$ eccentricity
 $(4-3.6, -1)$ $(.4, -1)$ $\frac{c}{a} = \frac{3.6}{4} = .9$
 $(4, -1)$ center
 (x^2)

11. $\frac{(x+5)^2}{9/4} + (y-1)^2 = 1$ major $(-5+1.5, 1)$ $(-3.5, 1)$
 $\uparrow$ $a^2 = 9/4$ $a = 1.5$ $\uparrow$ $b^2 = 1$ $b = 1$ $(-5-1.5, 1)$ $(-6.5, 1)$
 $(-5, 1+1.5)$ $(-5, 2.5)$ minor
 $(-5, 1-1.5)$ $(-5, -.5)$
 $(-5+1.12, 1)$ $(-3.88, 1)$ fori
 $(-5-1.12, 1)$ $(-6.12, 1)$
 $(-5, 1)$ center


$c^2 = a^2 - b^2$
 $9/4 - 1$
 $\sqrt{1.25} = 1.12$

eccentricity
 $\frac{c}{a} = \frac{1.12}{1.5} = .75$

15. $9x^2 + 4y^2 + 36x - 24y + 36 = 0$

$9x^2 + 36x + 4y^2 - 24y = -36$

$9(x^2 + 4x + 4) + 4(y^2 - 6y + 9) = -36 + 36 + 36$

$9(x+2)^2 + 4(y-3)^2 = 36$

$\frac{9(x+2)^2}{36} + \frac{4(y-3)^2}{36}$

$\frac{(x+2)^2}{4} + \frac{(y-3)^2}{9}$

$r_b = 2 \quad r_a = 3$

center $(-2, 3)$



$c^2 = a^2 - b^2$

$9 - 4$

$c = \sqrt{5}$

2.23

major $(-2, 3+3) \quad (-2, 6)$

$(-2, 3-3) \quad (-2, 0)$

minor $(-2+2, 3) \quad (0, 3)$

$(-2-2, 3) \quad (-4, 3)$

foci $(-2, 3+2.23) \quad (-2, 5.23)$

$(-2, 3-2.23) \quad (-2, .76)$

$\frac{c}{a} = \frac{2.23}{3} = .74$

19. $16x^2 + 25y^2 - 36x - 50y + 16 = 0$

$16x^2 - 36x + 25y^2 - 50y = -16$

$16(x^2 - 2x + 1) + 25(y^2 - 2y + 1) = -16 + 16 + 25$

$16(x-1)^2 + 25(y-1)^2 = 25$


$\frac{16(x-1)^2}{25} + \frac{25(y-1)^2}{25}$

$\frac{(x-1)^2}{(25/16)} + \frac{(y-1)^2}{1} = 1$

$r_b = 1 \quad r_a = 1.25$

$a = 1.25$

center $(1, 1)$



major $(1+1.25, 1) \quad (2.25, 1)$

$(1-1.25, 1) \quad (-.25, 1)$

minor $(1, 1+1) \quad (1, 2)$

$(1, 1-1) \quad (1, 0)$

foci $(1+1.75, 1) \quad (2.75, 1)$

$(1-1.75, 1) \quad (-.75, 1)$

$c^2 = a^2 - b^2$

$\frac{25}{16} - 1$

$c = \sqrt{.5625}$

$c = .75$

$\frac{c}{a} = \frac{.75}{1.25} = .6$

(V)

20. $9x^2 + 25y^2 - 36x - 50y + 61 = 0$

$9x^2 - 36x + 25y^2 - 50y = -61$


$9(x^2 - 4x + 4) + 25(y^2 - 2y + 1) = -61 + 36 + 25$

$9(x-2)^2 + 25(y-1)^2 = 0$

degenerate conic

↑ that's all - just write

23.



$$\frac{x^2}{2^2} + \frac{y^2}{4^2} = \frac{x^2}{4} + \frac{y^2}{16} = 1$$

the account of some

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100

9.2 Ellipses

Day 2

1/4

647

26. Find the standard equation of an ellipse
 center $(0,0)$ major axis is y axis
 vertex $(0, \pm 8)$ c
 foci $(0, \pm 4)$ c^2

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\begin{aligned} c^2 &= a^2 - b^2 \\ 16 &= 64 - b^2 \\ -48 &= -b^2 \\ b^2 &= 48 \end{aligned}$$

$$\frac{y^2}{64} + \frac{x^2}{48} = 1$$

28. Foci $(\pm 2, 0)$ $c=2$ c x axis
 major axis length $= 6 \in 2a$
 $a=3$

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

$$\begin{aligned} c^2 &= a^2 - b^2 \\ 4 &= 9 - b^2 \\ b^2 &= 5 \\ b &= \sqrt{5} \end{aligned}$$

$$\frac{x^2}{9} + \frac{y^2}{5} = 1$$

30. Center $\neq (0,0)$ c Center $(2,1)$
 Foci $(0,1)$ $(4,1)$ $c=2$
 major axis length $= 6 \in 2a$
 $a=3$

$$\begin{aligned} c^2 &= a^2 - b^2 \\ 4 &= 9 - b^2 \\ b^2 &= 5 \\ b &= \sqrt{5} \end{aligned}$$

$$\frac{(x-2)^2}{9} + \frac{(y-1)^2}{5} = 1$$

34. Foci: $(0,0)$ $(4,0)$ c center $(2,0)$
 major axis length $= 8 \in 2a$ x -axis
 $a=4$ $c=2$

$$\begin{aligned} c^2 &= a^2 - b^2 \\ 4 &= 16 - b^2 \\ b^2 &= 12 \\ b &= \sqrt{12} \end{aligned}$$

$$\frac{(x-2)^2}{16} + \frac{y^2}{12} = 1$$

36. center $(2,-1)$ y -axis
 vertex $(2, \frac{1}{2})$ $a = \frac{3}{2}$
 minor axis $= 2 \in b=1$

$$\frac{(y+1)^2}{9/4} + \frac{(x-2)^2}{1} = 1$$

h always goes with x^2

proper way $\frac{(x-2)^2}{1} + \frac{4(y+1)^2}{9} = 1$

38.

center $(3, 2)$

$$a = 3c$$

foci $(1, 2)$ $(5, 2)$ $\in x$ axiscenter $(3, 2)$

$$c = 2$$

$$a = 3(2)$$

$$a = 6$$

$$c^2 = a^2 - b^2$$

$$4 = 36 - b^2$$

$$b^2 = 32$$

$$b = \sqrt{32}$$

$$\frac{(x-3)^2}{36} + \frac{(y-2)^2}{32} = 1$$

42.

$$e = \frac{1}{2} \in \frac{c}{a} = \frac{1}{2} \quad c = 4$$

$$a = 8$$

vertex $(0, \pm 8)$ $\in$ center $(0, 0)$ $\in y$ axis

$$a = 8$$

$$c^2 = a^2 - b^2$$

$$16 = 64 - b^2$$

$$b^2 = 48$$

$$b = \sqrt{48}$$

$$\frac{x^2}{48} + \frac{y^2}{256} = 1$$

9.2 Ellipses

HLW Day 2

1/4

646

21. $12x^2 + 20y^2 - 12x + 40y - 37 = 0$

$$12x^2 - 12x + 20y^2 + 40y = 37$$

$$12(x^2 - x + \frac{1}{4}) + 20(y^2 + 2y + 1) = 37 + 3 + 20$$

$$12(x - \frac{1}{2})^2 + 20(y + 1)^2 = 60$$

$$12(x - \frac{1}{2})^2 + 20(y + 1)^2 = 1$$

①

$$\frac{(x - \frac{1}{2})^2}{\frac{5}{12}} + \frac{(y + 1)^2}{\frac{3}{20}} = 1$$


$$a^2 = \frac{5}{12}$$

$$b^2 = \frac{3}{20}$$

$$a = \sqrt{\frac{5}{12}}$$

$$b = \sqrt{\frac{3}{20}}$$

center $(\frac{1}{2}, -1)$



major $(\frac{1}{2} + \sqrt{\frac{5}{12}}, -1)$

$(\frac{1}{2} - \sqrt{\frac{5}{12}}, -1)$

minor $(\frac{1}{2}, -1 + \sqrt{\frac{3}{20}})$

$(\frac{1}{2}, -1 - \sqrt{\frac{3}{20}})$

foci $(\frac{1}{2} + \sqrt{\frac{5}{12}}, -1)$

$(\frac{1}{2} - \sqrt{\frac{5}{12}}, -1)$

$$c^2 = a^2 - b^2$$

$$c^2 = \frac{5}{12} - \frac{3}{20}$$

$$c^2 = \frac{1}{2}$$

$$c = \sqrt{\frac{1}{2}}$$

$$e = \frac{c}{a}$$

$$\frac{\sqrt{\frac{1}{2}}}{\sqrt{\frac{5}{12}}} = \frac{\sqrt{6}}{5}$$

647

25.

Vertices $(\pm 3, 0) \Rightarrow a = 3$

Foci $(\pm 2, 0) \Rightarrow c = 2$

center $(0, 0)$

$$c^2 = a^2 - b^2$$

$$4 = 9 - b^2$$

$$b = \sqrt{5}$$

$$\frac{x^2}{9} + \frac{y^2}{5} = 1$$

①

27.

Foci $(\pm 4, 0) \Rightarrow c = 4$

major length = 10 $\Rightarrow a = 5$

center $(0, 0)$

$$c^2 = a^2 - b^2$$

$$16 = 25 - b^2$$

$$b^2 = 9$$

$$b = 3$$

$$\frac{x^2}{25} + \frac{y^2}{9} = 1$$

29.

Vertices $(0, \pm 5) \Rightarrow a = 5$

passes through $(4, 2)$

$$\frac{x^2}{b^2} + \frac{y^2}{25} = 1$$

$$\frac{4^2}{b^2} + \frac{2^2}{25} = 1$$

$$-\frac{4}{25} - \frac{4}{25}$$

$$\frac{16}{b^2} = \frac{21}{25}$$

$$400 = 21b^2$$

$$400/21 \Rightarrow 19 = b^2$$

$$4.36$$

$$\frac{x^2}{400/21} + \frac{y^2}{25} = 1$$

proper way

$$\frac{21x^2}{400} + \frac{y^2}{25} = 1$$

647

Find the equation

33. vertices $(0,2)$ $(4,2)$ $\leftarrow$ center $(2,2)$
 $a=2$
 minor axis length $= 2 \rightarrow b=1$

$$\frac{(x-2)^2}{4} + \frac{(y-2)^2}{2} = 1$$

36. center $(2,-1)$
 vertices $(2, \frac{1}{2})$
 minor axis length $= 2 \rightarrow b=1$

did in class

39. center $(0,4)$
 $a=2c$ $4=2c$ $c=2$
 vertices $(-4,4)$ $(4,4)$
 $\leftarrow$ x axis $a=4$

$$c^2 = a^2 - b^2$$

$$4 = 16 - b^2$$


$$b = \sqrt{12}$$

$$\frac{x^2}{16} + \frac{(y-4)^2}{12} = 1$$

43. Architecture $\downarrow$ x-axis
 major axis $= 100$ ft $\rightarrow a=50$
 height $= 40$ ft $\rightarrow b=40$

$$\frac{x^2}{50^2} + \frac{y^2}{40^2} = 1$$

$$\frac{x^2}{2500} + \frac{y^2}{1600} = 1$$




c) $(45, y) \rightarrow \frac{45^2}{2500} + \frac{y^2}{1600} = 1$

$$\sqrt{304}$$

17.4 ft

647
45.

Geometry



Area of ellipse = $2 \times$ area of circle

Area of ellipse = πab

$\pi(10)^2$ ← wrong formula for area

area circle = $\pi(10)^2$

~~6283~~ 100π

area ellipse = ~~125.66~~ 200π

$200\pi + 25.66 = \pi a(10)$

$200\pi = 10a$

$20\pi = a$

length major axis = ~~40~~

46. Astronomy - Halley's Comet

elliptical orbit

sun = focus

eccentricity = .97 $\leftarrow \frac{c}{a}$

length major axis = 35.88 AU $\leftarrow 2a$

1 AU = 93 million miles

$a = 17.94$

$\frac{c}{.97} = \frac{c}{17.94}$

$c = 17.40$

Equation?

x axis

(0,0) center


$c^2 = a^2 - b^2$ 321.84 ← wrong #

$302.878 = 1287.3744 - b^2$

$b = \sqrt{984.4964} = 31.377$

$b = 31.377$ 4.36

$$\frac{x^2}{321.84} + \frac{y^2}{19.01} = 1$$



1000 - 1000

1000 - 1000

1000 - 1000

1000 - 1000


1000 - 1000

9.3 Hyperboles


Notes

1/7


asymptotes



$$\begin{cases} y = x \\ y = -x \end{cases} \begin{matrix} \text{rise} \\ \text{run} \end{matrix} X \rightarrow \pm \frac{a}{b} X$$




$$y - k = \pm \frac{b}{a} (x - h)$$



$$d_2 - d_1 = 2a$$

The set of all points (x, y) in a plane - the differences of whose 2 distinct fixed points (foci) is constant $= 2a$

$c > a$ then $c^2 > a^2 \rightarrow c^2 = a^2 + b^2$
 ellipse: $c^2 < a^2 \rightarrow c^2 = a^2 - b^2$



$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

a^2 is always positive (+)
so x is major axis

$$\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$$

a^2 is always (+)
so y is major axis

$$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$$

$$\frac{(y-k)^2}{a^2} - \frac{(x-h)^2}{b^2} = 1$$

note

656
8.

$$\frac{y^2}{9} - \frac{x^2}{1} = 1$$

$$a^2 = 9 \quad b^2 = 1$$


$$a = 3 \quad b = 1$$

$$c^2 = a^2 + b^2$$

$$c = \sqrt{10}$$

$$c = \pm 3.16$$

$$y = \pm \frac{a}{b}x = \pm \frac{3}{1}x = y = \pm 3x$$



14. $\frac{(y-1)^2}{1/4} - \frac{(x+3)^2}{1/16} = 1$ center $(-3, 1)$

$$a^2 = \frac{1}{4}$$

$$b^2 = \frac{1}{16}$$


$$a = \pm \frac{1}{2}$$

$$b = \pm \frac{1}{4}$$

$$c^2 = a^2 + b^2$$

$$c = \sqrt{5/16} \rightarrow \frac{\sqrt{5}}{\sqrt{16}} \rightarrow \frac{\sqrt{5}}{4}$$

$$c = \pm 1.559$$



vertices $(-3, 1 \pm \frac{1}{2})$

foci $(-3, 1 \pm \frac{\sqrt{5}}{4})$

asymptote $y - 1 = \pm \frac{a}{b}(x + 3)$

$$y - 1 = \pm 2(x + 3)$$

$$y = 2x + 7$$

$$y = -2x - 5$$

mod

18. $6y^2 - 3x^2 = 18$

$$\frac{6y^2}{18} - \frac{3x^2}{18} = 1$$

$$\frac{y^2}{3} - \frac{x^2}{6} = 1$$

$$a^2 = 3 \quad b^2 = 6$$

$$a = \sqrt{3} \quad b = \sqrt{6}$$

$$c^2 = a^2 + b^2$$


$$c = \sqrt{9}$$

$$c = 3$$

vertices $(0, \pm\sqrt{3})$

foci $(0, \pm 3)$

$$r y = \pm \frac{a}{b} x = \pm \frac{\sqrt{3}}{\sqrt{6}} \cdot \frac{\sqrt{6}}{\sqrt{6}} = \frac{\sqrt{18}}{6} = \frac{3\sqrt{2}}{6} = \boxed{\frac{\sqrt{2}}{2}}$$



20. $x^2 - 9y^2 + 36y - 72 = 0$

$$x^2 - 9(y^2 - 4y + 4) = 72 - 36$$

$$x^2 - 9(y - 2)^2 = 36$$

$$\frac{x^2}{36} - \frac{(y-2)^2}{4} = 1$$

$$a^2 = 36 \quad b^2 = 4$$

$$a = \pm 6 \quad b = \pm 2$$

center $(0, 2)$

$$c^2 = a^2 + b^2$$


$$c = \sqrt{40} = 2\sqrt{10}$$

$$c = \pm 6.32$$

center $(0, 2)$

vertices $(\pm 6, 2)$

foci $(\pm 6.32, 2)$



$$y - 2 = \pm \frac{b}{a} x$$

$$y - 2 = \pm \frac{2}{6} x$$

$$y = \pm \frac{1}{3} x + 2$$

$$y = \frac{1}{3} x + 2$$

$$y = -\frac{1}{3} x + 2$$

656

hyperbola
 y axis $\frac{y}{a} = 1$
 $a = 3$

28.

vertices $(0, \pm 3)$ asymptotes $y = \pm 3 \Rightarrow \frac{a}{b} = \frac{3}{1}$

$$\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$$

$$\frac{y^2}{9} - \frac{x^2}{1} = 1$$

30.

hyperbola
 x axis $\frac{x}{a} = 1$
 Foci $(\pm 10, 0)$ $c = 10$

asymptotes $y = \pm \frac{3}{4}x$ $\Rightarrow \frac{b}{a} = \frac{3}{4}$
 $b = 3$
 $a = 4$

$$10^2 = 3^2 + 4^2 \quad (\times)$$

$$10^2 = (3 \cdot 2)^2 + (4 \cdot 2)^2 \quad \downarrow \cdot 2$$

$$100 = 36 + 64 \quad (\checkmark)$$

$$\frac{x^2}{64} + \frac{y^2}{36} = 1$$

657

36.

hyperbola
 x axis $\frac{x}{a} = 1$
 Vertices $(2, 1)$ $(-2, 1)$ $\Rightarrow$ center $(0, 1)$
 Passes through $(5, 4)$

$$\frac{x^2}{a^2} - \frac{(y-1)^2}{b^2} = 1$$

$$\frac{5^2}{a^2} - \frac{(4-1)^2}{b^2} = 1$$

$$\frac{25}{a^2} - \frac{9}{b^2} = 1$$

$$-\frac{21}{4} = \frac{-9}{b^2}$$

$$-36 = -21b^2$$

$$b^2 = \frac{36}{21}$$

$$b = 1.30$$

$$\frac{x^2}{a^2} - \frac{(y-1)^2}{36/21} = 1$$

$$\frac{x^2}{a^2} - \frac{21(y-1)^2}{36} = 1$$

$$\frac{x^2}{4} - \frac{7(y-1)^2}{12} = 1$$

6.3 Hyperbola

HW

1/7

Q57 9.

Find stuff

$$\frac{y^2}{25} - \frac{x^2}{81} = 1$$

$$a^2 = 25 \quad b^2 = 81$$

$$a = 5 \quad b = 9$$

center (0,0)



$$c^2 = a^2 + b^2$$


$$c^2 = 25 + 81$$

$$c = \sqrt{106}$$

$$c = 10.3$$

vertices (0, ±5)

foci (0, ±10.3)



$$y = \pm \frac{a}{b} x = \pm \frac{5}{9} x$$

$$y = \frac{5}{9} x$$


$$y = -\frac{5}{9} x$$

11. $\frac{(x-1)^2}{4} - \frac{(y+2)^2}{1} = 1$

$$a^2 = 4 \quad b^2 = 1$$

$$a = 2 \quad b = 1$$

center (1, -2)



$$c^2 = a^2 + b^2$$


$$c^2 = 4 + 1$$

$$c = \sqrt{5}$$

$$c = 2.23$$

vertices (1 ± 2, -2)

foci (1 ± 2.23, -2)



$$y = \pm \frac{b}{a} (x-1) - 2$$

$$y+2 = \pm \frac{1}{2} (x-1)$$

$$y = \pm \frac{1}{2} (x-1) - 2$$

$$13, \quad \frac{(y+5)^2}{1/9} - \frac{(x-1)^2}{1/4} = 1$$

$$a^2 = 1/9 \quad b^2 = 1/4$$

$$a = 1/3 \quad b = 1/2$$

$$\text{center } (1, -5)$$



$$c^2 = a^2 + b^2$$

$$c = \sqrt{13/36}$$

$$c = .60$$


$$\text{vertices } (1, -5 \pm 1/3)$$

$$\text{foci } (1, -5 \pm \sqrt{13/36})$$

$$y = \frac{a}{b}x \quad y+5 = \pm \frac{1/3}{1/2}x - 1$$

$$y+5 = \pm 2/3(x-1)$$

$$y = -5 \pm 2/3(x-1)$$



$$17, \quad 2x^2 - 3y^2 = 6$$

$$\frac{2x^2}{6} - \frac{3y^2}{6} = 1$$

$$\frac{x^2}{3} - \frac{y^2}{2} = 1$$

$$a^2 = 3 \quad b^2 = 2$$

$$a = \sqrt{3} \quad b = \sqrt{2}$$

$$\text{center } (0, 0)$$


$$c^2 = a^2 + b^2$$

$$c = \sqrt{5}$$

$$c = 2.24$$

$$\text{vertices } (\pm\sqrt{3}, 0)$$

$$\text{foci } (\pm\sqrt{5}, 0)$$



$$y = \pm \frac{b}{a}x$$

$$y = \pm \frac{\sqrt{2}}{\sqrt{3}}x \quad y = \pm \frac{\sqrt{6}}{3}$$

19.

$$4x^2 - y^2 - 36x - 6y + 18$$

$$4x^2 - 36x - y^2 - 6y = -18$$

$$4(x^2 - 9x + 4) - (y^2 - 6y + 9) = -18 + 36 - 9$$

$$4(x-2)^2 - (y-3)^2 = 9$$

$$\frac{(x-2)^2}{\frac{9}{4}} - \frac{(y-3)^2}{9} = 1$$

$$a^2 = \frac{9}{4}$$

$$b^2 = 9$$

$$a = \frac{3}{2}$$

$$b = 3$$

center (2, 3)

H/H

$$c^2 = a^2 + b^2$$

$$c = \sqrt{10}$$


$$c = 3.16$$

vertici (2 ± 1.5, 3)

foci (2 ± 3.16, 3)

$$y - 3 = \pm \frac{3}{2}(x - 2)$$

$$y = 3 \pm 3(x - 2)$$



22. $16y^2 - x^2 + 2x + 64y + 63 = 0$

$$16y^2 + 64y - x^2 + 2x = -63$$

$$16(y^2 + 4y + 4) - (x^2 - 2x + 1) = -63 + 64 - 1$$

$$16(y+2)^2 - (x-1)^2 = 0$$


degenerate conic

↓ can go further → degenerate hyperbola
pair of intersecting lines

$$(x+2)^2 = (x-1)^2 \cdot \frac{1}{16} \in \text{set } \Rightarrow \text{to } y$$

$$x+2 = \pm \frac{(x-1)}{4}$$

$$x+2 = \pm (x-1) \cdot \frac{1}{4}$$



draw like that

23. $9y^2 - x^2 + 2x + 54y + 62 = 0$

$9y^2 + 54y - x^2 + 2x = -62$

$9(y^2 + 6y + 4) - (x^2 - 2x + 1) = -62 + 81 - 1$

$9(y+3)^2 - (x+1)^2 = -18$

$\frac{(y+3)^2}{2} - \frac{(x+1)^2}{18} = 1$

$a^2 = 2$

$b^2 = 18$

$a = \sqrt{2}$

$b = \sqrt{18}$

center $(-1, -3)$




$c^2 = a^2 + b^2$

$c = \sqrt{20}$

$c = 4.47$

vertex $(-1, -3 \pm \sqrt{2})$

foci $(-1, -3 \pm \sqrt{20})$



27. Vertex $(\pm 1, 0)$ $c = 1$ y axis

asymptote $y = \pm 5x$

$\frac{b}{a} = \frac{b}{1} = 5 \Rightarrow b = 5$

$\frac{y^2}{1} + \frac{x^2}{5^2} = 1$

$\frac{y^2}{1} + \frac{x^2}{25} = 1$

29. Foci $(0, \pm 8)$ $c = 8$, x axis

asymptotes $y = \pm 4x$

center $(0, 0)$

$\frac{a}{b} = 4$

$b^2 = c^2 - a^2$

$a = 4b$

$b^2 = 8^2 - 16b^2$

$a^2 = 16b^2$

$b^2 + 16b^2 = 64$

$b^2(1+16) = 64$

$17b^2 = 64$

$b = \sqrt{64/17}$

$\frac{x^2}{\frac{1024}{17}} + \frac{y^2}{\frac{64}{17}} = 1 \Rightarrow \frac{17x^2}{1024} + \frac{17y^2}{64} = 1$

$a^2 = 16(\sqrt{64/17})^2$

$16 \cdot (64/17)$

$a^2 = \frac{1024}{17}$

$\frac{1024}{17}$

657

31. Vertices $(2,0)(6,0)$ $\in a=2$ $c^2=a^2+b^2$ $\frac{(x^2-4)}{4} + \frac{y^2}{12} = 1$
 foci $(0,0)(8,0)$ $\in c=4$ $16=4+b^2$ $b=\sqrt{12}$
 $\uparrow$ x axis
 $\parallel$

35. Vertices $(2,3)(2,-3)$ $\in (2,0)$ center
 passes through pt $(0,5)$ $a=3$
 y axis
 $\neq$

$$\frac{y^2}{9} - \frac{(x-2)^2}{b^2} = 1$$

$$\frac{25}{9} - \frac{(-2)^2}{b^2} = 1$$

$$-\frac{16}{9} = -\frac{4}{b^2}$$

$$4 \cdot 4 = 16b^2$$

$$b^2 = 9/4$$

$$b = 3/2$$

$$\frac{y^2}{9} - \frac{(x-2)^2}{9/4} = 1$$

$$\frac{y^2}{9} - \frac{4(x-2)^2}{9} = 1$$

37. Vertices $(0,4)(0,0)$ $\in a=2$
 point $(\sqrt{5}, -1)$ y axis
 center $(0,2)$ $\neq$

$$\frac{(y^2-2)}{4} - \frac{x^2}{4} = 1$$

$$\frac{(y^2-2)}{4} - \frac{x^2}{b^2} = 1$$

$$-\frac{1}{4} - \frac{5}{b^2} = 1$$

$$\frac{5}{4} - \frac{5}{b^2} = -20 = 5b^2$$

$$b = \sqrt{4}$$

$$b = 2$$

38. Vertices $(1,2)(3,2)$ $\in$ center $(2,2)$

asymptote $y = x$ $a=1$

$y = 4 - x$ x axis $\parallel$

a point $(1,1)$

$$\frac{(y^2-2)}{1} - \frac{(x^2-2)}{b^2} = 1$$

$$\frac{-1}{1} - \frac{-1}{b^2} = 1$$

$$+1 = \frac{1}{b^2} + 1$$

$$-\frac{1}{b^2} = \frac{1}{b^2} \quad 2b^2 = 1$$

$$b = 1$$

also now $\frac{(y^2-2)}{1} - \frac{(x^2-2)}{1} = 1$

$$\frac{b}{a} = \frac{b}{1} = 1$$

$$b = 1$$

Don't do

41, Vertices $(0,2)(6,2)$ $\in$ center $(3,2)$
 $a=3$
 $asy = y = \frac{2}{3}x$
 $\wedge y = 4 - \frac{2}{3}x$
 point $(6,4)$
 Xaxis $\cancel{y}$

$$\frac{(x-3)^2}{9} - \frac{(y-2)^2}{b^2} = 1$$

$$\frac{(6-3)^2}{9} - \frac{(4-2)^2}{b^2} = 1$$

$$\frac{9}{9} - \frac{4}{b^2} = 1$$

$$- \frac{4}{b^2} = 1$$

$$- b^2 = -4$$

$$b = \sqrt{4}$$

$$b = 2$$

$$\frac{(x-3)^2}{9} - \frac{(y-2)^2}{4} = 1$$

(1)

9.3 Hyperbolas


More Notes

1/8

657

46. Hyperbolic mirror - reflects light coming to 1 focus to another - focus = $(24, 0)$

top edge $(24, 24)$ vertex = ?



$\uparrow$ x axis $\downarrow$
 $c = 24$

$$c^2 = a^2 + b^2$$

$$24^2 = a^2 + b^2$$

$$576 - a^2 = b^2$$

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

put in point

$$\frac{x^2}{a^2} - \frac{y^2}{(576 - a^2)} = 1 \quad \downarrow \quad \frac{24^2}{a^2} - \frac{24^2}{(576 - a^2)}$$

$$\frac{576}{a^2} - \frac{576}{576 - a^2} = 1 \quad \leftarrow \text{LCD}$$

$$\frac{576(576 - a^2)}{a^2(576 - a^2)} - \frac{576(a^2)}{(576 - a^2)a^2}$$

$$\frac{331776 - 576a^2}{576a^2 - a^4} - \frac{576a^2}{576a^2 - a^4} = \frac{297216 - 576a^2 - 576a^2}{576a^2 - a^4}$$

$$\frac{331776 - 576a^2}{+a^4} - \frac{576a^2}{-576a^2} = \frac{297216 - 576a^2 - 576a^2}{-576a^2 + a^4}$$

$$a^4 - 1728a^2 + 331776 = 0$$

$$a^2 = \frac{-(-1728) \pm \sqrt{(1728)^2 - 4(1)(331776)}}{2(1)}$$

$$a^2 = 1507.99$$

$$a = \pm 38.83$$

$$a^2 = 220.12$$

$$a = \pm 14.83$$

$c > a$

⊗


$(\pm 14.83, 0)$

⊙


eccentricity = $\frac{c}{a}$ because $c > a$, $e > 1$

ex $\frac{10}{4} = \frac{c}{a}$ = bigger c - the bigger the e

If eccentricity is large - the branches are flatter



If eccentricity is close to 1 - the branches are more narrow



654

ex 5


2 microphones are 1 mile (5280 ft) apart

Record explosion

Mic A receives sound 2 sec before mic b

Where explosion?

Sound travels at 1100 ft/sec



$$2c = 5280$$

$$c = 2640$$

$$d_2 - d_1 = 1100 \cdot 2 = 2a$$

$$2200 \text{ ft}$$

$$a = 1100 \text{ ft}$$

$$c^2 = a^2 + b^2$$

$$2640^2 = 1100^2 + b^2$$

$$b = \sqrt{5759600}$$

$$b = 2399.92$$

path

$$\frac{x^2}{1100^2} + \frac{y^2}{5759600} = 1$$

$$\frac{x^2}{1210000} + \frac{y^2}{5759600} = 1$$

Type of Conic

all conics have been parallel to x or y axis

$$Ax^2 + Cy^2 + Dx + Ey + F = 0$$

Circle $A = C$ $A \neq 0$

Parabola $AC = 0$ $A = 0$ or $C = 0$ - not both

Ellipse $AC > 0$ $A + C$ like signs

Hyperbola $AC < 0$ $A + C$ have unlike signs

657

48 $x^2 + 4y^2 - 6x + 16y + 21 = 0$

$AC > 0$

$4 > 0$ ellipse

658

50 $y^2 - 4y - 4x = 0$

$\because A = 0$ so $AC = 0$ parabola

54 $4x^2 + 4y^2 - 16y + 15 = 0$

$4x^2 + 4y^2$

circle

$C^2 = 0$
 $SCMO = 1100$
 $D = 100000$
 $H = 100000$
 100000

Types of Conic

all conics have three branches

$$Ax^2 + Cy^2 + Dx + Ey + F = 0$$

Circle $A=C$
 Ellipse $A \neq C$
 Hyperbola $A \neq C$
 Parabola $A=0$ or $C=0$

$$x^2 + y^2 - 6x + 10y + 25 = 0$$

$AC > 0$
 $AC < 0$

$$x^2 - y^2 - 10x + 12y - 16 = 0$$

Circle
 Ellipse
 Hyperbola
 Parabola

9.3 Hyperbole

th Day 2

1/8


657 43, Sound Location

You live 4 miles apart
East-West Street

You hear thunder

Friend hears 18 sec later

Sound travels 1100 ft/sec



On test
Study

$$d_2 - d_1 = 1100 \cdot 18 = 2a$$

$$19800 \Rightarrow a = 9900$$

$$c^2 = a^2 + b^2$$

$$10560^2 = 9900^2 + b^2$$

$$b^2 = 13503600$$

$$b = 3674.72$$

$$\text{path } \frac{x^2}{11513600} + \frac{y^2}{13503600} = 1$$

98010000 ↑ hyperbolas are =

44, Sound Location


3 listening stations Delay

(3300, 0) 0 sec

(3300, 1100) 1 sec

(-3300, 0) 4 sec

sound travels 1100 ft/sec



$$c = 3300$$

$$d_2 - d_1 = 2a = 4 \cdot 1100$$

$$2a = 4400$$

$$a = 2200$$

$$\text{Assume } x = 3300$$

Find y

$$\frac{3300^2}{2200^2} - \frac{y^2}{6050000} = 1$$

$$-1.25 = -\frac{y^2}{6050000}$$

$$y = \sqrt{7562500}$$

$$y = \pm 2750 \text{ } \leftarrow \text{only negative}$$

$$(3300, -2750)$$

The (3300, 1100) point only shows that point is in 4th quad

Classify

① 47. $x^2 + y^2 - 6x + 4y + 9 = 0$ circle
 $\nearrow$ same

① 49. $4x^2 - y^2 - 4x - 3 = 0$ hyperbola
 $AC = -4 < 0$

① 51. $4x^2 + 3y^2 + 8x - 24y + 51 = 0$ ellipse
 $4 \cdot 3 = 12 > 0$

① 53. $25x^2 - 10x - 200y - 119 = 0$ parabola
 $AC = 0$

9.4 Rotations + Systems Notes

1/8

Are Parallel


Parabola Graphing

ex

$$y^2 = 4(x-1)$$

$$y_1 = \sqrt{4(x-1)}$$

$$y_2 = -\sqrt{4(x-1)}$$



Ellipse


$$\frac{(x+3)^2}{4} + \frac{(y-1)^2}{1} = 1$$

Solve for y

$$(y-1)^2 = 1 - \frac{(x+3)^2}{4}$$

$$y-1 = \pm \sqrt{1 - \frac{(x+3)^2}{4}}$$


$$y = \pm \sqrt{1 - \frac{(x+3)^2}{4}} + 1$$



Not Parallel

(slanted)

new B term



$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

Use discriminant to classify conic
($B^2 - 4AC$)

Ellipse or Circle : $B^2 - 4AC < 0$

Parabola : $B^2 - 4AC = 0$

Hyperbola : $B^2 - 4AC > 0$

$$Ox \quad 3x^2 + 7xy + 5y^2 - 6x - 7y + 15 = 0$$

$$A = 3$$

$$B = 7$$

$$C = 5$$

$$B^2 - 4AC$$

$$7^2 - 4(3)(5)$$

$$-11$$

$$-11 < 0$$

① ellipse or circle
 $\uparrow A \neq C$

P1

$$x^2 - 6xy + 9y^2 - 24 + 1 = 0$$

$$A = 1$$

$$B = -6$$

$$C = 9$$

$$(-6)^2 - 4 \cdot 1 \cdot 9$$

$$0$$

$$0 = 0$$

parabola

Solve System of Conics

Elimination or Substitution method

$$\begin{aligned} & \begin{cases} x^2 + y^2 - 16x + 39 = 0 & \rightarrow \text{circle} \\ x^2 - y^2 - 9 = 0 & \rightarrow \text{hyperbola} \end{cases} \\ + & \rightarrow \begin{cases} 2x^2 - 16x + 30 = 0 \end{cases} \end{aligned}$$

$$2(x^2 - 8x + 15)$$

$$2(x-5)(x-3)$$

$$\uparrow x=5 \quad \uparrow x=3$$

$$\uparrow y = 5^2 - 16(5) + 39$$

$$y = \sqrt{16}$$

$$y = \pm 4$$


$$x^2 - y^2 - 9 = 0$$

$$3^2 - y^2 - 9 = 0$$

$$y^2 = 0 \quad (3, 0)$$

$$(5, 4)$$

$$(5, -4)$$



9.4 Notation + Systems of Quadratics

HLW

1/8

666

27.

Classify

$$16x^2 - 24xy + 9y^2 - 30x - 40y \rightarrow 9 + (-24 - 40)y + (16x^2 - 30x) = 0$$

$$A = 16$$

$$B^2 - 4AC$$

$$B = -24$$

$$24^2 - 4(16)(9)$$

$$C = 9$$

$$0$$

parabola

$$(24x + 40) \pm \sqrt{(-24 - 40)^2 - 4(16x^2 - 30x)}$$

$$2 \cdot 9$$

$$-576x^2 + 1080x$$

$$18$$

$$(24x + 40) \pm \sqrt{3000x + 1000}$$

$$1000$$

$$18$$

29.

$$15x^2 - 8xy + 7y^2 - 45 = 0$$

$$A = 15$$

$$B = -8$$

$$C = 7$$

$$8^2 - 4(15)(7)$$

$$-356 < 0$$

ellipse/circle

factoring

$$7y^2 - 8xy + (15x^2 - 45) = 0$$

$$y = 8x \pm \sqrt{64x^2 - 4(7)(15x^2 - 45)}$$

$$8x \pm \sqrt{64x^2 - 420x^2 + 1260}$$

$$14$$

$$y = 8x \pm \sqrt{-356x^2 + 1260}$$

$$14$$

don't have to factor

667

31.

$$x^2 - 6xy - 5y^2 + 4x - 22 = 0$$

$$A = 1$$

$$B = -6$$

$$C = -5$$

$$56$$

hyperbola

don't have to factor

factor

667

33.

$$x^2 + 4x + 4y^2 - 5x - y - 3 = 0$$

$$A = 1$$

$$B = 4$$

$$C = 4$$

0

parabola

r? factoring

don't have to factor

35.


Sketch Degenerate Conic

$$y^2 - 16x^2 = 0$$

$$y^2 = 16x^2$$

$$y = \pm \sqrt{16x^2}$$

$$y = \pm 4x$$



37.

$$x^2 + 2xy + y^2 - 1$$

$$y^2 - x^2 - 2xy + 1$$

$$y = \pm \sqrt{-x^2 - 2xy + 1}$$


$$\rightarrow (x^2 + 2xy + y^2) - 1$$

$$(x+y)^2 = 1$$

$$x+y = \pm 1$$

$$y = x+1$$

$$y = x-1$$



9.4 Rotation + System of Quadratics

Day 2 solving Systems

1/9

Elimination method

667/42

$$\begin{cases} x^2 - 4y^2 - 20x - 64y - 172 = 0 & \text{hyperbola} \\ 16x^2 + 4y^2 - 320x + 64y + 1600 = 0 & \text{ellipse} \end{cases}$$

$$17x^2 - 340x + 1428 = 0$$

factor or quadratic formula

$$(17x - 238)(x - 6)$$

$$x = 14$$

$$x = 6 \rightarrow$$

$$(6)^2 - 4y^2 - 20(6) - 64y - 172 = 0$$

↓ plug in

Plug in

$$(14)^2 - 4y^2 - 20(14) - 64y - 172 = 0$$

$$196 - 4y^2 - 280 - 64y - 172 = 0$$

$$-4y^2 - 64y - 256 = 0$$

$$-4(y^2 + 16y + 64)$$

$$(y + 8)^2$$

$$\uparrow y = -8$$

$$(14, -8)$$

$$36 - 4y^2 - 120 - 64y - 172 = 0$$

$$-4y^2 - 64y - 256 = 0$$

$$-4(y^2 + 16y + 64)$$

$$(y + 8)^2$$

$$\uparrow y = -8$$

$$(6, -8)$$

46.

$$16x^2 - y^2 + 16y - 128 = 0$$

$$y^2 - 48x - 16y - 32 = 0$$

$$16x^2 - 48x - 160 = 0$$

$$16(x^2 - 3x - 10)$$

$$x = 5$$

$$x = -2$$

↓

$$y^2 - 48(5) - 16y - 32 = 0$$

$$y^2 - 16y - 272 = 0$$

$$y = 26.33$$

$$x = -10.33$$

$$(5, 26.33)$$

$$(5, -10.33)$$

$$16 \pm 8\sqrt{21}$$

$$(5, 8 \pm 4\sqrt{21})$$

↓ pick term with most squared things

$$y^2 - 48(-2) - 16y - 32 = 0$$

$$y^2 - 16y + 64 = 0$$

$$(y - 8)^2$$

$$y = 8$$

$$(-2, 8)$$

$$44. \begin{cases} x^2 + 4y^2 - 2x - 8y + 1 = 0 \\ -x^2 + 2x - 4y - 1 = 0 \end{cases}$$

$$4y^2 - 12y = 0$$

$$4y(y-3) = 0$$

$$y = 0 \quad \vee \quad y = 3$$

$$y = 0 \quad x^2 + 4(0)^2 - 2x - 8(0) + 1 = 0$$

$$x^2 - 2x + 1 = 0 \quad x^2 + 36 - 2x - 24 + 1 = 0$$

$$x^2 - 2x + 1 = 0$$

$$(x-1)^2$$

$$(0, 1)$$

$$x^2 - 2x + 13 = 0$$

No real solution

Substitution

$$50. \begin{cases} 4x^2 - y^2 - 8x + 6y - 9 = 0 & (1) \rightarrow 4y^2 - 8x - 9 = y^2 - 6y \\ 2x^2 - 3y^2 + 4x + 18y - 43 = 0 & (2) \end{cases}$$

$$2x^2 - 3y^2 + 4x + 18y - 43 = 0$$

↓

$$2x^2 + 4x - 43 - 3y^2 + 18y$$

$$2x^2 + 4x - 43 - (3y^2 - 18y)$$

$$12x^2 - 24x - 27 = 3y^2 - 18y$$

$$2x^2 + 4x - 43 - (12x^2 - 24x - 27)$$

$$2x^2 + 4x - 43 - 12x^2 + 24x + 27$$

$$-10x^2 + 28x - 16 = 0$$

$$10x^2 - 28x + 16 = 0$$

$$2(5x^2 - 14x + 8)$$

$$x = 2$$

$$x = .8$$

∴ will be no solution

$$x = 3$$

$$48. \quad \left. \begin{aligned} 6x^2 + 3y^2 - 12 &= 0 \\ x + y - 2 &= 0 \end{aligned} \right\} \rightarrow y = 2 - x$$

$$\downarrow$$

$$6x^2 + 3(2-x)^2 - 12$$

$$6x^2 + 3(4 - 2x - 2x + x^2) - 12$$

$$6x^2 + 3(x^2 - 4x + 4) - 12$$

$$6x^2 + 3x^2 - 12x + 12 - 12$$

$$9x^2 - 12x$$

$$3x(3x - 4)$$

↓

$$\rightarrow x = 4/3$$

$$x = 0$$

$$6(0)^2 + 3y^2 - 12 = 0$$

$$3y^2 - 12 = 0$$

$$3y^2 = 12$$

$$y^2 = 4$$

$$y = \pm 2$$

$$(0, \pm 2)$$

$$6(4/3)^2 + 3y^2 - 12 = 0$$

$$6(16/9) + 3y^2 = 12$$

$$32/3 + 3y^2 = 12$$

$$\frac{4}{3} = \frac{3y^2}{3}$$

$$y^2 = 4/3$$

$$y = \pm 2/3$$

$$(4/3, \pm 2/3)$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

$$x + y = 15$$

9.4 Rotation Systems of Quadratics

HW Day 2

1/9

667. 41.

Combine by elimination

$$\begin{cases} -4x^2 - y^2 - 16x + 24y - 16 = 0 \\ 4x^2 + y^2 + 40x - 24y + 208 = 0 \end{cases}$$

$$24x + 192 = 0$$

$$24x = -192$$

$$x = -8 \Rightarrow 4(-8)^2 + y^2 + 40(-8) - 24y + 208$$

$$256 + y^2 - 320 - 24y + 208$$

$$y^2 - 24y + 144$$

$$y = 12$$

$$(-8, 12)$$

①

43.

$$x^2 - y^2 - 12x + 16y - 64 = 0$$

$$x^2 + y^2 - 12x - 16y + 64 = 0$$

$$2x^2 - 24x = 0$$

$$2x(x - 12)$$

$$x = 0$$

$$(0)^2 + y^2 - 12(0) - 16y + 64$$

$$y^2 - 16y + 64$$

$$y = 8$$

$$(0, 8)$$

$$x = 12$$

$$(12)^2 + y^2 - 12(12) + 16y - 64$$

$$144 + y^2 - 144 + 16y - 64$$

$$y^2 + 16y - 64$$

$$y = 8$$

$$(12, 8)$$

45.

$$-16x^2 - y^2 + 24y - 80 = 0$$

$$16x^2 + 25y^2 - 400 = 0$$

$$+ 24y^2 + 24y - 480 = 0$$

$$\text{cancel } 24(y^2 + y - 20)$$

$$x = 4$$

$$x = -5$$

$$\text{try } y = 0$$

$$-16x^2 - (0)^2 + 24(0) - 80 = 0$$

$$-16x^2 - 80 = 0$$

$$-16(x^2 - 5) \rightarrow$$

in
cont

Substitution

49. $10x^2 - 25y^2 - 100x + 160 = 0$

$$y^2 - 2x + 16 = 0 \longrightarrow$$

$$y^2 = 2x - 16 \text{ \textit{e just use}} \\ y = \sqrt{2x - 16}$$

$$10x^2 - 25(2x - 16) - 100x + 160 = 0$$

$$10x^2 - 50x + 400 - 100x + 160 = 0$$

$$10x^2 - 150x + 560 = 0$$

$$10(x^2 - 15x + 56)$$

$$x = 8 \quad \checkmark \quad \text{---} \quad 5x = 7$$

~~$$10(8)^2 - 25y^2 - 100(8) + 160 = 0$$~~

~~$$640 - 25y^2 - 800 + 160 = 0$$~~

~~$$-25y^2 - 320 = 0$$~~

~~$$-5(5y^2 - 64)$$~~

no answer

0?

~~$$10(7)^2 - 25y^2 - 100(7) + 160$$~~

~~$$490 - 25y^2 - 700 + 160$$~~

~~$$-25y^2 - 50 = 0$$~~

~~$$-25(y^2 - 2)$$~~

ending

Use simpler equation

$$(8, 0) \quad \checkmark$$

$$x^2 = 2(8) - 16$$

$$x^2 = 0$$

$$x = 0$$

$$x^2 = 2(7) - 16$$

$$x^2 = -2$$

$$2i$$

9.5 Parametrics

Notes


1/9

to this point - graphs have been represented by a single equation in 2 variables (x, y)
↑ rectangular equations

- can use a 3rd variable to represent a curve in a plane
↑ time

example: an object is propelled into the air at an angle of 45° with an initial velocity of 48 ft/sec
object follows a parabolic path

$$y = \frac{x^2}{72} + x$$



↑ tells you where object is
- doesn't tell you when the object was at a given point (x, y)

SO introduce a 3rd variable
 $t = \text{time}$ $\in \text{parameter}$

Objective $\rightarrow$ write both x and y as a function of t


$$x = f(t)$$

$$y = g(t)$$

A	0	1	2	3	$3\sqrt{2}$
x	0	$24\sqrt{2}$			36
y	0	$-16 + 24\sqrt{2}$			18

$$x = 24\sqrt{2}t$$

$$y = -16t^2 + 24\sqrt{2}t$$



9.5 Parametrics

Notes Day 2

1/10


t is the parameter

ex1 Sketch the curve given by the parametric equation

$$x = t^2 - 4 \quad -2 \leq t \leq 3$$

$$y = \frac{t}{2}$$

t	-2	-1	0	1	2	3
x	0	-3	-4	-3	0	5
y	-1	-0.5	0	1/2	1	3/2




same shape only faster

ex2 2 different sets of parametrics

$$x = 4t^2 - 4$$

$$y = t$$

$$-1 \leq t \leq 3/2$$




t	x	y
-1	0	-1
-1/2	3	-1/2
0	-4	0
1/2	-3	1/2
1	0	1
3/2	5	3/2

Calculator

Set to Radian mode

Set to Parametric mode (Par)

$$\begin{aligned} \text{Put in } X_1 &\rightarrow T \\ Y_1 &\rightarrow T^3 \end{aligned}$$



Graph (might have to adjust window)

$$T_{\min} = -4$$

$$T_{\max} = 4$$

$$T_{\text{step}} = .1$$

* Does matter which is x and y

Convert Parametric to Rectangular

ex 3

$$\begin{aligned} x &= t^2 - 4 \\ y &= \frac{1}{2}t \end{aligned} \quad \begin{array}{l} \text{take one and solve} \\ \text{for } t \end{array} \quad 2y = t$$

$$\begin{aligned} x &= (2y)^2 - 4 \\ x &= 4y^2 - 4 \end{aligned}$$

parabola

ex 4

$$x = \frac{1}{t+1}$$

$$t > -1$$

$$y = \frac{t}{t+1}$$

$$x^2 = \frac{1}{t+1}$$

$$y = \left(\frac{1}{x^2 - 1} \right)$$


$$\frac{1}{x^2} = t+1$$

$$\left(\frac{1}{x^2} - 1 \right) + 1$$

$$\boxed{\left(\frac{1}{x^2} - 1 \right) = t}$$

$$\frac{\frac{1-x^2}{x^2}}{\frac{1}{x^2} + x^2} \Rightarrow$$


$$y = 1 - x^2 \text{ is parabolic + vertex} = (0, 1)$$



673
8.

$$x = 4 \cos^2 \theta \quad y = 2 \sin \theta$$

θ	$-\frac{\pi}{2}$	$-\frac{\pi}{4}$	0	$\frac{\pi}{4}$	$\frac{\pi}{2}$
x	0	2	4	2	0
y	-2	$-\sqrt{2}$	0	$\sqrt{2}$	2



Convert to rectangular

$$x = 4 \cos^2 \theta$$

$$\frac{x}{4} = \cos^2 \theta$$



$$\cos^2 \theta + \sin^2 \theta = 1$$

$$y = 2 \sin \theta$$

$$\frac{y}{2} = \sin \theta$$

$$\frac{y^2}{4} = \sin^2 \theta$$

$$\frac{x}{4} + \frac{y^2}{4} = 1 \quad \text{parabola}$$

674

16.

$$x = \sqrt{t}$$

$$x^2 = t$$

$$y = 1 - t$$

add restriction
 $t \geq 0$

$$y = 1 - x^2$$

parabola

18.


$$x = |t - 1|$$

$$x = t + 2$$

$$y - 2 = t$$

$$x = |(y - 2) - 1|$$

$$x = |y - 3|$$



22.

$$x = e^{2t}$$


$$y = e^t$$

$$y^2 = e^{2t}$$

$$x = y^2$$

$$e^t > 0$$

$$y > 0$$



24.

$$x = \ln 2t$$

$$x = \log_e 2t$$

$$e^x = 2t$$

$$t = \frac{e^x}{2}$$

$$y = 2t^2$$

$$y = 2 \left(\frac{e^x}{2} \right)^2$$

$$y = 2 \left(\frac{1}{4} e^{2x} \right)$$

$$y = \frac{1}{2} e^{2x}$$

P1

$$y = 4 \sin \theta$$

$$\frac{y}{4} = \sin \theta$$

$$\frac{y^2}{16} = \sin^2 \theta$$

$$\frac{x}{3} = \cos \theta$$


$$\frac{x^2}{9} = \cos^2 \theta$$

$$\frac{x^2}{9} = \cos^2 \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\frac{y^2}{16} + \frac{x^2}{9} = 1$$

ellipse



$$26. \quad x = 4 + 2\cos\theta$$

$$x - 4 = 2\cos\theta$$

$$\frac{x-4}{2} = \cos\theta$$

$$\frac{(x-4)}{2} = \cos\theta$$

$$y = -1 + 2\sin\theta$$

$$y + 1 = 2\sin\theta$$

$$\frac{y+1}{2} = \sin\theta$$

$$\frac{(y+1)}{2} = \sin\theta$$

$$\left(\frac{x-4}{2}\right)^2 + \left(\frac{y+1}{2}\right)^2 = 1$$

$$\frac{(x-4)^2}{4} + \frac{(y+1)^2}{4} = 1 \quad \in \text{circle} \quad \text{Coefficients} =$$

28,

$$x = \sec\theta$$

$$x^2 = \sec^2\theta$$

$$y = \tan\theta$$

$$y^2 = \tan^2\theta$$

$$\tan^2\theta + 1 = \sec^2\theta$$

$$y^2 + 1 = x^2$$

$$1 = x^2 - y^2 \in \text{hyperbole}$$

Rectangular to Parametric

Find a set of parametric equations to represent the graph of $y = 1 - x^2$ with $t = x$

$$y = 1 - t^2 \quad \sim \quad \text{pair } \odot$$

with $t = 1 - x$ instead

$t - 1 = -x$ solve for x

$$\boxed{1 - t = x} \rightarrow y = 1 - (1 - t)^2$$

$$y = 1 - (1 - 2t + t^2)$$

$$y = 2t - t^2$$

44, $y = 5 - 7x$ (no starting parametric)

so start with $t = x$

$y = 5 - 7t$
or $x = t$

so $y = 5 - 7t$

48 $y = 6x^2 - 5$

try $t = x$

$y = 6(t)^2 - 5$

9.5 Parametric Equations


Day 2 HW

1/10

673

①


3. $x = \sqrt{t}$ $y = t$ $x = \sqrt{y}$ $y = x^2$ $x \geq 0$ (b)



674

①

5. $x = \ln t \Rightarrow e^x = t$
 $y = \frac{1}{2}t - 2 \Rightarrow y = \frac{1}{2}(e^x) - 2$ (f)




t continues (blue line)

7. $x = \sqrt{t}$ $x^2 = t$
 $y = 2 - t \Rightarrow y = 2 - x^2$

a)

t	0	1	2	3	4
x	0	1	$\sqrt{2}$	$\sqrt{3}$	2
y	2	1	0	-1	-2


$y = 2 - x^2$



Suppose to be the same if this is $x \geq 0$

13.

$x = \frac{1}{4}t \Rightarrow 4x = t$
 $y = t^2 \Rightarrow y = (4x)^2 = 16x^2$




t	x	y
-1	-1/4	1
0	0	0
1	1/4	1
2	1/2	4
3	3/4	9

674


15.

$x = t + 2 \Rightarrow t = x - 2$
 $y = t^2 \Rightarrow y = (x - 2)^2$
 $y = x^2 - 4x + 4$



t	x	y
-1	1	1
0	2	0
1	3	1
2	4	4
3	5	9
4	6	16

17. $x = 2t \Rightarrow t = \frac{x}{2}$
 $y = |t - 2| \Leftrightarrow y = \left| \frac{x}{2} - 2 \right| = \frac{1}{2} |x - 4|$




t	x	y
-2	-4	4
-1	-2	3
0	0	2
1	2	1
2	4	0
3	6	1

19. $x = 3 \cos \theta$ $y = 3 \sin \theta$
 $\frac{x}{3} = \cos \theta$ $\frac{y}{3} = \sin \theta$

$\frac{x^2}{9} = \cos^2 \theta$ $\frac{y^2}{9} = \sin^2 \theta$


$\cos^2 \theta + \sin^2 \theta = 1$

$\frac{x^2}{9} + \frac{y^2}{9} = 1$




t	x	y
0	3	0
1	0	3
2	-3	0
3	0	-3

21. $x = e^{-t}$ $y = e^{3t}$
 $\frac{1}{x} = e^+ \rightarrow y = (e^+)^3 \rightarrow \left(\frac{1}{x}\right)^3 \rightarrow \frac{1}{x^3}$



23. $x = t^3$ $y = 3 \ln t$
 $x = \sqrt[3]{y^3} = y$
 $x = y^e$



t	x	y
-1	-1	(X)
0	0	(X)
1	1	0
2	8	2.07
3	27	3.30
4	64	5.16


674

Graph with calculator

25.

$$x = -4 + 2 \cos \theta$$


$$y = -1 + \sin \theta$$



27.

$$x = 4 \sec \theta$$


$$y = 3 \tan \theta$$



29

$$x = t/2$$

$$y = \ln(t^2 + 1)$$



43. Find 2 parametric equations

$$y = 4x - 3$$

$$\textcircled{1} \begin{aligned} x &= t \\ y &= 4t - 3 \end{aligned}$$

$$x = \frac{1}{4}t$$

$$y = t - 3$$

45.

$$y = \frac{1}{x}$$

$$\begin{aligned} x &= t \\ y &= \frac{1}{t} \end{aligned}$$

$$\begin{aligned} x &= 5t \\ y &= \frac{5}{t} \end{aligned}$$

47.

$$y = x^2 + 4$$

$$\begin{aligned} x &= t \\ y &= t^2 + 4 \end{aligned}$$

$$\begin{aligned} x &= \sqrt{t} \\ y &= t + 4 \end{aligned}$$

49.

$$y = x^3 + 2x$$

$$\begin{aligned} x &= t \\ y &= t^3 + 2t \end{aligned}$$

$$\begin{aligned} x &= 10t \text{ or } \frac{x}{10} = t \\ y &= (10t)^3 + 20t \end{aligned}$$


Graph

53.

Witch of Agnesi

$$x = 2 \cot \theta$$

$$y = 2 \sin^2 \theta$$



675

59.

Projectile Motion

$$x = (v, \cos \theta)t$$

$$y = h + (v, \sin \theta)t - 16t^2$$

(center field fence = 10 ft high + 400 ft from home plate)

Ball hit when 3 ft above ground @ 100 mph @ θ angle

146.46 feet/sec


a) Write Parametric Equation

$$x = (146.46 \cos \theta)t$$

$$y = 3 + (146.46 \sin \theta)t - 16t^2$$

or set = to
400 to
find t

b) 15°



No

$$[146.46 \cos(15)]t$$

$$141.7t = x$$

$$3 + (146.46 \sin 15)t - 16t^2$$


$$3 + 38t - 16t^2 = y$$

what's
 θ so now

put

when x =
400
height?

c) 23°



Yes

$$400 = 146.46 \cos 23 t$$

$$t = 2.96 \text{ sec}$$

$$y = 3 + (146.46 \sin 23)(2.96) - 16(2.96)^2$$

$$x$$

$$y = 21.39$$

$$y > 0$$

d) Start by solving for t = $\frac{x}{146.46 \cos \theta}$

$$\text{Sub in } 3 + 146.46 \sin \theta \left(\frac{x}{146.46 \cos \theta} \right) - 16 \left(\frac{x}{146.46 \cos \theta} \right)^2$$

Rewrite
in terms of
tan

$$y = 3 + \tan \theta x - \frac{16 \sec^2 \theta}{(146.67)^2} x^2$$

$$y = 3 + \tan \theta x - \frac{16 (1 + \tan^2 \theta)}{146.67^2} x^2$$

Sub in
problem

$$10 = 3 + \tan \theta x - \frac{16 (1 + \tan^2 \theta)}{146.67^2} 400^2$$

$$10 = 3 + 400 \tan \theta - 119.0 \tan^2 \theta - 119$$

Quadratic
formula

$$119 \tan^2 \theta - 400 \tan \theta + 126 = 0$$

$$\tan \theta = .35 \rightarrow 19.9^\circ \quad \tan \theta = 3.01 \rightarrow 71.68^\circ$$

want min

9/ Conics + Parametrics

Test Topics

1/15

1. Determine all parts of hyperbole, ellipse, and parabola then graph
2. Finding equation of a conic given foci, length of axis, etc.
3. Determine type of conic using $A \cdot C$ parallel
 $B^2 - 4AC$ non parallel
4. Find equation of line tangent to parabola,
5. Systems of Conics
6. Given parametric equations, get rectangular equations
7. Given rectangular equations come up with parametric equation.
8. Word problems
 - going thru a tunnel
 - hearing a sound
 - hyperbolic mirror

Just Jones

(Circles of Influence)

1/15

1. Influence and Goals of Stakeholders

2. Understanding of a team's role

3. Defining the team's mission

4. Finding solutions to the team's problems

5. Defining the team's structure

6. Creating a supportive environment

7. Encouraging team members to work together

8. Working together

9. Doing the work

10. Evaluating the work


11. Reporting the work

9 Analytic Geometry Review

1/15


699 Find the equation of the tangent line

7. $x^2 = -2y$ $\leftarrow$ $4p = -2$
 $(2, -2)$ $p = -\frac{1}{2}$
 vertex $(0, 0)$



$d_1 = \sqrt{(0-2)^2 + (-\frac{1}{2}-2)^2}$
 $\sqrt{4 + 6.25}$
 2.5
 $d_2 = d_1 = 2.5$
 $\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$
 $\frac{y - (-2)}{-2 - (-2)} = \frac{x - 2}{0 - 2}$
 $\frac{y + 2}{0} = \frac{x - 2}{-2}$
 $y + 2 = -2(x - 2)$
 $y + 2 = -2x + 4$
 $y = -2x + 2$

9. $2x = ?$




$x^2 = 4p(y - 12)$
 $16 = 4p(10 - 12)$
 $16 = 4p(-2)$
 $16 = -8p$
 $p = -2$
 $x^2 = 4(-2)(y - 12)$
 $x^2 = -8(y - 12)$
 $x^2 = -8y + 96$
 $x = \sqrt{96 - 8y}$
 $x = 9.79 \cdot 2 = 19.59$

13. Find standard form of ellipse

$(0, 1)$ minor points $(2, 0)$ $eb = 1$
 $(4, 1)$ $a = 2$ $(2, 2)$ x axis major $\bigcirc$
 vertex $(2, 1)$

$\frac{(x - 2)^2}{2^2} + \frac{(y - 1)^2}{1^2} = 1$
 $\frac{(x - 2)^2}{4} + \frac{(y - 1)^2}{1} = 1$

15. Architecture



switch

$$b = 4$$

$$\text{Foci} = ?$$

$$c = ?$$

$$c = ?$$

$$2b = 10$$

$$a = 5$$

$$c^2 = a^2 - b^2$$

$$c^2 = 5^2 - 4^2$$

$$c^2 = 9$$

$$c = 3$$

$$17. \quad 16x^2 + 9y^2 - 32x + 72y + 16$$

$$16x^2 - 32x + 9y^2 + 72y = -16$$

$$16(x^2 - 2x + 1) + 9(y^2 + 8y + 16) = -16 + 16 + 144$$

$$16(x-1)^2 + 9(y+4)^2 = 144$$

$$\frac{(x-1)^2}{9} + \frac{(y+4)^2}{16} = 1$$

$$\frac{(x-1)^2}{9} + \frac{(y+4)^2}{16} = 1$$

$$\text{Center } (1, -4)$$

$$a = 4$$

$$\text{major } (1, -4 \pm 4)$$

$$b = 3$$

$$\text{minor } (1 \pm 3, -4)$$

$$c^2 = a^2 - b^2$$


$$c^2 = 4^2 - 3^2$$

$$c = \sqrt{7}$$

$$c = 2.65$$

$$\text{foci } (1, -4 \pm \sqrt{7})$$

$$e = \frac{c}{a} = \frac{\sqrt{7}}{4}$$



$$19. \quad 3x^2 + 8y^2 + 12x - 112y + 403 = 0$$

$$3x^2 + 12x + 8y^2 - 112y = -403$$

$$3(x^2 + 4x + 4) + 8(y^2 - 14y + 49) = -403 + 12 + 392$$

$$3(x+2)^2 + 8(y-7)^2 = 1$$

$$\frac{(x+2)^2}{1/3} + \frac{(y-7)^2}{1/8} = 1 \quad \text{ex axis}$$

$$\text{center } (-2, 7)$$

$$a^2 = 1/3 \Rightarrow \frac{1}{\sqrt{3}} \Rightarrow \frac{\sqrt{3}}{3}$$

$$b^2 = 1/8 \Rightarrow \frac{1}{\sqrt{8}} \Rightarrow \frac{\sqrt{8}}{8}$$

$$c^2 = a^2 - b^2$$

$$1/3 - 1/8$$

$$c = \sqrt{5/24} = \frac{\sqrt{5}}{2\sqrt{6}}$$

$$\text{major } (-2 \pm \frac{\sqrt{3}}{3}, 7)$$

$$\text{minor } (-2, 7 \pm \frac{\sqrt{2}}{4}) \quad e = \frac{\sqrt{2}}{4}$$

$$\text{Focus } (-2 \pm \frac{\sqrt{5}}{2\sqrt{6}}, 7)$$

$$\nearrow \sqrt{30}/12$$

699

Find standard form of hyperbole

23. Foci (0,0) & vertex (4,0)

(8,0) $c=4$ $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

asy = $y = \pm 2(x-4)$

slope = $\frac{b}{a}$

$b=2a$

$b^2=4a^2$

$a^2+b^2=c^2$

$a^2+4a^2=16$

$5a^2=16$

$5a^2=16$

$a^2=\frac{16}{5}$

$\frac{16}{5} + b^2 = 16$

$b^2 = 12\frac{8}{5}$

$6\frac{4}{5}$

remember vertex

$(x-4)^2$

$\frac{x^2}{\frac{16}{5}} - \frac{y^2}{\frac{64}{5}} = 1$

700

25. $9x^2 - 16y^2 - 18x - 32y - 151 = 0$

$9x^2 - 18x - 16y^2 - 32y = 151$

$9(x^2 - 2x + 1) - 16(y^2 + 2y + 1) = 151 + 9 - 16$

$9(x-1)^2 - 16(y+1)^2 = 144$ remember!

$\frac{(x-1)^2}{16} - \frac{(y+1)^2}{9} = 1$

$\frac{(x-1)^2}{16} - \frac{(y+1)^2}{9} = 1$

$a^2=16$

$b^2=9$

$a=4$

$b=3$

$c^2=a^2+b^2$

$c^2=16+9$

$c=\sqrt{25}$

$c=5$

asymptotes $y+1 = \pm \frac{b}{a}(x-1)$

$y = \pm \frac{3}{4}(x-1) - 1$


vertex (1, -1)

major (1 ± 4, -1)

minor (1, -1 ± 3)


foci (1 ± 5, -1)

$e = \frac{c}{a} = \frac{5}{4}$



701

29.



travels 186,000 miles per sec

A gets 10065 sec sooner

How far north is ship?

$\frac{60^2}{120^2} + \frac{y^2}{b^2}$

$186,000 \cdot 0.0005 = 93 \text{ miles} = 2a$

$a = 93/2$

$2a = 240$

$a = 120$

$2c = 200$

$c = 100$

$100^2 = \left(\frac{93}{2}\right)^2 + b^2$

$A=C$ circle
 $AC=0$ parabola
 $AC>0$ ellipse
 $AC<0$ hyperbola

Classify ✓

33. $5x^2 - 2y^2 + 10x - 4y + 17 = 0$

① $5 \cdot 10 = 50$
 $50 - 20 = 30$
 $30 < 0$ hyperbola

39. $16x^2 - 8xy + y^2 - 10x + 5y = 0$

① $16 = A$
 $-8 = B$
 $1 = C$
 $(-8)^2 - 4(16)(1) = 0$ parabola

$y^2 + (-8x + 5)y + (16x^2 - 10x) = 0$ Solve for y

$$y = \frac{-(8x - 5) \pm \sqrt{(8x - 5)^2 - 4(1)(16x^2 - 10x)}}{2(1)}$$

$$y = \frac{-(8x - 5) \pm \sqrt{64x^2 + 25 - 80x - 64x^2 + 40x}}{2}$$

41. $x^2 + y^2 + 2xy + 2\sqrt{2}x - 2\sqrt{2}y + 2 = 0$

① $A=1$
 $B=2$
 $C=1$
 $(2)^2 - 4 \cdot 1 \cdot 1 = 0$ parabola

43. Solve

$-4x^2 - y^2 - 32x + 24y - 64 = 0$

$4x^2 + y^2 + 66x - 24y + 304 = 0$

$24x + 240 = 0$

$24x = -240$

$x = -10$

find y $-4(-10)^2 - y^2 - 32(-10) + 24y - 64 = 0$

$-400 - y^2 + 320 + 24y - 64 = 0$

$-y^2 + 24y - 144 = 0$

$y = 12$
 $(-10, 12)$

44, $x^2 + y^2 - 25 = 0 \Rightarrow y^2 = -x^2 + 25$
 $9x - 4y^2 = 0 \Rightarrow 9x - 4(-x^2 + 25) = 0$
 $4x + 4x^2 - 100 = 0$
 $4x^2 + 9x - 100 = 0$

$x = 4$
 $9(4) - 4y^2 = 0$
 $-36 - 36$

$-4y^2 = -36$

$y^2 = 9$

$y = 3$

$(4, 3)$

$x = -6.25$

$9(-6.25) - 4y^2 = 0$
 $+56.25 + 56.25$


$-4y^2 = 56.25$

$y^2 = -14.0625$

non real

47, $x = 1 + 4t$
 $y = 2 - 3t$


t	1	2	3	4	5
x	5	9	13	17	21
y	-1	-4	-7	-10	-13



701

51, $x = 6 \cos \theta$
 $y = 6 \sin \theta$


t	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π
x	4.24	0	-4.24	-6
y	4.24	6	4.24	0



53. $x = \sec \theta$

$y = \tan \theta$


θ	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π
x	1	X	-1	-1
y	1	X	-1	0



54. $x = 2 - \sin \theta$

$y = 2 - \cos \theta$

θ	$\frac{\pi}{4}$	$\frac{\pi}{2}$	$\frac{3\pi}{4}$	π
x	1.29	1	1.29	2
y	1.29	2	2.70	3



703

5.

Find standard form of parabola

Focus $(8, -2)$

Directrix $x = 4$

center $(6, -2)$

$p = 2$

$y^2 = 4px$

$(y-k)^2 = 4p(x-h)^2$

$(y+2)^2 = 4p(x-6)^2$


$(y+2)^2 = 4(2)(x-6)^2$

$(y+2)^2 = 8(x-6)^2$

Everything You Ever Wanted to Know about the Conic Sections

Use the following tables to guide you on your road to mathematical dominance of conic sections.

Circle Centered at (h, k)	
Standard Form Equation	$(x - h)^2 + (y - k)^2 = r^2$
Center	(h, k)
Radius	r

Ellipse Centered at the (h, k) , with $a > b$		
Standard Form Equation	$\frac{(x - h)^2}{a^2} + \frac{(y - k)^2}{b^2} = 1$	$\frac{(y - k)^2}{a^2} + \frac{(x - h)^2}{b^2} = 1$
Center	(h, k)	(h, k)
Major Axis Vertices	$(h \pm a, k)$ 	$(h, k \pm a)$ 
Minor Axis Vertices	$(h, k \pm b)$	$(h \pm b, k)$
Foci	$(h \pm c, k)$	$(h, k \pm c)$
Eccentricity	$e = \frac{c}{a}$	$e = \frac{c}{a}$
$a^2 = b^2 + c^2$		

Parabola Centered at (h, k)		
Standard Form Equation	$(x - h)^2 = 4p(y - k)$	$(y - k)^2 = 4p(x - h)$
Vertex	(h, k) 	(h, k) 
Axis of Symmetry	$x = h$	$y = k$
Focus	$(h, k + p)$	$(h + p, k)$
Directrix	$y = k - p$	$x = h - p$
Eccentricity	$e = \frac{c}{a}$	$e = \frac{c}{a}$

Hyperbola Centered at (h, k)		
Standard Form Equation	$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$	$\frac{(y-k)^2}{a^2} - \frac{(x-h)^2}{b^2} = 1$
Center	(h, k)	(h, k)
Transverse Axis Vertices	$(h \pm a, k)$	$(h, k \pm a)$
Conjugate Axis Vertices	$(h, k \pm b)$	$(h \pm b, k)$
Foci	$(h \pm c, k)$	$(h, k \pm c)$
Asymptotes	$(y-k) = \pm \frac{b}{a}(x-h)$	$(y-k) = \pm \frac{a}{b}(x-h)$
Eccentricity	$e = \frac{c}{a}$	$e = \frac{c}{a}$
$a^2 + b^2 = c^2$		

Conic Section	Equation	Slope of Tangent at $P(x_0, y_0), y_0 \neq 0$
Circle	$x^2 + y^2 = r^2$	$-\frac{x_0}{y_0}$
Ellipse	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b$ $\frac{y^2}{a^2} + \frac{x^2}{b^2} = 1, a > b$	$-\frac{b^2 x_0}{a^2 y_0}$ $-\frac{a^2 x_0}{b^2 y_0}$
Hyperbola	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$	$\frac{b^2 x_0}{a^2 y_0}$ $-\frac{a^2 x_0}{b^2 y_0}$
Parabola	$y^2 = 4cx$ $x^2 = 4cy$	$\frac{2c}{y_0}$ $\frac{x_0}{2c}$

Reciprocal Identities

$$\sin x = \frac{1}{\csc x} \quad \sec x = \frac{1}{\cos x} \quad \tan x = \frac{1}{\cot x}$$

$$\csc x = \frac{1}{\sin x} \quad \cos x = \frac{1}{\sec x} \quad \cot x = \frac{1}{\tan x}$$

Tangent Identities

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x}$$

Pythagorean Identities

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \tan^2 x = \sec^2 x$$

$$1 + \cot^2 x = \csc^2 x$$

Cofunction Identities

$$\sin\left(\frac{\pi}{2} - x\right) = \cos x \quad \cos\left(\frac{\pi}{2} - x\right) = \sin x$$

$$\csc x\left(\frac{\pi}{2} - x\right) = \sec x \quad \tan\left(\frac{\pi}{2} - x\right) = \cot x$$

$$\sec\left(\frac{\pi}{2} - x\right) = \csc x \quad \cot\left(\frac{\pi}{2} - x\right) = \tan x$$

Negative Angle Identities

$$\sin(-x) = -\sin x \quad \cos(-x) = \cos x$$

$$\csc(-x) = -\csc x \quad \tan(-x) = -\tan x$$

$$\sec(-x) = \sec x \quad \cot(-x) = -\cot x$$

Sum and Difference Formulas

$$\sin(u \pm v) = \sin u \cos v \pm \cos u \sin v$$

$$\cos(u \pm v) = \cos u \cos v \mp \sin u \sin v$$

$$\tan(u \pm v) = \frac{\tan u \pm \tan v}{1 \mp \tan u \tan v}$$

Double Angle Formulas

$$\sin 2x = 2 \sin x \cos x$$

$$\cos 2x = \cos^2 x - \sin^2 x = 2 \cos^2 x - 1 = 1 - 2 \sin^2 x$$

$$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$$

Power Reducing

$$\sin^2 u = \frac{1 - \cos 2u}{2}$$

$$\cos^2 u = \frac{1 + \cos 2u}{2}$$

$$\tan^2 u = \frac{1 - \cos 2u}{1 + \cos 2u}$$

6 Circular function definitions where θ is any angle

$$\sin \theta = \frac{y}{r}$$

$$\csc \theta = \frac{r}{y}$$

$$\cos \theta = \frac{x}{r}$$

$$\sec \theta = \frac{r}{x}$$

$$\tan \theta = \frac{y}{x}$$

$$\cot \theta = \frac{x}{y}$$

$$r = \sqrt{x^2 + y^2}$$

CHART OF THE 3 TRIG FUNCTIONS OF SPECIAL ANGLES

Degrees	Radians	$\sin \theta$	$\cos \theta$	$\tan \theta$
0°	0	0	1	0
30°	$\frac{\pi}{6}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
45°	$\frac{\pi}{4}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
60°	$\frac{\pi}{3}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$

Additional Trigonometric Formulas/Identities

Half - Angle Identities:

$$\sin^2 u = \frac{1 - \cos 2u}{2}$$

$$\cos^2 u = \frac{1 + \cos 2u}{2}$$

$$\tan^2 u = \frac{1 - \cos 2u}{1 + \cos 2u}$$

Half - Angle Formulas:

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$$

Alternative Half - Angle Formulas for Tangent:

$$\tan \frac{\theta}{2} = \frac{1 - \cos \theta}{\sin \theta}$$

$$\tan \frac{\theta}{2} = \frac{\sin \theta}{1 + \cos \theta}$$

Product - to - Sum Formulas:

$$\sin u \cos v = \frac{1}{2} [\sin(u + v) + \sin(u - v)]$$

$$\cos u \sin v = \frac{1}{2} [\sin(u + v) - \sin(u - v)]$$

$$\cos u \cos v = \frac{1}{2} [\cos(u + v) + \cos(u - v)]$$

$$\sin u \sin v = \frac{1}{2} [\cos(u - v) - \cos(u + v)]$$

Sum - to - Product Formulas:

$$\sin u + \sin v = 2 \sin \left(\frac{u + v}{2} \right) \cos \left(\frac{u - v}{2} \right)$$

$$\sin u - \sin v = 2 \cos \left(\frac{u + v}{2} \right) \sin \left(\frac{u - v}{2} \right)$$

$$\cos u + \cos v = 2 \cos \left(\frac{u + v}{2} \right) \cos \left(\frac{u - v}{2} \right)$$

$$\cos u - \cos v = -2 \sin \left(\frac{u + v}{2} \right) \sin \left(\frac{u - v}{2} \right)$$

IS Test

1/16

1. Sum or difference identity to find exact value (2)

2. Determine exact value of trig expressions given $\sin v$ and $\cos v$ and a restriction on the quadrant

Don't have
to do

power 3. Use double or half angle identities to find exact
reducing value of trig expression (2)

4. Express sum as a product and give exact value

Have 5. Solving trig equations that have identities in them
formulas (sum identities, double angle) - etc

6. Law of sines + cosines (4 word problems)

7. Herons area formula

13.14

1/10

1. The first part of the problem is to find the value of x such that $x^2 + 1 = 0$.

2. The second part is to find the value of y such that $y^2 + 1 = 0$.

3. The third part is to find the value of z such that $z^2 + 1 = 0$.

4. The fourth part is to find the value of w such that $w^2 + 1 = 0$.

5. The fifth part is to find the value of v such that $v^2 + 1 = 0$.

6. The sixth part is to find the value of u such that $u^2 + 1 = 0$.

7. The seventh part is to find the value of t such that $t^2 + 1 = 0$.

8. The eighth part is to find the value of s such that $s^2 + 1 = 0$.

9. The ninth part is to find the value of r such that $r^2 + 1 = 0$.

10. The tenth part is to find the value of q such that $q^2 + 1 = 0$.

11. The eleventh part is to find the value of p such that $p^2 + 1 = 0$.

12. The twelfth part is to find the value of o such that $o^2 + 1 = 0$.

13. The thirteenth part is to find the value of n such that $n^2 + 1 = 0$.

14. The fourteenth part is to find the value of m such that $m^2 + 1 = 0$.

15. The fifteenth part is to find the value of l such that $l^2 + 1 = 0$.

16. The sixteenth part is to find the value of k such that $k^2 + 1 = 0$.

17. The seventeenth part is to find the value of j such that $j^2 + 1 = 0$.

18. The eighteenth part is to find the value of i such that $i^2 + 1 = 0$.

19. The nineteenth part is to find the value of h such that $h^2 + 1 = 0$.

20. The twentieth part is to find the value of g such that $g^2 + 1 = 0$.

B4 Practice

1/10

1. $\frac{-1 + \frac{\sqrt{3}}{3}}{1 + \frac{\sqrt{3}}{3}}$ common denom $\rightarrow \frac{-\frac{3}{3} + \frac{\sqrt{3}}{3}}{\frac{3}{3} + \frac{\sqrt{3}}{3}} \rightarrow \frac{\frac{\sqrt{3}-3}{3}}{\frac{\sqrt{3}+3}{3}} \rightarrow \frac{\sqrt{3}-3}{3} \cdot \frac{3}{\sqrt{3}+3}$

$\frac{3\sqrt{3}-3}{3\sqrt{3}+3} \rightarrow \frac{\sqrt{3}-3}{\sqrt{3}+3} \cdot \frac{(\sqrt{3}-3)}{(\sqrt{3}-3)} = \frac{3-3\sqrt{3}-3\sqrt{3}+9}{3+3\sqrt{3}-3\sqrt{3}-9}$

$\frac{3-6\sqrt{3}+9}{-6} \rightarrow \frac{12-6\sqrt{3}}{-6} \rightarrow \boxed{-2+\sqrt{3}}$ $\leftarrow$ proper answer

2. Express the sum $\sin \frac{7\pi}{12} + \sin \frac{\pi}{12}$ as a product

$\sin u + \sin v = \sin(u+v)$

~~$\sin u \cos v + \cos u \sin v$~~

not sum + diff formula $\sin\left(\frac{7\pi}{12}\right) \cos\left(\frac{\pi}{12}\right) + \cos\left(\frac{7\pi}{12}\right) \sin\left(\frac{\pi}{12}\right)$

$\sin \frac{7\pi}{12} = 105^\circ$ $\sin \frac{\pi}{12} = 15^\circ$
 $60+45^\circ$ $60-45^\circ$

$\sin(u+v) + \sin(u-v) = 2\sin u \cos v$

$\sin\left(\frac{\pi}{3} + \frac{\pi}{4}\right) + \sin\left(\frac{\pi}{3} - \frac{\pi}{4}\right)$

$\sin \frac{\pi}{3} \cos \frac{\pi}{4} = \sqrt{6}$

$\boxed{2 \cdot \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} = \frac{\sqrt{6}}{2}}$

3. Find the exact value of $\sin\left(\frac{\pi}{12}\right) = 15^\circ$

$60-45^\circ$
 $\frac{\pi}{3} - \frac{\pi}{4}$

$\sin u - v = \sin u \cos v - \cos u \sin v$

$\sin 60^\circ \cos 45^\circ - \cos 60^\circ \sin 45^\circ$

$\frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2} = \frac{\sqrt{6}-\sqrt{2}}{4}$

4. Find exact value of $\sin(u+v)$

$\sin u = \frac{4}{5} \quad 0 < u < \frac{\pi}{2} \rightarrow 53.13^\circ$ ✓ limit

$\cos v = -\frac{12}{13} \quad \frac{\pi}{2} < v < \pi \rightarrow 157.38^\circ$ ✓ limit


$$\sin(u+v) = \sin u \cos v + \cos u \sin v$$

$$\frac{4}{5} \cdot -\frac{12}{13} + \cos(53.13^\circ) \sin(157.38^\circ)$$

~~$\cos(53.13^\circ) = 3/5$~~ ~~$\sin(157.38^\circ) = 5/13$~~

Do this

Find $\downarrow$
 $5/13$



Find $\rightarrow 3$

Use graph

$$-\frac{48}{65} + \frac{15}{65} = -\frac{33}{65}$$

5. Evaluate using double angle + half angle

$\cos \theta = \frac{5}{13} \quad \frac{3\pi}{2} < \theta < 2\pi \rightarrow \boxed{474^\circ}$
exact value

$$\cos 2\theta = 2\cos^2 \theta - 1$$

$$2\left(\frac{5}{13}\right)^2 - 1 = \frac{50}{169} - \frac{169}{169} = -\frac{119}{169}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}} = \pm \sqrt{\frac{1 + 5/13}{2}} = \pm \sqrt{\frac{18/13}{2}} = \pm \sqrt{\frac{9}{13}}$$

$$\pm \frac{\sqrt{9}}{\sqrt{13}} = \pm \frac{3}{\sqrt{13}} = \pm \frac{3\sqrt{13}}{13} \in 4^{\text{th}} \text{ quad } (+)$$

6. Find all solutions of $\cos x = 0$ for $[0, 2\pi)$

$$2 - \sin^2 x = 2 \cos^2 \left(\frac{x}{2}\right)$$

$$2 - (1 - \cos^2 x) = 2 \left(\frac{1 + \cos x}{2}\right)^2$$

$$2 - 1 + \cos^2 x = 2 \left(\frac{1 + \cos x}{2}\right)^2$$

$$\cos^2 x + 1 = 1 + \cos x$$

$$\cos^2 x - \cos x + 0 = 0$$

Roots $x = 0$
 $x = 1$

$\cos x = 0$
 $\cos^{-1} \cos^{-1}$
 $\frac{\pi}{2}, \frac{3\pi}{2}$

$\cos x - 1 = 0$

$\cos x = 1$
 $\cos \cos$

$x = 0$

~~2π~~ $\in$ not included

7. Two planes leave DC, at same time


Plane 1 $\rightarrow$ 425 mph 355° bearing

Plane 2 $\rightarrow$ 530 mph 67° bearing

Distance after 2 hrs $\uparrow$

$$\rightarrow .2 = 850 \text{ miles}$$

$$\rightarrow .2 = 1060 \text{ miles}$$



$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$850^2 + 1060^2 - 2(850)(1060) \cos 72$$


$$c^2 = 1289251.376$$

$$1135.45 \text{ miles}$$

8. Pilot landing on a 9000 ft runway

Angle of depression at ends of runway $= 17.5^\circ + 18.8^\circ$

Find air distance plane must travel before touching beginning of runway



$$\frac{\sin 1.3}{9000} = \frac{\sin 17.5}{x} = 119289 \text{ feet}$$

Or get height
 $\sin 20.5 = \frac{h}{31} = 10.85$
 if $h < B < A \rightarrow 2 \text{ cases}$

9. 2 solutions?

$$a = 12$$

$$b = 31$$

$$A = 20.5^\circ$$

$$B / \frac{\sin 20.5}{12} = \frac{\sin B}{31} \quad B_1 = 64.78^\circ$$

$$C / \rightarrow 180 - 20.5 - 64.78 = 94.72^\circ$$

$$\text{Test } B_2 / 180 - 64.78 = 115.22$$


$$C_2 / 180 - 20.5 - 115.22 = 44.28^\circ$$

If another
 $A + (180 - B) < 180$

$$c_1 / \frac{\sin 20.5}{12} = \frac{\sin 94.72}{c} \quad c = 34.15$$

$$c_2 / \frac{\sin(20.5)}{12} = \frac{\sin 44.28}{c} \quad c = 23.922$$


10.



$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\cos A = 0.376, 53125$$

$$A = 67.91, 57.91$$



$$\sin 67.91 = \frac{h}{6}$$

$$h = 5.56 = h$$

$$5.08$$

$$A = 57.91 \cdot 8$$

$$44.56$$

$$10.66$$

Formulas

Sum and Difference Identities

$$(a) \cos(\alpha \pm \beta) = \cos \alpha \cdot \cos \beta \mp \sin \alpha \cdot \sin \beta$$

$$(b) \sin(\alpha \pm \beta) = \sin \alpha \cdot \cos \beta \pm \cos \alpha \cdot \sin \beta$$

$$(c) \tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \cdot \tan \beta}$$

Double-Angle Identities

$$(d) \sin(2\alpha) = 2 \cdot \sin \alpha \cdot \cos \alpha$$

$$(e) \cos(2\alpha) = \cos^2 \alpha - \sin^2 \alpha = 2 \cdot \cos^2 \alpha - 1 = 1 - 2 \cdot \sin^2 \alpha$$

$$(f) \tan(2\alpha) = \frac{2 \cdot \tan \alpha}{1 - \tan^2 \alpha}, \tan \alpha \neq 1$$

Half-Angle Identities

$$(a) \sin\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos \alpha}{2}}$$

$$(b) \cos\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 + \cos \alpha}{2}}$$

$$(c) \tan\left(\frac{\alpha}{2}\right) = \pm \sqrt{\frac{1 - \cos \alpha}{1 + \cos \alpha}} = \frac{\sin \alpha}{1 + \cos \alpha}, \cos \alpha \neq -1$$

$$(d) \tan\left(\frac{\alpha}{2}\right) = \frac{1 - \cos \alpha}{\sin \alpha}, \sin \alpha \neq 0$$

Product/Sum Identities

$$(a) 2 \cdot \cos \alpha \cdot \cos \beta = \cos(\alpha - \beta) + \cos(\alpha + \beta)$$

$$(b) 2 \cdot \sin \alpha \cdot \sin \beta = \cos(\alpha - \beta) - \cos(\alpha + \beta)$$

$$(c) 2 \cdot \sin \alpha \cdot \cos \beta = \sin(\alpha + \beta) + \sin(\alpha - \beta)$$

$$(d) 2 \cdot \cos \alpha \cdot \sin \beta = \sin(\alpha + \beta) - \sin(\alpha - \beta)$$

		$\sin$	$\cos$	$\tan$
$\frac{\pi}{6}$	30	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{3}}{3}$
$\frac{\pi}{4}$	45	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{2}}{2}$	1
$\frac{\pi}{3}$	60	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$

Formulas

- Law of Sines:
$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

- Law of Cosines:
$$\begin{aligned}a^2 &= b^2 + c^2 - 2bc \cdot \cos A \\b^2 &= a^2 + c^2 - 2ac \cdot \cos B \\c^2 &= a^2 + b^2 - 2ab \cdot \cos C\end{aligned}$$

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

- "Heading" : Angle made in a clockwise direction from the north
"Bearing" :

- Area of a Triangle:
$$Area = \frac{1}{2} ab \cdot \sin C$$

$$Area = \frac{1}{2} ac \cdot \sin B$$

$$Area = \frac{1}{2} bc \cdot \sin A$$

$$Area = \frac{1}{2} b \cdot h$$

- Heron's Formula:
$$K = \sqrt{s(s-a)(s-b)(s-c)}, \text{ where } s = \frac{a+b+c}{2}$$

- Area of a Sector:
$$A_{\text{sector}} = \frac{C.A.}{360} \cdot \pi r^2 \text{ or } A_{\text{sector}} = \frac{1}{2} \cdot r^2 \cdot \theta$$

Final

Part I EXAM CONCEPTS

Chapters 1, 2, 4, 5, 6

1) Modeling - curve of best fit

2) Functions

Domain, Range, Inverse, Composite Functions

Piecewise Functions

Graphing, translations, reflections (shifts)

3) Symmetry

4) Zeros of a function

5) Distance between 2 pts, pt and a line

6) Rational Root Theorem, Factor Theorem

~~Remainder Theorem~~

10) Complex numbers

7) Asymptotes

9) Trig (9 questions)

8) Radical Equations

Exam - Part II (Chapters ^{3, 7, 8, 9} ~~1, 2, 3, 4, 5, 6~~)

- 1) Exponents
- 2) Logs
- 3) ~~Polar Equations~~
- 4) Conic sections
- 5) Matrices | Systems | Partial Fraction
Decomposition
- 6) Sequences and Limits, Binomial
Series Theorem
- 7) ~~Vectors~~ & Parametric Equations

Directions: Welcome to PreCalc POWER. PreCalc POWER is a worksheet that you will help prepare you for the final exam in this course. There will always be 10 questions, and you must try to complete all questions **correctly, with work (when necessary)**. You may use your notes and discuss with others but you may copy from others!! Good Luck!!

1. Find the inverse of the function $f(x) = \frac{1}{\sqrt{x+2}}$. Restrict the domain of the original function, if necessary.

$$x = \frac{1}{\sqrt{y+2}}$$

$$x\sqrt{y+2} = 1$$

$$\sqrt{y+2} = \frac{1}{x}$$

$$y+2 = \left(\frac{1}{x}\right)^2$$

$$y = \left(\frac{1}{x}\right)^2 - 2$$

Answer: $y = \left(\frac{1}{x}\right)^2 - 2$

$$x, x \in \mathbb{R}$$

3

2. Find the domain and range of the function $f(x) = \frac{3}{x^2 - x - 20}$.

Domain: $x \in \mathbb{R} \neq 5, -4$

Range: $y \in \mathbb{R} \neq 0$

3. Find a real quadratic equation in standard form with real coefficients that has $1+2i$ as a solution.

$$\begin{matrix} 1+2i & 1-2i \\ [x - (1+2i)] & [x - (1-2i)] \\ (x-1)+2i & (x-1)-2i \end{matrix}$$

$$(x^2 - 2x + 1) + (2i \cdot -2i) = x^2 - 2x + 1 + 4 = x^2 - 2x + 5$$

Answer: $x^2 - 2x + 5$

4. Find the equation of a line parallel to the line $3x - 2y + 6 = 0$ and passing through the point $(4, 3)$.

$$-2y = -3x + 6$$

$$y = \frac{3x+6}{2}$$

$$y = \frac{3x+6}{2}$$

Answer: $y = \frac{3}{2}x - 3$

$$0 = 3x - 2y - 6$$

5. Solve the quadratic equation $8x^2 + 22x - 21 = 0$ by factoring.

$$\begin{array}{r} a \cdot c \\ 28 \times -168 \\ -6 \\ 22 \\ b \end{array}$$

$$\begin{array}{c} 2x + 7 \\ 4x \times \frac{1}{3} \\ \begin{array}{|c|c|} \hline 8x^2 & 28x \\ \hline -6x & -21 \\ \hline \end{array} \end{array}$$

$$(2x+7)(4x-3)$$

$$2x = -7 \quad 4x - 3 = 0$$

$$x = -\frac{7}{2} \quad 4x = 3$$

$$x = -\frac{7}{2} \quad x = \frac{3}{4}$$

Answer: $-\frac{7}{2}, \frac{3}{4}$

6. Bo Finkle is at the Phillies game, and he is sitting in the first row of the outfield bleachers (about 26 feet off of the ground). His favorite hitter, Ace Einhorn, has just launched a monster homerun from home plate, and it is heading directly to Bo. If Ace hits the ball at a speed of 105 feet per second and at a height of 3 feet off of the ground, how long will Bo have to react in order to catch the homerun? Use $h(t) = -16t^2 + vt + s$, where $h(t)$ is the height of the ball in feet after t seconds, v is the initial velocity of the ball in feet per second, and s is the starting height for the ball in feet.


$$26 = -16t^2 + 105t + 3$$

find intersection

← solve on calc

Answer: 6.33 sec

7. The terminal side of an angle θ in standard position passes through the point $(-7, 24)$. Determine the **exact** values for $\sin\theta$ and $\cos\theta$.



$$r^2 = 7^2 + 24^2$$

$$r = 25$$

$$\sin\theta = \frac{24}{25}$$

$$\sin = \frac{y}{r} \quad \frac{24}{25} \quad \cos \frac{x}{r} = \frac{-7}{25}$$

$$\cos\theta = \frac{-7}{25}$$

8. For $0 \leq \theta < 2\pi$, determine the values for θ when $\sin\theta = -0.4312$.

↓ inverse sin

$$-25.54 = \frac{x}{r}$$

$$+360 = 334.456$$

⊕ ⊕
⊖ ⊖
3rd

$$180 + 25.54 \rightarrow$$

$$205.34^\circ$$

$$5.84r$$

Answer: 334.456

$$3.59r$$

9. Determine the zeros of the polynomial function $f(x) = 2x^3 - 17x^2 + 11x + 15$. Show your work! Put in calc to find start

$$\begin{array}{r|rrrr} +1.5 & 2 & -17 & 11 & 15 \\ & \downarrow & & & \\ & 2 & -14 & -10 & 0 \end{array}$$

$$2x^2 - 14x - 10$$

$$\text{roots: } \approx 7.65$$

$$\approx -0.65$$

write out + keep roots

Answer: -1.5, 1.5, 7.65

write as roots

10. **Double Bonus:** The dimensions of a rectangular box are given by three consecutive integers, and its volume is 1716 cm^3 . Determine the dimensions of the box.

$$\begin{array}{ccccccc} 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ \downarrow & & & & & & & \\ 858 & & & & & & & \end{array}$$

$$429 \cdot 2 \cdot 2 = 1716$$

$$11/12/13$$

1

PreCalculus

PreCalc POWER II

6. A company wishes to make boxes with volume 48 in.^3 from sheets of cardboard 8 in. by 10 in. by cutting squares from each corner. What size squares should they cut? What are the dimensions of the boxes?

$6 \cdot 4 \cdot 2$

Answer: $6 \cdot 4 \cdot 2 \text{ in}$

7. Determine the zeros and the number of relative extrema of following polynomial function: $h(x) = x^4 + x^3 - 12x^2 - 28x - 16$. $\leftarrow 4^\circ$, so 3 extrema

$$\begin{array}{r} -1 \overline{) 1 \ 1 \ -12 \ -28 \ -16} \\ \underline{1 \ 1 \ -12 \ -28} \\ 0 \ 0 \ 0 \ 0 \end{array}$$

$$\begin{array}{r} 4 \overline{) 1 \ 4 \ 16 \ 16} \\ \underline{1 \ 4 \ 16 \ 16} \\ 0 \ 0 \ 0 \ 0 \end{array}$$

Answer: $(x+1)(x-4)(x+2)^2$
 $\min(-2, 0)(2, 59, \infty)$
 $\max(1, 34, 79)$

8. Evaluate $(g \circ f)(2)$ given $f(x) = x^2 + 1$ and $g(x) = 3x - 4$.

$3(x^2 + 1) - 4$
 11

Answer: 11

9. Divide using synthetic division: $(x^4 + 2x^2 - 10) \div (x + 1)$. Give your answer as a quotient only.

$$\begin{array}{r} -1 \overline{) 1 \ 0 \ 2 \ 0 \ -10} \\ \underline{1 \ 1 \ 3 \ -3} \\ 0 \ 0 \ 0 \ 0 \end{array}$$

Answer: $x^3 - x^2 + 3x - 3 - \frac{7}{x+1}$

10. Double Bonus: If $p(x) = 3x^3 + 2x^2 + Kx + M$, $p(-3) = -41$, and $p(2) = 29$, find K and M .

$-41 = 3(-3)^3 + 2(-3)^2 + K(-3) + M$
 $-81 + 18 - 3K + M$
 $22 = -3K + M$

$29 = 3(2)^3 + 2(2)^2 + K(2) + M$
 $24 + 8 + 2K + M$
 $32 = 2K + M$
 $K = -5$
 $M = 7$

Calc: $\begin{bmatrix} -3 & 1 \\ 2 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 22 \\ -3 \end{bmatrix} = \begin{bmatrix} -5 \\ 7 \end{bmatrix}$

Directions: Directions: Welcome to PreCalc POWER. PreCalc POWER is a worksheet that you will help prepare you for the final exam in this course. There will always be 10 questions, and you must try to complete all questions **correctly, with work (when necessary)**. You may use your notes and discuss with others but you may copy from others!! Good Luck!!

1. Express the following in expanded form: $\log_b \sqrt[3]{4x^5y^7}$

$$\frac{1}{3} \log_b (4x^5y^7)$$

$$\frac{1}{3} \log_b 4 + \frac{5}{3} \log_b x + \frac{7}{3} \log_b y$$

Answer: $\frac{1}{3} \log_b 4 + \frac{5}{3} \log_b x + \frac{7}{3} \log_b y$

2. Find all zeros (including complex zeros) of $f(x) = x^4 + x^2 - 22x - 24$. Show all work algebraically! Seed on calc

$$\begin{array}{r|rrrr} 1 & 1 & 0 & 1 & -22 & -24 \\ & & 1 & 1 & 30 & +24 \\ \hline & & 1 & 1 & 30 & +24 \\ & & & 1 & 3 & 10 \\ & & & & +8 & 0 \\ \hline & & & & & 0 \end{array}$$

$$-1) \downarrow -1 \quad -2 \quad -8$$

$$1 \quad 2 \quad 8 \quad 0$$

$$x^2 + 2x + 8$$

$$\frac{-2 \pm \sqrt{-28}}{2}$$

$$\frac{-2 \pm 2\sqrt{7}i}{2}$$

$$(-1 \pm \sqrt{7}i)$$

Answer: $(0, 3)(0, -1)(0, -1 + \sqrt{7}i)(0, -1 - \sqrt{7}i)$

3. Determine the asymptotes for each of the following functions:

(a) $f(x) = \frac{2x-1}{4x+3}$

horiz asy = $\frac{2}{4} = \frac{1}{2}$

vert asy = $4x+3=0$

$4x = -3$

$x = -\frac{3}{4}$

Answer:

$y = \frac{1}{2}, x = -\frac{3}{4}$

no horiz asy for logs

(b) $g(x) = \log_2(2x-3)$

$\log 2^y = 2x-3$

Answer:

$x = \frac{3}{2}$

find domain (where $\log(0)$)

4. Determine the exact value for each of the following:

(a) $\cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$

30°

$\oplus 1st \ 30^\circ \ \frac{\pi}{6}$
 $\oplus 4th \ 330^\circ \ \frac{11\pi}{6}$

Answer:

$\frac{\pi}{6}, \frac{11\pi}{6}$ b/c domain + range $(0, \pi)$
domain $(-1, 1)$

(b) $\arccos\left(\cos \frac{5\pi}{3}\right)$

$\cos^{-1}(1.5)$ $\oplus 1st \ 60^\circ \ \frac{\pi}{3}$
 $\oplus 4th \ 300^\circ \ \frac{5\pi}{3}$

Answer:

$\frac{\pi}{3}, \frac{5\pi}{3}$ outside of $(0, \pi)$

5. A communications system is modeled on the coordinate plane. The top of the transmitting tower is located at $(-6, -25)$, and the receiving dish is located at $(4, 11)$. If the signal travels on a straight path from the tower to the dish, find the distance from an antenna located at $(9, 5)$ to the path of transmission.



$\frac{-25-11}{-6-4} = \frac{-36}{-10} = 3.6$

$-3.6x + y + 3.4 = 0$
y intercept

Answer:

$(6, 4)$
 $-3.6(9) + 5 + 3.4$
 $\sqrt{(-3)^2 + (1)^2}$
 $\sqrt{9+1}$

6. The owner of a small restaurant called, *No Soy Grande*, can prepare a particular gourmet meal at a cost of \$10. He estimates that if the menu price of the meal is x dollars, then the number of customers who will order that meal at that price in an evening is given by the function $N(x) = 40 - x$. Express his nightly revenue, cost, and profit on this meal as functions of x .

multiply it out

Revenue:

$$N(x) \cdot x \rightarrow 40x - x^2$$

Cost:

$$10x$$

Profit:

$$N(x)x - 10x$$

$$50x - x^2 - 400$$

7. Derive a polynomial function of lowest degree with rational coefficients that has the zeros $2 - \sqrt{2}$ and 4.

$$(x-4)(x-(2-\sqrt{2}))(x-(2+\sqrt{2}))$$

$$(x-4)(x^2-2^2-2)$$

$$(x-4)(x^2-4x+4-2)$$

$$x^3-4x^2+2x-4x^2+16x-8$$

Answer $x^3 - 8x^2 + 18x - 8$

8. Solve the trig equation $2\sin^2 x + 3\cos x - 3 = 0$

$$2(1-\cos^2 x) + 3\cos x - 3 = 0 \quad (-\sin x + 1)(2\sin x - 1)$$

$$2 - 2\cos^2 x + 3\cos x - 3 = 0$$

$$-2\cos^2 x + 3\cos x - 1 = 0$$

$$2x^2 - 3x + 1 = 0$$

$$\sin x = 1$$

$$90^\circ, \frac{\pi}{2}$$

$$2\sin x = 1$$

$$\sin x = \frac{1}{2}$$

$$30^\circ$$

Answer

$$\frac{\pi}{6}, \frac{\pi}{2}, \frac{5\pi}{6}$$

no restriction

9. Solve $\cot x \cos^2 x = 2 \cot x$

$$\cot x \cos^2 x - 2 \cot x = 0$$

$$\cot x (\cos^2 x - 2) = 0$$

$$\cot x = 0$$

$$\tan x = \frac{1}{0}$$

$$\text{und.}$$

$$\cos^2 x = 2$$

$$\cos x = \pm \sqrt{2}$$

$$\text{und.}$$

$$\frac{\pi}{2}, \frac{3\pi}{2}$$

Answer

$$\text{undefined}$$

$$\frac{\pi}{2} + n\pi$$

10. Double Bonus: What is the number of factors when $x^{13} - x$ is factored completely into polynomials and monomials with real integral coefficients? To prove your answer, list all of the factors.

7 factors

$$x(x^{12}-1)$$

$$x(x^6-1)(x^6+1)$$

$$x(x^3-1)(x^3+1)(x^6+1)$$

$$x(x-1)(x^2+x+1)(x+1)(x^2-x+1)(x^2+1)(x^4-x^2+1)$$

$$x(x-1)(x^2+x+1)(x+1)(x^2-x+1)$$

$$x(x-1)(x^2+x+1)(x+1)(x^2-x+1)$$

Sum of cubes $\rightarrow$ factoring pattern x^3+y^3

Difference of cubes $\rightarrow x^3-y^3 = (x-y)(x^2+xy+y^2)$

$$x^3-y^3 = (x-y)(x^2+xy+y^2)$$

$$x^3-y^3 = (x-y)(x^2+xy+y^2)$$

$$x^3-y^3 = (x-y)(x^2+xy+y^2)$$

$$x^3-y^3 = (x-y)(x^2+xy+y^2)$$

$$x^3-y^3 = (x-y)(x^2+xy+y^2)$$

Directions: Directions: Directions: Welcome to PreCalc POWER. PreCalc POWER is a worksheet that you will help prepare you for the final exam in this course. There will always be 10 questions, and you must try to complete all questions **correctly, with work (when necessary)**. You may use your notes and discuss with others but you may copy from others!! Good Luck!!

- 4.5 (1) State the domain and range of the function $f(x) = 3 \cdot \cos(2x) + 2$.

Domain: $[-\pi, \pi]$

Range: $[-1, 1]$

- (2) Decompose the following into partial fractions: $\frac{6x^2 + 25x - 24}{x^3 + x^2 - 12x}$.

(1) $x(x^2 + x - 12)$

Answer: _____

- (3) Find the equation of the slant asymptote of the graph of this rational function

$$\frac{2x^3 + 5x^2 - 23x - 14}{x^2 - 2x - 3}$$

$$\begin{array}{r} 2x - 9 \\ x^2 - 2x - 3 \overline{) 2x^3 + 5x^2 - 23x - 14} \\ \underline{-(2x^3 - 4x^2 - 6x)} \\ 9x^2 - 17x - 14 \end{array}$$

Answer: $2x - 9$

4. Prove the following trig identity: $\cos^2 \theta - \sin^2 \theta = \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}$. Show all work.

(1) $(1 - \sin^2 x) - \sin^2 x$
 $1 - 2\sin^2 x$

$$\frac{\cos^2 x}{\cos^2 x} - \frac{2}{\cos^2 x}$$

Answer: _____

5. Find the real zeros, algebraically, of the following polynomial function

$$f(x) = x^4 + x^3 - x - 1$$

$$\frac{-1 \pm \sqrt{-3}}{2}$$

$$\frac{-1 \pm \sqrt{3}i}{2}$$

Answer: $x = 1, x = -1, x = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i$

6. Divide $\frac{6-i}{8+3i}$ and write the result in standard form.

$$\frac{6-i}{8+3i} \times \frac{8-3i}{8-3i} = \frac{48-8i^2-18i+3i^2}{64-24i+24i-9i^2} = \frac{48-8(-1)-18i+3(-1)}{64-24i+24i-9(-1)} = \frac{48+8-18i-3}{64-24i+24i+9} = \frac{53-18i}{73}$$

Answer: $\frac{53}{73} - \frac{18i}{73}$

7. Expand the following binomial $(3x^2 - 4y^3)^5$. Show all steps.

$$(3x^2)^5 + 5(3x^2)^4(-4y^3) + 10(3x^2)^3(-4y^3)^2 + 10(3x^2)^2(-4y^3)^3 + 5(3x^2)(-4y^3)^4 + (-4y^3)^5$$

$$243x^{10} + 5 \cdot 81x^8 \cdot (-4y^3) + 10 \cdot 27x^6 \cdot 16y^6 + 10 \cdot 9x^4 \cdot (-64y^9) + 5 \cdot 3x^2 \cdot 256y^{12} - 1024y^{15}$$

Answer: $243x^{10} - 1620x^8y^3 + 4320x^6y^6 - 3840x^4y^9 + 1280x^2y^{12} - 1024y^{15}$

8. If $f(x) = 3x^2$ and $g(x) = 7 - x$, determine $(f \circ g)(-x - 4)$.

$$f(g(x)) = 3(7-x)^2 = 3(49 - 14x + x^2) = 147 - 42x + 3x^2$$

$$f(g(-x-4)) = 3(-x-4)^2 = 3(x^2 + 8x + 16) = 3x^2 + 24x + 48$$

Answer: $3x^2 + 24x + 48$

9. Determine the domain of the logarithmic equation $\log_2(x-4) + \log_2(5) = 3$, and then solve for x .

$$\log_2((x-4)(5)) = 3$$

$$\log_2(5x-20) = 3$$

$$2^3 = 5x-20$$

$$8 = 5x-20$$


$$28 = 5x$$

$$x = 5.6$$

Domain: $x > 4$

Answer: $x = 5.6$


10. Double-Bonus: Given $E(3, 4)$ and $F(-3, -4)$, write an equation in terms of x and y for all points $P(x, y)$ such that segment PE is perpendicular to segment PF . Simplify the equation and explain your result.



Directions: Welcome to PreCalc POWER. This POWER is worth **20 points**, with each question being worth **2 points**. Read each question carefully and show all work necessary. Good Luck.

NAME: _____

1. Solve the triangle below for x .



Answer: 44.81°

$$\frac{\sin 110}{20} = \frac{\sin x}{15}$$

$$20 \sin x = 14.10$$

$$44.81^\circ$$

2. Determine the period and phase shift of the function $f(x) = 2 \cdot \tan\left(-\frac{3}{4}x - 5\right)$.

$$y = a^x$$

$$\frac{2\pi}{b} \quad \begin{matrix} \text{sin} \\ \text{cos} \end{matrix}$$

Period: π

Phase Shift: $-\frac{20}{3}$

3. Find the inverse of the function $f(x) = x^2 - 6x + 9$. Restrict the domain of the original function, if necessary, in order to make the inverse a function.

$$x = y^2 - 6y + 9$$

$$x = (y - 3)^2$$

$$\sqrt{x} = y - 3$$

Answer: $\sqrt{x} + 3$

4. Find the general equation of a line that is perpendicular to $4x - 9y - 4 = 0$ and passes through the point $(9, -2)$.

$$y - y_1 = m(x - x_1)$$

$$y - -2 = \frac{9}{4}(x - 9)$$

Answer: 2

5. Solve the exponential equation $x^2 e^{x+1} + 4x e^{x+1} - 5e^{x+1} = 0$.

$$y = \log_a x$$

$$x = a^y$$

$$e^{x+1} (x^2 + 4x - 5)$$

$$x + 1 = e^0$$

Answer: $x = 1$

6. Sketch the graph of $f(x) = -3\csc(\frac{4}{5}x)$. Label your axes and scale completely.

Sketch:

7. Solve the equation $\sqrt{x+30} = x+10$.

$$x+30 = x^2 + 100$$

$$x^2 - x - 70 = 0$$

Answer: _____


8. Determine all remaining zeros of the polynomial

$f(x) = 3x^4 - 16x^3 + 24x^2 + 44x - 39$ given $3 + 2i$ is a zero.

$$\begin{array}{r} 3+2i \overline{) 3 \quad -16 \quad 24 \quad 44 \quad -39} \\ \underline{3} \\ 0 \end{array}$$

Answer: _____

9. A bridge has a parabolic arch that is 20 feet high in the center and 50 feet wide at the bottom. If a two-lane highway passes underneath the bridge, and the end of each lane is 15 feet from the center of the arch, how much clearance would a truck or large vehicle have for passing under the bridge (assuming that the driver of the vehicle obeys the law and stays in his or her proper driving lane)?



Answer: 13.28 feet

$$\frac{x^2}{25^2} + \frac{y^2}{20^2}$$

$$14375 = \frac{x^2}{20^2}$$

$$\sqrt{175}$$

$$13.28$$


10. Double-Bonus: Determine the **two** equations, in conic form (Chapter 9 Style), for a parabola containing the points (1, 3), (-1, 9), and (5, 15).

Equation #1: _____


Equation #2: _____

Directions: Welcome to PreCalc POWER. This POWER is worth **20 points**, with each question being worth **2 points**. Read each question carefully and show all work necessary. Good Luck!!.

1. Determine the area of the segment of the circle with arc degree measure 80° and radius 26 cm.



2. Determine the domain and range of the function $f(x) = \frac{3}{\sqrt{x-2}}$.



Domain: $x > 2$
Range: $y > 0$

3. What is the equation of the function that results when $f(x) = \log_7 x$ is reflected through the line $y = x$?

$$x' = 7^{f(x)}$$

Answer: _____

4. Determine the rational zeros of the function $f(x) = x^3 - 3x^2 - 6x + 8$. Show all of your work.

$$\begin{array}{r|rrrr} 1 & 1 & -3 & -6 & 8 \\ & & 1 & -2 & -8 \\ \hline & 1 & -2 & -8 & 0 \end{array}$$

Handwritten notes: $x^2 - 2x - 8$, $x = 4$, $x = -2$

Answer: $1, 4, -2$

5. Simplify the expression: $\left(\frac{x^{\frac{11}{2}} \cdot y^{\frac{-1}{2}}}{x^{\frac{2}{3}} \cdot y^{\frac{5}{3}}} \right)^5$

$$\frac{x^{\frac{11}{2} \cdot 5} \cdot y^{\frac{-1}{2} \cdot 5}}{x^{\frac{2}{3} \cdot 5} \cdot y^{\frac{5}{3} \cdot 5}} = \frac{x^{\frac{55}{2}} \cdot y^{\frac{-5}{2}}}{x^{\frac{10}{3}} \cdot y^{\frac{25}{3}}}$$

$$x^{\frac{145}{6}} \cdot y^{\frac{-125}{6}}$$

Answer: $x^{\frac{145}{6}} \cdot y^{\frac{-125}{6}}$

6. Determine the distance between the graphs of the equations $2x - 5y = 9$ and $-10x + 25y = 14$.

$$+10x \quad +10x$$

$$25y = 10x + 14$$

$$y = \frac{2}{5}x + \frac{14}{25}$$

$$-\frac{9}{5} = -\frac{14}{25}$$

Answer: 59/25

7. If $\tan \theta = -\frac{12}{5}$ and θ is in Quadrant IV, find the exact value of $\cos \theta$.

$$\sqrt{5^2 + 12^2}$$

$$\cos \theta = \frac{5}{13}$$

Answer: 5/13

8. Determine the exact value of $\sin(\tan^{-1} \sqrt{3})$.

$$\sin(60^\circ)$$

$$\frac{\sqrt{3}}{2}$$

Answer: √3/2

9. Express $\log_b \left(\frac{\sqrt[3]{mt}}{w^k} \right)$ in expanded form.

$$\log_a a^x = x \quad \log_b (\sqrt[3]{mt}) = \log_b (w^k)$$

$$a^{\log_a x} = x \quad \frac{1}{3} \log_b (mt) = k \log_b w$$

$$\frac{1}{3} [\log_b (m) + \log_b (t)] - k \log_b w$$

Answer: 1/3 [log_b (m) + log_b (t)] - k log_b w

10. Double-Bonus: In $\triangle ABC$, $\angle A = 120^\circ$, $c + a = 21$, and $b + a = 20$. Find a . (Don't cry; you need to know the law if you want this one).

Answer: _____

Directions: Welcome to PreCalculus POWER. PreCalculus POWER is a worksheet that you will complete once a week to prepare you for the final exam in this course. There will always be 10 questions, and you must complete all questions **correctly, with work (when necessary)**. IA POWER will always be worth **20 points**, with each question being worth **2 points**. Finally, some weeks there will be a final question worth double the points if you get it correct (but not a double loss of points if wrong). Good Luck..

1. Find the inverse of the function $f(x) = \frac{\sqrt{x}}{2}$. Restrict the domain of the original function, if necessary.

$$x = \frac{\sqrt{y}}{2}$$

$$2x = \sqrt{y}$$

$$(2x)^2 = y$$

$$y = 4x^2$$

Answer: $4x^2$

2. The relation f is : $\{(2, -3), (5, -4), (8, -2)\}$. Give the domain and determine what the inverse of this relation is.

Domain: _____

Inverse: _____

3. Find the minor M_{32} of matrix A : $A = \begin{bmatrix} 7 & -1 & -2 \\ -4 & 8 & 1 \\ 9 & 5 & 6 \end{bmatrix}$.

determinant of only certain columns
row 3 and col 2 removed

Answer: -1

4. Find the equation of a line perpendicular to the line $3x - 2y + 6 = 0$ and passing through the point $(4, 3)$.

$$y - 3 = -\frac{2}{3}(x - 4)$$

$$y - 3 = -\frac{2}{3}x + \frac{8}{3}$$

$$y = -\frac{2}{3}x + \frac{17}{3}$$

Answer: $y = -\frac{2}{3}x + \frac{17}{3}$

5. Given $1^3 + 2^3 + 3^3 + 4^3 + \dots$, give the sigma notation for the first 20 terms of the series.

Answer: $\sum_{i=1}^{20} i^3$


SHOW ALL WORK! 20 points

NAME: _____

1. Determine a function represented by this graph:

$$\text{amp} = 4$$

$$\text{pd} = \frac{2\pi}{6}$$



Answer: $y = 5 \sin\left(\frac{1}{2}x\right)$

2. A record store sells CDs at \$9.50 each. For a CD on the top of the chart, the store expects to sell 150 CDs per week. For each 50 cent decrease in the price, the store will sell approximately 20 more CDs per week. The store wants to make a profit of \$1050. If each CD costs the \$3.00, determine the price at which they should sell the CDs.

price - people

$$(9.50 - 150(x)) \cdot (150 + 20(x)) = 1050$$


$$975 + 130x - 75x - 10x^2 = 1050$$

$$-10x^2 + 55x - 75 = 0$$

factor

Answer: \$8.25 or \$8

3. From this graph, determine the open intervals for which $f(x)$ is increasing, decreasing, or constant.



Increasing: $P \in x < -3, P \in 0 < x < 4$

Decreasing: $P \in x > 4$

Constant: $P \in -3 < x < 0$

4. Determine k so that $h(x) = 2x^3 + 5x^2 + \boxed{kx} - 16$ has $x - 2$ as a factor.

2)
$$\begin{array}{r} 2x^3 + 5x^2 + kx - 16 \\ \underline{-(2x^3 - 4x^2 + 4x - 8)} \\ 9x^2 + 18x - 8 \end{array}$$

Answer: _____

5. Solve this equation $4^{2x-1} = 5^{x+2}$.

$$\log_{10}(4^{2x-1}) = \log_{10}(5^{x+2})$$

$$2x - 1(\log 4) = x + 2(\log 5)$$

Answer: _____

6. In a geometric series if $|r| < 1$ the series converges (has a definite limit or sum). If $|r| > 1$ the geometric series diverges or has no sum. Does this series

$6 + 2 + \frac{2}{3} + \frac{2}{9} + \dots$ converge or diverge. If it converges find the sum. Please show all work.

$$\frac{2}{15} + \frac{2}{1} + \frac{2}{3} + \frac{2}{9} + \frac{2}{\infty}$$

$r = \frac{1}{3}$ Converge

$$\frac{a_1}{1-r} = \frac{6}{1-\frac{1}{3}} \rightarrow$$

Answer: 9

7. Find all solutions of $\sin 5x + \sin 3x = 0$ in the interval $[0, 2\pi]$.

Calc
Zoom trig
find 0s

Answer: $0, 45^\circ, 90^\circ, 135^\circ, 180^\circ, 225^\circ, 270^\circ, 315^\circ$

8. Simplify this expression $\frac{(xy)^{-2}(yz)^{-3} \frac{1}{2}}{(xz)^3(zw)^{-2}}$. Show all work!

Answer: _____

9. Determine all asymptotes of this function $f(x) = \frac{2x^2 - 3x}{x+1}$. Show your work!

horiz asy $\rightarrow$ degree(top) < bottom

$$y=0$$

degree $\rightarrow$ leading coefficients

* top > bottom $\rightarrow$ none

$$\text{vert asy} = x+1=0 \quad x=-1$$

$$\text{Slant } \frac{2x^2-3x}{x+1}$$

$$x+1 \overline{) 2x^2-3x}$$

$$-(2x^2+2x)$$

Answer: $x=-1, y=2x-5$

$$-5x+0$$

$$-(-5x-5)$$

5 $\in$ remainder

10. Solve this equation $665,400 = 560,774e^{7r}$ for r . Show all work.

Answer: _____

Final Review

Extra Problem

1/18

$$\begin{aligned} 1. \quad & x - 2y + 3z = 9 \\ & -x + 3y - z = -6 \\ & 2x - 5y + 5z = 17 \end{aligned}$$

Calc

$$\begin{bmatrix} 1 & -2 & 3 \\ -1 & 3 & -1 \\ 2 & -5 & 5 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 9 \\ -6 \\ 17 \end{bmatrix}$$

$$[A]^{-1}[B]$$

Final Review
Extra Problems

$$x - 2y + 3z = 4$$

$$x + y - z = 1$$

$$2x + 3y - 4z = 7$$

Col

$$\begin{bmatrix} 1 & -2 & 3 \\ 1 & 1 & -1 \\ 2 & 3 & -4 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 \\ 0 & 3 & -4 \\ 0 & 5 & -10 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 \\ 0 & 3 & -4 \\ 0 & 0 & -2 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -2 & 3 \\ 0 & 3 & -4 \\ 0 & 0 & -2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 \\ 0 & 1 & -2 \\ 0 & 0 & -2 \end{bmatrix}$$

