


Lecture 20

Vector Fields + Line Integrals 2D

3/30

Geometrically (in 2D)



← at x, y there is a vector

$$\vec{F}(x, y)$$

Analytic Representation

$$\vec{F}(x, y) = M(x, y) \hat{i} + N(x, y) \hat{j}$$
$$= \langle M, N \rangle$$


↑ functions of 2 variables

"Vector (valued) function of 2 variables"

- not just one vector, system of vectors
- = a vector field
- in a plane
- 2D

Gravitational field (force field)

- 2D



← unit mass at x, y

further away it is, the smaller it is

$$\text{Magnitude } (x, y) = \frac{GM}{r^2} \quad \leftarrow r \text{ is distance to origin}$$

$$\text{Direction } -\frac{\langle x, y \rangle}{r}$$

Magnitude $\cdot \frac{1}{r}$


Field $\vec{F} = -\frac{GM}{r^2} \cdot \frac{\langle x, y \rangle}{r}$

$$= -\frac{GM \cdot (x\hat{i} + y\hat{j})}{(x^2 + y^2)^{3/2}}$$

$$= -\frac{GM \langle x, y \rangle}{r^3}$$

Velocity Field

- of rotating disk
- counter clockwise
- ω = angular velocity



direction $\frac{\langle -y, x \rangle}{r}$

magnitude $\omega \cdot r$


$$\vec{F} = \omega \langle -y, x \rangle$$

$$\omega(-y\hat{i} + x\hat{j})$$

Line Integrals

Calculate work done by a force field $\vec{F}$ along a curve


- Simplest constant force over straight line
= Force $\cdot$ distance
- constant force $\rightarrow$ does not depend on x and y



Work by $\vec{F}$
 P_1 to P_2 (unit mass)

$$\begin{aligned} &= \vec{F} \cdot \hat{U} |\vec{P_1 P_2}| \leftarrow \hat{U} \text{ component of } \vec{F} \cdot \text{distance} \\ &= \vec{F} \cdot \vec{P_1 P_2} \end{aligned}$$

- $\vec{F}$ non constant
 C is real curve
now more complex
* view curve as tiny little line segment



parametrize curve
 $\vec{r}(t) = \langle x(t), y(t) \rangle$

$$\vec{F} = \langle M(x, y), N(x, y) \rangle$$

Work by $\vec{F}$ on Δr_i

↓
Choose a point on line



$$\approx \langle M(x_i, y_i), N(x_i, y_i) \rangle \cdot \Delta \vec{r}_i$$

the arrow
vector

$$\text{total work} \approx \sum_{i=1}^n \langle M(x_i, y_i), N(x_i, y_i) \rangle \cdot \Delta \vec{r}_i$$

$$\Delta \vec{r}_i = \Delta x \vec{i} + \Delta y \vec{j}$$

Total work along curve

$$\approx \sum_{i=1}^n M(x_i, y_i) \Delta x_i + N(x_i, y_i) \Delta y_i$$

Let n go to ∞

$$= \int_C \vec{F} \cdot d\vec{r}$$

Express everything in terms of t

$$\begin{aligned} x &= x(t) & x_i &= x(t_i) \\ y &= y(t) & y_i &= y(t_i) \end{aligned}$$

$$= \int_{t_1}^{t_2} \left[M(x(t), y(t)) \frac{dx}{dt} + N(x(t), y(t)) \frac{dy}{dt} \right] dt$$

$$\int_C \vec{F} \cdot d\vec{r}$$

$$d\vec{r} = dx \hat{i} + dy \hat{j}$$

$$\int_C M dx + N dy$$

← what most people write
for line integral

use t to do parametrisation

$$\int_{t_1}^{t_2} \left[\vec{F}(x(t), y(t)) \cdot \frac{d\vec{r}}{dt} \right] dt$$

use t w/ this equation

$$\int_{t_1}^{t_2} \left[M(x(t), y(t)) \frac{dx}{dt} + N(x(t), y(t)) \frac{dy}{dt} \right] dt$$

Get one massive function of t ,
integrate w/ respect to t

3rd Way

Intrinsic Calculation

insight + speed (if it works)
arc length on curve


Use $t = s$

$$\int_C \vec{F} \cdot d\vec{r} = \int_C \left(\vec{F} \cdot \frac{d\vec{r}}{ds} \right) ds$$


$\hat{T}$

the unit tangent vector

$$= \int_C \vec{F} \cdot \vec{T} ds$$




$\oint$ same



$\vec{F} = -y\vec{i} + x\vec{j}$
 $\omega = 1$
 rotating velocity field
 counter clockwise

$$\int_C \vec{F} \cdot d\vec{r} \rightarrow \underline{C_1} \int_C \vec{F} \cdot \vec{T} ds = 0$$



always
perpendicular
along line

$$\underline{C_2} \int_{C_2} \vec{F} \cdot \vec{T} ds = \int_{C_2} \left[M \frac{dx}{dt} + N \frac{dy}{dt} \right] dt$$

$$= \int_{C_2} -y dx + x dy$$

use x as parameter

$$= \int_0^1 -x^2 dx + x \cdot 2x dx$$

$$= \int_0^1 x^2 dx$$

$$= \frac{1}{3}$$

C₃ Most complex of all

$$\int_{C_3} -y dx + x dy$$

$$= \int_{C_3'} -y dx + x dy + \int_{C_3''} -y dx + x dx$$

$$= \int_{C_3'} 0 + x \cdot 0 + \int_{C_3''} 0 + 1$$

since $x=1$
along vertical
line

$$= \int_0^1 dy$$

$$= 1$$

G₀

$$C_1 = 0$$

$$C_2 = \frac{1}{3}$$

$$C_3 = 1$$

Lecture 21

Gradient Fields + Conservative Fields
Potential Functions, FTC for line integrals

4/2

$$\vec{F} = \langle M, N \rangle$$

Test 2


Average

14

Mei 72

$$C = \vec{r}(t) \\ = \langle x(t), y(t) \rangle$$

t_0



Fields

Differentials

Symbol

$$\int_C \vec{F} \cdot d\vec{r} = \int_C M dx + N dy$$

Calc

$$\int_{t_0}^{t_1} \left(\vec{F} \cdot \frac{d\vec{r}}{dt} \right) dt = \int_{t_0}^{t_1} \left(M \frac{dx}{dt} + N \frac{dy}{dt} \right) dt$$

↑
Physics + electrical
engineering

↑ everyone else

↑ I will use this for calculations

↑ this quick answer
see what's going on
* intuition *

↑ calculations *

Learn all 4 forms

5th form

$$\int_C (\vec{F} \cdot \vec{T}) ds$$

? cleaner parametrization
Symbolic calculation

↳ when $t = s$ arclength

$$\text{since } \vec{T} = \frac{d\vec{r}}{ds}$$

1. Remember

Math work done by the field
Physics against

2. Direction you are transversing



If you are travelling the other way




C prime

$$\int_C \vec{F} \cdot d\vec{r} = - \int_{C'} \vec{F} \cdot d\vec{r}$$

3.

$$\int_{t_1}^{t_0} \text{---} dt = - \int_{t_0}^{t_1} \text{---} dt$$

$$\vec{F} = \langle x^2, 2xy \rangle$$



parametrization

$$x = t^2$$

$$y = t^3$$

(only one can calc arc length for)

* ans to
Scalar product
is a scalar

$$\int_C y^2 dx + 2xy dy \quad (\text{differential form})$$

↓ replace

$$\int_0^1 t^6 \cdot 2t dt + 2t^2 \cdot 3t^2 dt$$

$$dx = 2t dt$$

$$dy = 3t^2 dt$$

Now normal $(8,0)$ integral

$$\int_0^1 8t^7 dt$$

$$t^8 \Big|_0^1$$

$$(1)$$


Fundamental theory of line integrals

Gradient field - field $\vec{F}$ which has $\vec{\nabla} f(x, y)$

$$\vec{F} = \left\langle \underset{\substack{\uparrow \\ M}}{\frac{\partial f}{\partial x}}, \underset{\substack{\uparrow \\ N}}{\frac{\partial f}{\partial y}} \right\rangle$$

Fundamental
theory of
calculus
for line
integrals

If f is a gradient field $\vec{F} = \vec{\nabla} f$



$$\begin{aligned} \int_C \vec{F} \cdot d\vec{r} &= f(x, y) \Big|_{P_0}^{P_1} \\ &= f(x_1, y_1) - f(x_0, y_0) \end{aligned}$$

$$\langle y^2, 2xy \rangle = \vec{\nabla} (xy^2)$$

$$\begin{aligned} \int_C y^2 dx + 2xy dy \\ \downarrow \\ = xy^2 \Big|_{(0,0)}^{(1,1)} \\ = 1 \end{aligned}$$

A small diagram of a coordinate system with a curve starting at the origin $(0,0)$ and ending at $(1,1)$.

That's the simple idea
But some more stuff to say about it
Basis for a lot of stuff

Mattuck i
not what you
expect

Proof (to memorize)

Hypothesis $\vec{F} = \nabla f$

$$\int_C \vec{F} \cdot d\vec{r} =$$
$$= \int_{t_0}^{t_1} \left(\frac{\partial f}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial f}{\partial y} \cdot \frac{dy}{dt} \right) dt \quad \text{calculating form}$$

Now 18.01 integral

$$= \int_{t_0}^{t_1} \left[\frac{df(x(t), y(t))}{dt} \right] dt$$

$$= f(x(t), y(t)) \Big|_{t_0}^{t_1}$$

fundamental theorem of
calculus (first)

$$= f(x_1, y_1) - f(x_0, y_0)$$


$$\int_0^1 \text{sect } dt$$

problem is finding anti deriv
Should know

Problems for Tve

1. - given $\vec{F} = \langle M, N \rangle$
how do you tell its a gradient field?
- extremely rare, but important
2. If it is $\vec{F} = \nabla f$ how do you find f ?
3. Translate everything into language of differentials

hypothesis $\vec{F} = \nabla f \Rightarrow$ [if true/implies] #1 and #2



#1,

start + end
at same
pt = same
value of
line integral

$$\int_{C_1} \vec{F} \cdot d\vec{r}$$

$$= \int_{C_2} \vec{F} \cdot d\vec{r}$$


"path independent"

#2.

$$\oint_C \vec{F} \cdot d\vec{r}$$

that is closed

P₀ sign means
path is
closed




$$P_0 = P_1$$

$$= 0$$

"F is conservative"

$$\int_{P_1}^{P_1} = \int_{P_0}^{P_0} = 0$$

relation #1 $\Rightarrow$ #2 = we did it
#2 $\Rightarrow$ #1



$$\text{hyp} \Rightarrow \oint_C \vec{F} \cdot d\vec{r} = 0$$

$$\int_{C_1} + \int_{C_2'} = 0$$

by hypothesis since $C_1 + C_2$
is closed curve

$$\int_{C_1} - \int_{C_2} = 0$$

thus
$$\int_{C_1} = \int_{C_2}$$

Recitation

Test Correction + Line Integrals

4/5

Test 2

#5

On test paper

did not spend much
time on

Lectures

Line integrals

$\int_C \vec{F} \cdot d\vec{r}$

$\vec{F}$ is field $\vec{F}(x,y) = \text{vector}$

$\vec{F}$ vector $d\vec{r}$ vector

dot product

$d\vec{r} = \langle x'(t), y'(t) \rangle$

is "position vector" moving
along C - parameterization

$\int_C P dx + Q dy$

P is function $P(x,y) = \text{number}$

P number Q number

$dy = y'(t) dt$

ex1

Let C be curve parametrized by

$\vec{r}(t) = \langle t+1, t^2-1 \rangle$ $t \in [0,1]$

a) Compute $I = \int_C \vec{F} \cdot d\vec{r}$ where $\vec{F} = \langle x+y, x \rangle$

b) Compute $J = \int_C P dx + Q dy$ where $P(x,y) = y$
 $Q(x,y) = 1$

↑ 2 different equations want to compute, not 2 different
witnesses

So curve has been given to us in terms of t already
know

$$x = t+1$$

$$y = t^2 - 1$$

a Compute I

$$\vec{F} = \langle x+y, x \rangle$$
$$\langle t+1+t^2-1, t+1 \rangle$$
$$\langle t^2+t, t+1 \rangle$$

Now $\int_0^1 \langle t^2+t, t+1 \rangle \cdot \langle t+1, t^2-1 \rangle$

~~so just dot product now~~

$$(t^2+t) \cdot (t+1) + (t+1) \cdot (t^2-1)$$

derivative of original x and y

Need

$$x'(t) = 1$$


$$y'(t) = 2t$$

$$\vec{r}' = \langle 1, 2t \rangle$$

~~and then $t=1 \rightarrow t=0$~~

Can draw the curve

but don't have to



now find dot product

$$F \cdot dr = \langle t^2+t, t+1 \rangle \cdot \langle 1, 2t \rangle$$
$$(t^2+t) \cdot 1 + (t+1) \cdot 2t$$
$$t^2+t + 2t^2+2t$$
$$3t^2+3t$$

$$\begin{aligned}
 I &= \int_0^1 (3t + 3t^2) dt \\
 &= \left. \frac{3t^2}{2} + t^3 \right|_0^1 \\
 &= \frac{3}{2} + 1 = \left(\frac{5}{2} \right) \checkmark
 \end{aligned}$$

b) Now a problem with differentials

$$p = y = t^2 + 1$$

$$Q = 1$$

$$dx = x'(t) dt = 1 dt$$

$$dy = y'(t) dt = 2t dt$$

$$\int_0^1 (t^2 - 1) dt + 2t dt$$

$$\left. \frac{t^3}{3} + t^2 - t \right|_0^1$$

$$\left(\frac{1}{3} \right)$$


Remark

If $F = \langle M, N \rangle$ then

from definition

$$\int_C \vec{F} \cdot d\vec{r} = M dx + N dy$$

ex 2



$$\vec{F} = \langle y, -1 \rangle$$

a) $I = \int_{C_1} \vec{F} \cdot d\vec{r}$

b) $J = \int_{C_2} \vec{F} \cdot d\vec{r}$

Now need to parametrize the circle myself

a) $x(t)$ look for candidates
 $y(t)$ what makes sense

$$x(t) = t$$

$$y(t) = (t-1)^2$$

Then can check that it satisfies

$$y = (x-1)^2$$

Choose a parameter that identifies a point

Let's say x is trial pt

$$\text{So } t = x$$

So what is $y \rightarrow (x-1)^2$

So plug in $t=x$ for $(t-1)^2$



With this curve x is not a good choice

Could use y as parameter

$$y(t) = t$$

$$x(t) = \dots$$

For our function

$$y(t) = (1-t) \quad \leftarrow \text{to go right direction}$$

$$x(t) = \sqrt{1-t}$$

But don't use - too complicated

$$t \in (0,1)$$

So now calculate

$$\left(\frac{4}{3} \right)$$

b) What parameter here?

$$x^2 + y^2 = r^2 \quad \text{unit circle}$$

$$x^2 + y^2 = 1^2 \quad \rightarrow$$

Lets declare $x(t) = t$

$$y(t) = \frac{1-x^2}{\sqrt{1-t^2}}$$

Could also do angle as parameter θ

$$x(t) = \cos t$$

$$y(t) = \sin t$$

t is CCW

t is not time

But we are going clockwise, so

$$x(t) = \cos\left(\frac{\pi}{2} - t\right)$$

$$y(t) = \sin\left(\frac{\pi}{2} - t\right)$$

$$t \in \left[0, \frac{\pi}{2}\right]$$

get from graph
or convert

$$J = \boxed{1 + \pi/4}$$

Free to take whatever parametrization

- it will work out in the end

- Relation between 2 parametrizations is
relation of changing speed

$$\text{new } x(t) = x(f(t))$$

$$y(t) = y(f(t))$$

?

relation of parametrization

Result does depend on curve

Not path independent

since field is not conserved

Lecture 22

Criterion for conservative fields + potential functions

4/6

Special kind

Fund. The. Calculus

Fields

$$M\vec{i} + N\vec{j}$$

Function of 2 variables

gradient

$$\vec{\nabla}f = f_x\vec{i} + f_y\vec{j}$$

$$\int_{P_0}^{P_1} \vec{\nabla}f \cdot d\vec{r} = f(P_1) - f(P_0)$$

Diff - entials

$$Mdx + Ndy$$

$$df = f_x dx + f_y dy$$

exact/total differential

that make it

path independent

$$\int_{P_0}^{P_1} df = f(P_1) - f(P_0)$$

path independent

Problem 1

Given $\begin{cases} \langle M, N \rangle \\ Mdx + Ndy \end{cases}$

how do we decide if it is special?

$$\begin{cases} \vec{\nabla}f \\ df \end{cases}$$

If it does pass the test, how to find the function?

Criterion $M_y = N_x$ ← partial deriv

$$\begin{cases} \langle M, N \rangle = \vec{\nabla}f \\ Mdx + Ndy = df \end{cases}$$

But not true always

- it must be true in the whole x, y plane
- then it is always true

Ex $axy dx + (bx^2 + cy^2) dy$
 $\langle axy, bx^2 + cy^2 \rangle$

exact
 when is it a gradient field
 aka when conservative
 aka when path independent

$$ax \stackrel{?}{=} 2bx$$

$$a = 2b$$

Hypothesis:

$$\begin{cases} M = f_x \\ N = f_y \end{cases} \text{ true in some region of a plane}$$

Calculate

$$\begin{matrix} M_y = f_{xy} \\ N_x = f_{yx} \end{matrix} \quad \left. \begin{matrix} \text{thus} \\ f_{xy} = f_{yx} \end{matrix} \right\}$$

does not really look like it should be true - but it is

- if all 4 2nd order partial derivatives f_{xx} f_{xy} f_{yx} f_{yy} are continuous

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots$$

$$f(x, y) = f(0, 0) + \left[(f_x)_0 + (f_y)_0 \right] + \frac{1}{2!} \left[(f_{xx})_0 x^2 + 2(f_{xy})_0 xy + (f_{yy})_0 y^2 \right] + \dots$$

Its all numbers...

$$= a_0 + (b_1 x + b_2 y) + (c_1 x^2 + c_2 xy + c_3 y^2) + \dots x^m y^n$$

$$\text{For } x^m y^n = f$$

$$f_{xy} = m n x^{m-1} y^{n-1} = f_{yx}$$

Note totally cheating argument - circular

Problem #2 $\langle M, N \rangle$ If it passes the test, how do we find $f(x, y)$

2 ways $\frac{1}{3}$ like first way (of people)
 $\frac{2}{3}$ " second "

Method 1

$$f(P) = f(P_0) + \int_{P_0}^P M dx + N dy$$

Know $\langle M, N \rangle = \nabla f$


constant, not important when differentiating

Gradient field - so over any path

$$\vec{F} = \langle 2xy, x^2 + y^2 \rangle$$

test $2x = 2x \odot \rightarrow$ so gradient field

$$f(x_1, y_1) = \int_{P_0}^{P_1} 2xy dx + (x^2 + y^2) dy$$



C_1 simplest path to calculate

$$= \int_{C_1} 2xy dx + (x^2 + y^2) dy + \int_{C_2} 2xy dx + (x^2 + y^2) dy$$

$y=0 \quad dy=0$
 so it all disappears

$\left[x = x_1 \right] \quad dx=0$
 drops out what is left

$$\int_0^{y_1} (x_1^2 + y^2) dy$$

Parametrization

$$x = x_1$$

$$y = y_1$$

$$= x_1^2 y_1 + \frac{1}{3} y_1^3$$

$$f(x, y) = x^2 y + \frac{1}{3} y^3$$

$$\vec{\nabla} f = \langle 2xy, x^2 + y^2 \rangle \text{ checks } \checkmark$$

Method 2

$$\langle 2xy, x^2 + y^2 \rangle$$

↓

$$f_x = 2xy$$

$$f_y = x^2 + y^2$$

What was original function

$$f = x^2 y + c \quad \leftarrow \text{any functions of } y$$

$$x^2 y + g(y)$$

↳ what is $g'(y)$ ↺

$$f_y = x^2 + y^2$$

$$= x^2 + g'(y)$$

↙ know

$$g'(y) = y^2$$

$$g(y) = \frac{1}{3} y^3$$

$$f(x, y) = x^2 y + \frac{1}{3} y^3$$

do it
systematically
for when 3
variables

Pros of each method

Method 1 - theoretical $\rightarrow$ gives formula for answer
Calculation is easy, straight forward

Method 2 - algorithm for calculation
requires no knowledge of line integrals

Recitation


4/7

Lecture

Def Field $\vec{F}$ is a "gradient" field / "conservative" field
if $\vec{F} = \nabla f$

Prop $\vec{F} = \langle M, N \rangle$ is ∇f if $M_y = N_x$
everywhere

Theorem If $\vec{F} = \nabla f$ then
 $\int_C \vec{F} \cdot d\vec{r} = f(B) - f(A)$
for any curve from $A \rightarrow B$



Same with differentials

Def $Mdx + Ndy$ is "exact" if f such that
 $df = Mdx + Ndy$

Prop $Mdx + Ndy$ exact and only if $M_y = N_x$

Theorem $\int_C Mdx + Ndy = f(B) - f(A)$
if $Mdx + Ndy = df$

ex 1 Let $\vec{F} = \langle e^y, x e^y + 1 \rangle$

a) Is $\vec{F}$ a gradient field?

~~$M_x = 0$~~
 ~~$N_y = 0$~~

* can't switch *

$0 = 0$

yes

I did
wrong

$M_y = e^y$
 $N_x = e^y$

yes

b) Find a function, such that $\vec{F} = \nabla f$ by FTC (method 1)


Let $P_0 = (x_0, y_0)$

$$f(P_0) = f(0) + \int_C \vec{F} \cdot d\vec{r}$$

for any curve C from 0 to P_0

- can pick any curve

- but we always use same curve



$$\int_{C_1} \vec{F} \cdot d\vec{r} + \int_{C_2} \vec{F} \cdot d\vec{r} + f(0)$$

r_{constant}
take as 0

Need to parametrize C_1 and C_2

$C_1 \quad x(t) = t$

$C_1 \quad y(t) = 0$ since constant

$t \in [0, x_0]$

$dx = 1$

$dy = 0$

$C_2 \quad x(t) = x_0$

$C_2 \quad y(t) = t$

$t \in [0, y_0]$

$dx = 0$

$dy = 1$

doing this
slightly
differently
more work

that is always
the same for
that curve

w/ surfaces need 2 parameters

$$\int_{C_1} \langle t, t+1 \rangle \cdot \langle 1, 0 \rangle dt + \int_{C_2} \langle x_0 e^t + 1, 0 \rangle \cdot \langle 0, 1 \rangle dt$$

$$= \int_0^{x_0} 1 dt + \int_0^{y_0} x_0 e^t + 1 dt$$

$$x_0 + x_0 e^{y_0} + y_0 - x_0$$

$$\boxed{x_0 e^y + y} \quad \text{Drop the sub 0}$$

Method 2

We want $\vec{\nabla} f = \vec{F}$
 that is $\begin{cases} f_x = e^y \\ f_y = x e^y + 1 \end{cases}$

Step 1

a) $f = x e^y + c(y)$
 (constant w/ respect to x
 = function of y alone

Step 2

$$f_x = x e^y + c'(y)$$

Step 3

$$c'(y) = 1$$

$$c(y) = y$$

← since $\boxed{f_y = x e^y + c'(y)} = x e^y + 1$

Step 4

$$f = \boxed{x e^y + y} + c$$

? can take this = 0
 so forget about it

Notice it matches

ex2 Let $\vec{F} = \langle \log y + y, \frac{x}{y} + \alpha x + y \rangle$

a) For which value of α is it a gradient field?

$$M_y = \frac{1}{y} + 1$$

$$\frac{1}{y} + 1 = \frac{1}{y} + \alpha$$

$$M_x = \frac{1}{y} + \alpha$$

$$\boxed{\alpha = 1} \quad \checkmark$$

b) For that value find a function f such that $\vec{F} = \vec{\nabla} f$

$$f_x = \ln(y) + y$$

$$f_y = \frac{x}{y} + x + y$$

$$f = \int (\ln(y) + y) dx$$

$$= \cancel{y \ln(y) + y - y} + \cancel{c(y) \cdot x \ln(y) + xy} + c(y)$$

$$= \cancel{y \ln(y) + c(y)}$$

$$f_y = \frac{x}{y} + x + c'(y)$$

$$c'(y) = y \quad \text{copy wrong}$$

$$c(y) = y \quad \frac{y^2}{2}$$


$$f = \cancel{y \ln(y) + y} + c$$

→ should check both differentiate

- think screwed this up somewhere

$$x \ln(y) + xy + \frac{y^2}{2} + c$$

ex 3



$$\vec{F} = \langle x, y \rangle$$

max value of $\int_C \vec{F} \cdot d\vec{r}$ over all possible
curves contained in R

$$F \text{ gradient} \Rightarrow \int_C \vec{F} \cdot d\vec{r} = f(B) - f(A)$$

out of the

(1)

18.02A Problem Set 6B (DUE 4/8/10 10:45AM 2-106)

Hand this in along with Problem Set 6A; staple 6A and 6B together, in that order.

Part I (15 pts. + 10 pts. for Pset 6A)

Lecture 20. Tues. Mar. 30 Vector fields and line integrals in 2D

Read: Notes V1, Textbook: 21.1 (Work: 4A-3bd, 4; 4B-1bef, 2, 3)

Exam 2. Thurs. April 1 11:05-11:55 Walker 3rd floor

The exam will cover Lectures 9 - 19 on Partial Differentiation and Double Integration.
No books, notes, calculators; no use of cell-phones.

Lecture 21. Fri. Apr. 2 Gradient fields, conservative fields, path-independence;
Fundamental theorem of calculus for line integrals.

Read: 21.2 to p. 762 (Work: 4C-1, 2, 3)

Lecture 22. Tues. Apr. 6 Criterion for conservative fields; potential functions.

Read: Notes V2 (Work: 4C-5a, 6ab (do all three both by Method 1 and Method 2))

Lecture 43. Thurs. Apr. 8 Green's theorem (tangential form).

Read: 21.3 to middle p. 768

Part II (20 pts.)

Directions: Try each problem alone for 15 minutes. If you subsequently collaborate, solutions must be written up independently. It is illegal to consult problem sets from previous semesters. With each problem is the day it can be done.

Problem 1. (Tues. 3/30 3 pts.: 1,2)

a) Write down in xy -coordinates the vector field $\mathbf{F}$ whose vector at $P : (x, y)$ runs in the vertical direction from P to the line $L : x + y = 1$.

b) Show *without calculation* that for this field, $\int_C \mathbf{F} \cdot d\mathbf{r} \neq 0$, if C is any positively oriented right triangle with legs parallel to the axes and hypotenuse on L .

Problem 2. (Tues. 3/30 3 pts.)

Imagine the x -axis is an infinitely long heavy wire, of uniform density. The gravitational force it exerts on a unit mass placed at a point $(y, z) \neq (0, 0)$ in the yz -plane is given by

$$\mathbf{F}(y, z) = -c \frac{y\mathbf{j} + z\mathbf{k}}{y^2 + z^2},$$

where c is a positive constant. Taking $c = 1$, find by direct calculation of the line integral (i.e., by using a parametrization of C) the work done *by* the gravitational field in moving a unit mass along each of the following paths C in the yz -plane:

- half-line $z = 1, y \geq 0$; starting point $(0, 1)$;
- circle of radius a with center at the origin, traced counterclockwise;
- line from $(0, 1)$ to $(1, 0)$.

Problem 3. (Fri. 4/2 4 pts.: 1,1,2)

a) Let $\mathbf{F} = -\nabla(\ln r)$, where r is the distance from the origin in the yz -plane. Show that $\mathbf{F}$ is the field of Problem 2, when $c = 1$.

b) Show that for any path C joining two point P_1 and P_2 in the yz -plane, the value of $\int_C \mathbf{F} \cdot d\mathbf{r}$ depends only on the ratio r_2/r_1 , where r_i is the distance of P_i from the origin. What points P_i should be excluded?

c) Let C be a directed line segment of length 1.

(i) If $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$, how is C placed?

(ii) What position of C gives $\int_C \mathbf{F} \cdot d\mathbf{r}$ the largest value?

Problem 4. (Fri. 4/2 3 pts: 1,2) Let $\mathbf{F} = \nabla(x^2y - 3xy^2)$. Let C be the quarter of the ellipse $9x^2 + 4y^2 = 1$ running from the positive x -axis to the positive y -axis.

a) Write the integral $\int_C \mathbf{F} \cdot d\mathbf{r}$ in the $\int_C M dx + N dy$ form

b) Evaluate it in two different but equally easy ways: using the Fundamental Theorem and using path-independence. (Both ways can be done mentally, but write down enough to indicate the reasoning.)

Problem 5. (Tues. 4/6 4 pts.: 1, 1.5, 1.5) Let $\mathbf{F} = xy\mathbf{i} + x^2\mathbf{j}$.

a) Verify that $\mathbf{F}$ is not a conservative field.

b) Try to find a potential function for $\mathbf{F}$ by using Method 1 in the Notes. What goes wrong?

c) Answer the same question for Method 2.

Problem 6. (Tues. 4/6 3 pts.: 1.5, 1.5)

The most general two-dimensional field $\mathbf{F}$ whose direction is radially outward, and whose magnitude depends only on r ($r = \sqrt{x^2 + y^2}$), is

$$\mathbf{F} = f(r)(x\mathbf{i} + y\mathbf{j}).$$

a) Assuming $f(r)$ is differentiable for all r , show that any such $\mathbf{F}$ is conservative.

(Use the chain rule and the standard formulas $\partial r/\partial x = x/r$, $\partial r/\partial y = y/r$.)


b) Find a potential function of the form $g(r)$, if $f(r) = r^n$, where n is a positive or negative integer.

(Solve $\nabla g(r) = \mathbf{F}$ for the function $g(r)$. One value of n has to be handled differently from all the others.)

Part 1

Lecture 20 Vector Fields + line integrals in 2D

4A-3b Write down an explicit equation for

The vector at (x, y) directed radially in towards origin w/ magnitude r^2 

So, what?

Field = magnitude $\cdot$ direction
 $r^2 \langle \text{what is radially inward} \rangle$
 - in polar coords

$$r^2 \langle -r, 0 \rangle$$


$$-r \langle x\hat{i} + y\hat{j} \rangle$$

~~so~~ sh $\frac{-\langle x, y \rangle}{r}$ is radially inward ~~*~~

$$\text{so } r^2 \frac{\langle x\hat{i} + y\hat{j} \rangle}{r} \quad \checkmark$$

- d. Each vector is parallel to $\hat{i} + \hat{j}$ but magnitude varies.

$$\hat{i} + \hat{j}$$



field_(x,y) = magnitude $\times$ distance


$$m \propto \langle \hat{i} + \hat{j} \rangle$$

Is that all?

$$\text{well } f(x,y) \propto \langle \hat{i} + \hat{j} \rangle$$

so m is a function of x and y

4. The electromagnetic force field of a long straight wire along the z -axis carrying a uniform current is a 2D field, tangent to horizontal circles in dir of right hand rule
Magnitude $\frac{k}{r}$



$$\text{so } \frac{k}{r} \propto \langle \hat{i} + \hat{j} \rangle$$

$$\frac{k \langle \hat{i} + \hat{j} \rangle}{r^2}$$

$$k \frac{\langle -y\hat{i} + x\hat{j} \rangle}{r^2}$$

why that $-y\hat{i} + x\hat{j}$

4B-1b

For each field $\vec{F}$ and curve C
 evaluate $\int_C \vec{F} \cdot d\vec{r}$
 Line Integrals in Plane use differentials $\int_C M dx + N dy$

$$\vec{F} = x y \hat{i} - x^2 \hat{j} \quad C = \text{[diagram of a quarter circle in the first quadrant from (0,1) to (1,0)]}$$

$$\vec{F} = \langle M(x,y), N(x,y) \rangle$$

$$= \int_{t_1}^{t_2} \left[M(x(t), y(t)) \frac{dx}{dt} + N(x(t), y(t)) \frac{dy}{dt} \right] dt$$

But how do you actually solve, did not do
 example in class. will wait post recitation
 Never really got that "parametrising" thing

Ok 4/6 - but what function did they give you?

have $\vec{F}$ but no actual function

well that is given by the line as it moves

$$x^2 + y^2 = 1$$

$$x = \sqrt{1-y^2}$$

$$y = \sqrt{1-x^2}$$

So just say

$$x = t$$

$$y = \sqrt{1-t^2}$$

or is it (from notes)


$$x = t$$

$$y = (t-1)^2$$

well same thing right?

No

so its my original
 thought



Now rewrite $\vec{F}$ in terms of t

$$\langle t \sqrt{1-t^2}, -t^2 \rangle$$

now $\int_0^1 \langle t \sqrt{1-t^2}, -t^2 \rangle \cdot \langle x'(t), y'(t) \rangle dt = \int_0^1 \langle t \sqrt{1-t^2}, -t^2 \rangle \cdot \left\langle 1, \frac{-t}{\sqrt{1-t^2}} \right\rangle dt$

$$\left(t \sqrt{1-t^2} \right) + \left(-t^2 \cdot \frac{-t}{\sqrt{1-t^2}} \right) \Big|_0^1$$

$$t \sqrt{1-t^2} + \frac{t^3}{\sqrt{1-t^2}} \Big|_0^1$$

$$1 \sqrt{1-1^2} + \frac{1^3}{\sqrt{1-1^2}}$$

0 + divide by 0 error

They have totally different parameter

these are these problems where you have to guess a parameter

$$x = \cos t \rightarrow dx = -\sin t dt$$

$$y = \sin t \rightarrow dy = \cos t dt$$

And they use differentials

$$\int_C xy dx - x^2 dy$$

$$\int_{\pi/2}^0 -\sin^2 t \cos t - \sin t - \cos^2 t \cdot \cos t$$

$$\int_{\pi/2}^0 -\sin^2 t \cos t - \cos^3 t$$

Now just integrate,
well simplify first

$$= \int_{\pi/2}^0 \cos t \, dt$$

I don't get how they are
simplifying - worked
out on calculator

Now integrate

$$= \sin t \Big|_{\pi/2}^0$$

$$= \sin 0 - \left[-\sin \pi/2 \right]$$

$$= 0 + 1$$

①



But my question is how do you pick the
parameterization. It does always = the same
thing in the end, but the amount of work
is different. I prob screwed mine up somewhere

e $F = 6y \, x + x \, y$ $x = t^2$ $y = t^3$ from $(1,1)$ to $(4,8)$

ok first write that

$$x = t^2 \rightarrow dx = 2t \, dt$$

$$y = t^3 \rightarrow dy = 3t^2 \, dt$$

was already parametrized for C

$$\int 6y \, dx + x \, dy$$

These are x right?

$$\int_1^2 6 \cdot t^3 \cdot 2t + t^2 \cdot 3t^2$$

$$12t^4 + 3t^4$$

$$15t^4$$

$$\frac{15t^5}{5} \Big|_1^2$$

$$3t^5$$

$$3(2)^5 - 3(1)^5$$

$$96 \quad 3072 - 3$$

$$\cancel{3069}$$

$$93$$

✓


So I picked the wrong integral bounds - what I was not sure of - how did I screw up?

$$\text{so it is } t = \sqrt{x} \quad \sqrt{4} = 2$$

$$\sqrt{1} = 1$$

$$f \quad \vec{F} = (x+y)\hat{i} + xy\hat{j}$$

C broken line



$x=t$ ← how do we pick

$y=$

Well for this write it out first
 ↳ no clear eq of the line

$$\int_{C_1} (x+y)dx + xydy + \int_{C_2} (x+y)dx + xydy$$

~~$x=0$~~ ~~$dx=0$~~ ~~$x=2$~~ ~~$dy=0$~~

$$\int_{C_1} x dx + \int_{C_2} 2y dy$$

$$\int_0^2 x dx + \int_0^1 2y dy$$

What did I do now?
 - went wrong way
 in picture

~~$$\frac{x^2}{2} \Big|_0^2 + 2 \frac{y^2}{2} \Big|_0^1$$~~

~~$$\frac{2^2}{2} + \frac{2}{2} = 0$$~~

~~$$2 + 1$$~~

~~$$(3)$$~~

$$\frac{x^2}{2} + 2x \Big|_0^1$$


$$\left(\frac{5}{2}\right)$$



So far have not gotten a single problem, but gotten better

2. For the following fields and curves eval with geometry, not calculation

a) $\vec{F} = x\hat{i} + y\hat{j}$



So closed path so 0

$\vec{F}$ points radially outwards
tangent always perpendicular to radius
 $\vec{F} \cdot \vec{t} = 0$

$$\int_C \vec{F} \cdot d\vec{r} = \int_C \vec{F} \cdot \vec{t} \, ds = 0$$

I thought it was just because it was a closed circle

only true if $\vec{F} = \nabla f$

which means that $f_{xy} = f_{yx}$?

and/or means field is conservative, or not dependent on xy

check

In this case the field is constant as you walk along the circle - which is the argument they gave.

b) $\vec{F} = y\vec{i} - x\vec{j}$ C is counterclockwise circle


$\vec{F}$ always tangent to circle in clockwise
 $\vec{F} = -a\vec{t}$ so is that the parameterization

$$\int_C \vec{F} \cdot d\vec{r}$$

$$\int_C F \cdot \vec{t} ds$$

$$= - \int_C a ds$$

$= -2\pi a^2$ so that is the line integral
adding up force at each pt




Oh

mn
→
what's that?

3. Let $F = \hat{i} + \hat{j}$, how would you place a directed line segment C of length 1 so that $\int_C \vec{F} \cdot d\vec{r}$ would be a

a) maximum



Make it as large as possible, as far away from the origin as possible.

Maximum C if C in dir of field
 $C = \frac{\hat{i} + \hat{j}}{\sqrt{2}}$

Wait where did we cover this?
 It sounds familiar from physics

- what is the

$\sqrt{2}$


b) minimum


dir opposite field $C = \frac{-\hat{i} - \hat{j}}{\sqrt{2}}$

c) zero?

perpendicular to the field $C = \pm \frac{\hat{i} - \hat{j}}{\sqrt{2}}$

But let's look at other regions





But then how do we know where it starts
- oh it also says of height 1

Is that picture even what a field is?

Well direction $\rightarrow \frac{\langle x, y \rangle}{r}$ - so that is what they did

So their max ans was $\nearrow T + J$

min $\swarrow -T - J$

zero $\searrow T - J$ or $\nearrow -T + J$

So that was kinda what I drew
- quadrants 2, 3, 4 may have been wrong



better

yeah that was direction

it was not $x\uparrow + y\uparrow$
which I drew

↳ think how we are doing in physics

d) What is max + min value of integral

$\rightarrow$

$$\begin{aligned} \text{max} &= \sqrt{2} \\ \text{min} &= -\sqrt{2} \end{aligned}$$

right, same but opposite d'r

$$\int_C \vec{F} \cdot d\vec{r} = \pm |\vec{F}| \cdot |\vec{C}| = \pm \sqrt{2}$$

So what did they do
well did integral
but over what?



Lecture 21

Gradient fields, conservative fields, path-independent
Fundamental theorem of calculus for line integrals

4C-1 Let $f(x, y) = x^3 y + y^3$
 $C = y^2 = x$ between $(1, -1)$, $(1, 1)$ upwards

a) Calculate $F = \nabla f$



$$\int_C x^3 y dx + y^3 dy$$

Now set some parameter
 $x = t^2$ ← how do we choose these?
 $y = t^3$

Wait don't we just want gradient

$$\langle f_x, f_y \rangle \quad (\checkmark)$$

$$\langle 3x^2 y, x^3 + 3y^2 \rangle \text{ or } 3x^2 y \uparrow + (x^3 + 3y^2) \uparrow$$

b) Now calc $\int_C \vec{F} \cdot d\vec{r}$ directly

Use y as parameter
 $x = y^2$
 $y = y$

$$\text{thus } dx = 2y dy$$

So how do you decide what to pick again?
as parameter - don't say - ask

$$\int_C \vec{F} \cdot d\vec{r} = \int_{-1}^1 3(y^2)^2 y + 2y \, dy + [(y^2)^3 + 3y^2] dy$$

replace $x + y^5$

combine
simplify

$$\int_{-1}^1$$

$$6y^6 \, dy + y^6 + 3y^2 \, dy$$

$$\int_{-1}^1$$

$$7y^6 + 3y^2 \, dy$$


$$\left. \frac{7y^7}{7} + \frac{3y^3}{3} \right|_{-1}^1$$

$$y^7 + y^3$$

$$1^7 + 1^3 - [(-1)^7 + (-1)^3]$$

~~4~~ addition error

- ii) by using path independence to replace C w/
a simpler path



so new path is 2

$$\int \vec{F} \cdot d\vec{r}$$

$$x^3 y + y^3 \cdot 2$$

what do I do
w/ this?

They still have some formal process

Use y as parameter

$$C_2: x=1 \quad dx=0$$

$$\int_{-1}^1 (1+3y^2) dy$$
$$y + y^3 \Big|_{-1}^1$$

(4)

← where all in world
do they get this??

From here
need to describe path in
terms of parameter

iii) By FTC for line integrals

$$\int_C \vec{F} \cdot d\vec{r} = f(x, y) \Big|_{P_0}^{P_1}$$
$$= f(x_1, y_1) - f(x_0, y_0)$$

?? So how in all world does that work

$$\int_C \nabla f \cdot d\vec{r} = f(B) - f(A)$$

$$A = (1, -1)$$

$$B = (1, 1)$$

$$\int_C = (1+1) - (-1-1)$$

(4)

that was easy - but what
did they do?

- will rest after office hrs

2. Let $f(x, y) = x e^{xy}$
 $C = 1/x$ from $(1, 1)$ to $(0, \infty)$

So gradient field means constant/conservative field = means path independence

$$\begin{aligned} f_x &= 1 e^{xy} & \langle e^{xy}, 0 \rangle \\ f_y &= 0 \end{aligned}$$

Are these type of problems solved any different?

Need a parameter

$$\begin{aligned} y &= 1/x \\ x &= t \\ y &= 1/t \end{aligned}$$

Now dx, dy

$$dx = 1 dt$$

$$dy = -\frac{1}{t^2} dt \quad \checkmark \quad (\text{they do } x, \text{ but it's comparable})$$

Ask: how
 do gradient field
 problems differ
 from other
 problems

Now write with this

$$\int_C x e^{xy} dx \quad \leftarrow ??$$

Wrong $\nabla f = x y e^{xy} + e^{xy} \hat{i} + x^2 e^{xy} \hat{j}$

? where in all world did they get this??

$$\int_C \vec{F} \cdot d\vec{r}$$

$$\int_1^0 (e + e) dx + (x^2 e) \cdot \frac{-dx}{x^2}$$

So they must have done
written in gradient field

$$\int_1^0 x y e^{xy} + e^{xy} \uparrow dx + x^2 e^{xy} \uparrow dy$$

replaced w/ parameters

$$\int_1^0 t \cdot \frac{1}{t} e^{t \cdot \frac{1}{t}} + e^{t \cdot \frac{1}{t}} dt + t^2 e^{t \cdot \frac{1}{t}} \cdot -\frac{1}{t^2} dt$$

$$\int_1^0 e + e dt - t^2 e \cdot \frac{1}{t^2} dt$$

$$\int_1^0 2e dt - e dt$$

$$\int_1^0 e dt \text{ can't do}$$

$$2et - et \Big|_1^0$$

$$- [2e - e]$$

$$(-e)$$



Once I got started, I got it
One more like this

3. $f(x, y) = \sin x \cdot \cos y$

$$f_x = \cos x \cos y$$

$$f_y = -\sin x \sin y$$

$$\nabla f = \langle \cos x \cos y, -\sin x \sin y \rangle \quad \checkmark$$

$$x = \sin x \quad \text{e? Is that right}$$

$$y = \cos y \quad \text{- do they have to work out to something}$$

$$dx = \cos x$$

$$dy = -\sin y$$

$$\int_C \cos x \cos y \, dx + -\sin x \sin y \, dy$$

$$\int_C \cos(\sin x) \cos(\cos y) \cdot \cos x - \sin(\sin x) \sin(\cos y) \cdot \sin y$$

$$\int_C \cos(\sin x) \cdot \cos^2 y = \cos x + \sin^2 x \sin(\cos y) \sin y$$

$$\int_C \cos(\sin x) \cdot \cos x + \sin(\cos y) \sin y$$

No limits given

∴ Is that correct - can't go further

Path independent of course

$$\int \sin x, \cos, -\sin x_0 \cos y_0$$

FTC of line integrals

↳ oh the really easy way to do it

The difference on right hand side is maximized
if $\sin x_1 \cos y_1$ is maximized and
 $\sin x_0 \cos y_0$ is minimized - yeah duh

$$\text{Since } |\sin x \cos y| \leq |\sin x| |\cos y| \leq 1$$

max difference is $1 - (-1) = 2$

↳ Running from $(-\frac{\pi}{2}, 0)$ to $(\frac{\pi}{2}, 0)$ ✓

Lecture 22 Criterion for conservative fields, potential functions

4C-5a via both methods 1+2

For each of the following tell for what value of the constants the field will be a gradient field and for this value find a corresponding mathematical potential function.

$$\vec{F} = (y^2 + 2x)\vec{i} + axy\vec{j} = \vec{\nabla}f$$

$\vec{F}$ is gradient field if $M_y = N_x$

$$\begin{array}{ll} M_y = 2y & 2y = ay \\ N_x = ay & a = 2 \end{array}$$

Ok so now we know what a is

$$\vec{F} = (y^2 + 2x)\vec{i} + 2xy\vec{j}$$

Potential function $= f(x, y) = xy^2 + x^2 + C$

what is?

that long formula?


or

$$\begin{aligned} f(P_1) &= f(P_0) + \int_{P_0}^{P_1} M dx + N dy \\ &= f(P_0) + \int_{P_0}^{P_1} y^2 + 2x dx + 2xy dy \end{aligned}$$

need to find this
- can graph

this is
all method 1

Gradient field so path independent



$$= \int_{C_1} (x^2 + y^2) dx + 2xy dy \quad \substack{y=0 \quad dy=0} + \int_{C_2} (x^2 + y^2) dx + 2xy dy \quad \substack{x=x_1 \quad dx=0}$$

$$\int_0^{x_1} x^2 dx + \int_0^{y_1} 2x_1 y dy$$

$$\left. \frac{x^3}{3} \right|_0^{x_1} + \left. \frac{2x_1 y^2}{2} \right|_0^{y_1}$$

$$\frac{x_1^3}{3} + x_1 y_1^2$$

should be x^2

$$x^2 + xy^2 + c$$

-close, but why not exact

Now Method 2

Handwritten text at the top of the page, possibly a title or header.

Handwritten text in the upper middle section.

Handwritten text in the middle section, possibly a paragraph or list item.

Handwritten text in the lower middle section.

Handwritten text in the lower middle section, possibly a list or table.

Handwritten text in the lower middle section.

Handwritten text in the lower middle section.

Handwritten text in the lower middle section.

Handwritten text in the lower middle section.

Handwritten text in the lower middle section.

Handwritten text in the lower middle section.

Handwritten text at the bottom of the page.

$$f = \int y^2 + 2x \, dx$$

$$f = xy^2 + x^2 + \boxed{g(y)}$$

$$f = \int 2xy \, dy$$

$$f = xy^2$$

$$f_y = 2x = g'(y)$$

$$g'(y) = y$$

$$g(y) = \frac{y^2}{2}$$

$$f(x, y) = xy^2 + x^2 + \frac{y^2}{2} \quad (\checkmark) \text{ checks out cool}$$

like method 2 better - faster



6a. Decide which of the following differentials is exact. For the one that is exact, express it in terms of df .

$$y \, dx - x \, dy$$


$$\text{So is it } \nabla f = \langle y, -x \rangle$$

Method 1

$$f(P) = f(P_0) + \int_{P_0}^P M \, dx + N \, dy$$

$$f(P_0) + \int_{P_0}^P y \, dx - x \, dy$$

Now need to draw a path independent way



(Can always use this method?)

$$\int_{c_1} y dx - x dy \quad + \quad \int_{c_2} y dx - x dy$$

$x=0 \quad dy=0$
 $x=x_1 \quad dx=0$

$$0 \quad + \quad \int_0^{y_1} -x_1 dy$$

Now integrate

$$-x_1 y \Big|_0^{y_1}$$

$$\boxed{-x_1 y_1}$$

actually that is easier to figure out, but more writing

Method 2

$$f_x = y$$

$$f_y = -x$$

$$f = xy + g(y)$$

$$f = -x + g(y)$$

$$g(y) = y$$

$$g'(y) = 1$$

$$\boxed{f = xy + y}$$

$ydx - xdy$ is not exact
since $M_y = 1$ $N_x = -1$

(so that is why I had sign problems)

So is it just undanswerable.

✓

6b. $y(2x+y)dx + x(2y+x)dy$

Method 1

$$f(p) = f(p_0) + \int_{p_0}^p M dx + N dy$$

$$\langle y(2x+y), x(2y+x) \rangle$$

$$\int_{C_1} y(2x+y)dx + x(2y+x)dy + \int_{C_2} y(2x+y)dx + x(2y+x)dy$$

$y=0 \quad dy=0$ $x=x_1 \quad dx=0$

$$0 + \int_0^{x_1} 0 + x_1(2y+x_1)dy$$

$$\int_0^{x_1} 2x_1 y + x_1^2 dy$$

$$\left. \frac{2x_1 y^2}{2} + x_1^2 y \right|_0^{x_1}$$

$$\boxed{x_1 y_1^2 + x_1^2 y_1 + c} \quad (\checkmark) \text{ Perfect}$$

Method 2

$$\langle y(2x+y), x(2y+x) \rangle$$

This will be complicated

$$f = \int y(2x+y) dx$$

$$f = x(x+y)y + g(y)$$

$$\begin{aligned} f_y &= x(2y+x) \\ f &= y(y+x)x \\ &= x(x+y)y \end{aligned}$$

How is this going to go?

$$\begin{aligned} x^2 + xy + xy + y^2 + g(y) \\ x^2 + 2xy + y^2 + g(y) \end{aligned}$$

- Not close to what I got above

Did not check if exact
 $M_y = 2x + 2y = N_x$

But what went wrong here?


✓

Should do more
 practice to mem-
 orize test cases

✓

Part 2

(working outside!) la, Write down in x - y coordinates the vector field $\vec{F}$ whose vector at $P(x,y)$ runs from P to $L \rightarrow x+y=1$



So we are going from any point $\rightarrow$ down
but the vector is not $\vec{F}$
instead it is a curve C which runs vertically

$$\int_C \vec{F} \cdot d\vec{r}$$


r is the curve
 $- < 0, y >$

oliver otl

$$< 0, y_0 - (1-x) >$$

from y_0 to $1-x$

$$\vec{F} = < 0, 1-x-y >$$



Or is it just a field - which has magnitude + direction

What would magnitude be

$$\sqrt{0^2 + (y_0 - (1-x))^2}$$

$$y_0 - (1-x) < 0, y_0 - (1-x) >$$

b. Show that w/o calculation - for this field

$\int_C \mathbf{F} \cdot d\mathbf{r} \neq 0$ if C is a right triangle



-2.5

Well the field is non constant. It is dependent on x in the calculation of it. This means that the line integral along a closed path is not necessarily 0.

? should we write anything else

- there are parts which are perpendicular to the field - these don't count at all

only the degree each line is parallel counts (dot product) There are segments of the line which meet this criteria

extright


2. Imagine the x -axis is an ∞ long heavy wire of uniform density. The gravitational force exerted on it is

$$\vec{F}(y, z) = -\frac{(y\vec{j} + z\vec{k})}{y^2 + z^2} \quad (y, z) \neq (0, 0)$$

C is a positive constant $= 1$

Find the integral - by using a parametrization of C - the work done by moving field along some path in yz plane

- a) half-line $z=1$ $y \geq 0$ start $(0, 1)$



$$\vec{r} = \langle x, 0 \rangle \times [0, \infty)$$

$$\begin{aligned} y &= t & dy &= 1 \\ z &= 1 & dz &= 0 \end{aligned}$$

$$\int_0^{\infty} \vec{F} \cdot \vec{T}$$

$$\frac{\langle -y, -z \rangle}{y^2 + z^2} \cdot \langle 1, 0 \rangle$$

$$\frac{\langle -1, -1 \rangle}{t^2} \cdot \langle 1, 0 \rangle$$


$$-\frac{x}{x^2} \cdot 1$$

$$\int_0^{\infty} -\frac{1}{x} dt \rightarrow -\ln(x) \Big|_0^{\infty}$$

$$-\ln(\infty) + \ln(0)$$

∞ $\checkmark$ not closed loop $\checkmark$

b) circle w/ radius a , centered at origin, counterclockwise



so this one should be $\cos, \sin$

$$a^2 = y^2 + z^2$$

$$a = \sqrt{y^2 + z^2}$$

$$y = \cos t \quad dy = -\sin t$$

$$z = \sin t \quad dz = \cos t$$

$$\int_0^{2\pi} \frac{\langle -1y, -1z \rangle \cdot \langle dx, dy \rangle}{y^2 + z^2}$$

$$\int_0^{2\pi} \frac{\langle -\cos t, -\sin t \rangle \cdot \langle -\sin t, \cos t \rangle dt}{\cos^2 t + \sin^2 t}$$

$$(-\cos t \cdot -\sin t) + (-\sin t \cdot \cos t) dt$$

$$\int_0^{2\pi} -2 \sin t \cos t dt$$

$$+ \cos^2 x \Big|_0^{2\pi}$$


$$= \cos^2(2\pi) - [\cos^2(0)]$$

$$= 1 - 1$$

0




Closed loop - field prob constant
Something about field being tangent



- in that model of vector

$$\text{dir}(-\hat{y}, \hat{z})$$

c) line from (0,1) to (1,0)



$$r = \frac{\langle 0-1 \rangle}{\langle 1-0 \rangle} = \frac{\langle \hat{y}, -\hat{z} \rangle}{\langle 0, 1 \rangle}$$

~~$$y = t$$~~
~~$$z = -t + 1$$~~

what is parameter

~~$$y = t$$~~
~~$$z = -t + 1$$~~

✓ bingo (took like 10 min to find)
hate these problems

$$dy = 1$$

$$dz = -1$$

~~parameter~~
oliver OH

$$\int \frac{\langle -y, z \rangle}{y^2 + z^2} \cdot \langle dy, dz \rangle$$

$$\int_0^1 \frac{\langle t, -t+1 \rangle}{t^2 + (-t+1)^2} \cdot \langle 1, -1 \rangle dt$$

$$\begin{aligned} & t^2 + (-t+1)^2 \\ & t^2 + t^2 - 2t + 1 \\ & 2t^2 - 2t + 1 \end{aligned}$$

$$\int_0^1 \left(\frac{t}{2t^2 - 2t + 1} \cdot 1 \right) + \left(\frac{-t+1}{2t^2 - 2t + 1} \cdot -1 \right) dt$$

$$\int_0^1 \frac{t + t - 1}{2t^2 - 2t + 1} dt$$

$$\int_0^1 \frac{2t - 1}{2t^2 - 2t + 1} dt$$

$$\frac{-\ln(2t^2 - 2t + 1) + 2 \tan^{-1}(2t - 1)}{2} \Big|_0^1$$

$$\cancel{\left(\frac{\pi}{4} \right)} \quad (0)$$

$$(-0.5)$$

should be same for a, b, c

3a Let $\vec{F} = -\nabla(\ln r)$ where r is the distance from the origin in the y, z plane. Show that F is field of problem 2 when $c=1$

So we need to show $M_y = N_x$
or $M_z = N_y$ in this case

$$\vec{F} = \frac{y \hat{j} + z \hat{k}}{y^2 + z^2}$$

$$\vec{F} = \left\langle \frac{y}{y^2 + z^2}, \frac{z}{y^2 + z^2} \right\rangle$$

$$y(y^2 + z^2)^{-1}$$

$$M_z = \frac{-2zy}{(y^2 + z^2)^2} \quad N_y = \frac{-2y \cdot z}{(y^2 + z^2)^2}$$

know
already $\rightarrow$

They are the same
So is a gradient field

Or did I screw this up?

M and N are from $\nabla f = \langle M, N \rangle$
but I calculated them
-so its ok - should be good

Wait am I good?

~~Take
gradient~~

~~$$\left\langle \frac{-(y^2 - z^2)}{(y^2 + z^2)^2}, \frac{-(z^2 - y^2)}{(y^2 + z^2)^2} \right\rangle$$~~

Should = $\ln r$
 $r = \sqrt{y^2 + z^2}$

oiler
OH

~~$$\left\langle \frac{-y^2+z^2}{r}, \frac{-z^2+y^2}{r} \right\rangle$$~~

Want $\nabla(\ln r)$
 $\nabla(\ln \sqrt{y^2+z^2})$

g_1

g_2

$$\left\langle \frac{y}{y^2+z^2}, \frac{z}{z^2+y^2} \right\rangle \quad \checkmark$$

solved now

matches $\vec{F}$ $\odot$

$\vec{F}$ = field
 f = function
 $\vec{F} = \nabla(f)$

$$\left\langle \frac{y}{r^2}, \frac{z}{r^2} \right\rangle$$

3b Show that for any path C joining 2 points P_1 and P_2 in the x, y plane the value of $\int_C F \cdot dr$ depends on $\frac{r_2}{r_1}$ where r_i is

distance from origin.
What pts P_i should be excluded?

↳ this first

P_0 which is $(0,0)$ should be ✓

Well FTC (line integrals) says that

$$\int_{P_1}^{P_2} F \cdot d\vec{r} = f(y_2, z_2) - f(y_1, z_1)$$

where
gradient
field

Well if you plug points in

$$y_2, z_2 \rightarrow \ln \sqrt{y_2^2 + z_2^2} = \ln r_2$$

$$y_1, z_1 = \ln \sqrt{y_1^2 + z_1^2} = \ln r_1$$

$$= \ln(r_2) - \ln(r_1) \quad \checkmark$$

$$= \frac{r_2}{r_1}$$

rule of $\ln$

$$\ln a + \ln b = \ln ab$$

$$\ln a - \ln b = \ln \frac{a}{b}$$

oh I'm getting confused.

oliver oh
fixed
self

c) Let C be a directed line segment of length 1


i) If $\int_C \vec{F} \cdot d\vec{r} = 0$, how is C placed?

C must be placed perpendicular to the field ✓


ii) What position of C gives $\int_C \vec{F} \cdot d\vec{r}$ the largest value?

In the direction of the field, parallel with it

-0.5



4. Let $\vec{F} = \nabla(x^2y - 3xy^2)$ Let C be $1/4$ ellipse $9x^2 + 4y^2 = 1$ from $x \rightarrow y$ axis



a) Write integral $\int_C \vec{F} \cdot d\vec{r}$ in the $\int_C Mdx + Ndy$ form

$$f_x = 2xy - 3y^2 \quad \text{copy error}$$

$$f_y = x^2 - 3x \cdot 2y$$

$$\vec{\nabla} f = \langle 2x - 3y^2, x^2 - 6xy \rangle$$

$$\int_C (2x - 3y^2) dx + (x^2 - 6xy) dy$$

no parametrizing
done, right?

✓ correct
- oliver oh

b) Evaluate using FTC + Path Independent

$$\text{FTC} \rightarrow f(x_2, y_2) - f(x_1, y_1)$$

$$x_2^2 y_2 - 3x_2 y_2^2 - [x_1^2 y_1 - 3x_1 y_1^2]$$

$$x_2 = 0$$

$$x_1 = 1/3$$

$$y_2 = 1/5$$

$$y_1 = 0$$


$$0 - 0 - 0 - 0$$

$$\textcircled{0}$$


✓

Capital + lower
f confusing
me - see
chart
a few pg
back

b) Path independence



Lets just split it



$$\int_{C_1} (2xy - 3y^2) dx + (x^2 - 6xy) dy + \int_{C_2} (2xy - 3y^2) dx + (x^2 - 6xy) dy$$

Can keep dx ?
- pretty sure

$$x=0$$

$$dx = 0$$

$$y=0 \quad dy=0$$

$$\int_{C_1} 0 + 0 dy + \int_{C_2} 2x dx + 0$$

$$\int_0^{1/3} 2x dx = \left. \frac{2x^2}{2} \right|_0^{1/3} = \frac{1}{6}$$

⓪

✓ copy error in part a

$$s \quad \downarrow \frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y}$$

$$5. \quad F = \underbrace{xy}_{M} \hat{i} + \underbrace{x^2}_{N} \hat{j}$$

a) Verify $\vec{F}$ is not conservative

FYI

$$M_y = f_{xy}$$

You can see that x and y are terms of the field this means it is not conservative

~~$$f_x = y \hat{i} + 2x \hat{j}$$~~

~~$$f_{xy} = 1 \hat{i}$$~~

~~$$f_y = x \hat{i}$$~~

~~$$f_{yx} = 1 \hat{i}$$~~

But they are =


$$M = f_x$$

You can take the gradient of a vector field w/o it being a gradient field

$$So \quad M_y = x \hat{j} \neq N_x = 2x \hat{j} \quad \checkmark$$

b) Try to find a potential function for $\vec{F}$ by using method 1. What goes wrong?

$$\text{Potential function} = \int_{P_0}^P M dx + N dy$$



$$\int_{c_1} xy \, dx + x^2 \, dy + \int_{c_2} xy \, dx + x^2 \, dy$$

$y=0 \quad dy=0$
 $x=x_1 \quad dx=0$

$$0 + \int_0^{x_1} 0 + x^2 \, dy$$

$$x^2 y \Big|_0^{y_1}$$

$$\boxed{x^2 y}$$

✓

- But I found it?? What was supposed to go wrong? (Now, if you take $\nabla(x^2y)$, you don't get F)

probably blew past some assumption I was supposed to make

but as I am doing the integral it is not an even field

Or is it in the way I write the integral w/ $Mdx + Ndy$

Yeah I assumed

$$\vec{F} = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right\rangle \text{ which was invalid}$$

-only true for gradient fields

But then why did we bother trying it?

Or is it that result = variable < but that is ok??

why is it working

c. Now Method 2

-this will prob fail more

$$f_x = xy$$

$$f_y = x^2$$

$$f_x = \int xy \, dx$$

$$f = x^2 y + g(y)$$

$$f_y = x^2 +$$

$$= x^2 + g'(y)$$

$$g'(y) = 0$$

$$g(y) = C$$

$$f(x, y) = x^2 y + C$$


↑

same answer

Why is it working?


It's not, again $\nabla(x^2 y)$ is not F

6. The most 2D fields whose direction is radially outward and whose magnitude depends only on r ($r = \sqrt{x^2 + y^2}$) is



$$\vec{F} = f(r)(x\hat{i} + y\hat{j})$$

$\uparrow$ varies
magnitude



- a) Assuming $f(r)$ is differentiable for all r , show that any such $\vec{F}$ is conservative

$$\frac{\partial r}{\partial x} = \frac{x}{r} \quad \frac{\partial r}{\partial y} = \frac{y}{r}$$

Well $f(r)$ just disappears. coefficient - No, indirectly depends on x, y

$$\frac{\partial F}{\partial r} = \frac{\partial x}{\partial r} + \frac{\partial y}{\partial r}$$

(-1.5)

- but how does that use our hint?

well $\vec{F}$ is conservative if $\vec{F} = \nabla f$

$$\text{or } f_x = 1 \hat{i} \\ f_y = 1 \hat{j}$$

$$\langle 1, 1 \rangle$$

$$N_y = 0 \quad \checkmark \\ N_x = 0 \quad \checkmark$$

$$\text{or } f_x = \frac{\partial f}{\partial x}$$

$$\text{want } \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial r} = \frac{\partial f}{\partial r}$$

or is it $\frac{\partial f}{\partial r}$
so it would be

$$h = \left\langle 1 + \frac{c}{x}, 1 + \frac{c}{y} \right\rangle$$

And check

$$\begin{aligned} h_y &= 0 \text{ still same?} \\ h_x &= 0 \end{aligned}$$

??

Does that answer the question?


- b) Find a potential function of the form $g(r)$
if $f(r) = r^n$ where n is $\oplus \ominus$ integer

(Hint: Solve $\nabla g(r) = \vec{F}$ for the function $g(r)$. One value of n has to be handled differently from all of the others)
p so $\int$ back

1.5
see solu

$$g(r) = \int_{c_1} r^n x dx + r^n y dy + \int_{c_2} r^n x dx + r^n y dy$$

$y=0 \quad dy=0$ $x=x, \quad dx=0$



$$\int_{c_1} \sqrt{x^2+y^2}^n x dx + \sqrt{x^2+y^2}^n y dy$$

$$+ \int_{c_2} \sqrt{x^2+y^2}^n x dx + \sqrt{x^2+y^2}^n y dy$$

$$\int_0^{x_1} x^{n+1} dx + \int_{c_2} \sqrt{x_1^2+y^2}^n y dy$$

$$\frac{x^{n+2}}{n+2} \Big|_0^{x_1} + \int_0^{y_1} (x^2+y^2)^{n-1} y dy$$

$$\frac{x^{n+2}}{n+2} + \frac{(y^2+x^2)^n}{2n} \Big|_0^{y_1}$$


Well when $n-2=0$ it will be different since

$$n=2$$


the answer will always be 1. Is this what
is different?

11


what is different




$$\vec{F} = \vec{PQ} = (1-x-y)\hat{j} \quad \text{or} \\ = \langle 0, 1-x-y \rangle$$




On $\vec{PR}$, $\vec{F} \cdot \vec{T} = 0$ since $\vec{F} \perp \vec{PR}$
 On $\vec{RQ}$, $\vec{F} \cdot \vec{T} = 0$ since $\vec{F} = \vec{0}$ on $\vec{RQ}$
 On $\vec{QP}$, $\vec{F} \cdot \vec{T} < 0$ since $\vec{F}$ and $\vec{QP}$ have opposite directions.
 $\therefore \int_C \vec{F} \cdot \vec{T} ds = \int_{\vec{QP}} \vec{F} \cdot \vec{T} ds < 0.$



$$\text{Work} = \oint_C \frac{y dy + z dz}{y^2 + z^2} = - \int_0^\infty \frac{y dy}{y^2 + 1} \\ = - \frac{1}{2} \ln(y^2 + 1) \Big|_0^\infty = -\infty$$



Using same integral as in (a), $\otimes$
 $\text{Work} = - \int_0^{2\pi} \frac{-a^2 \sin t \cos t + a^2 \sin t \cos t}{a^2} dt \\ = - \int_0^{2\pi} 0 dt = 0.$



$$\text{Work} = - \int_0^1 \frac{y dy - (1-y) dy}{2y^2 - 2y + 1} \quad [\otimes] \\ = - \int_0^1 \frac{(2y-1)}{2y^2 - 2y + 1} dy \\ = - \frac{1}{2} \ln(2y^2 - 2y + 1) \Big|_0^1 = 0 - 0 = 0.$$

3 a) $r = \sqrt{y^2 + z^2}$
 $-\nabla \ln(y^2 + z^2)^{1/2} = -\frac{1}{2} \nabla \ln(y^2 + z^2) \\ = -\frac{1}{2} \left\langle \frac{2y}{y^2 + z^2}, \frac{2z}{y^2 + z^2} \right\rangle \\ = - \left(\frac{y\hat{i} + z\hat{j}}{y^2 + z^2} \right)$

b) Using the Fundamental Thm of Calculus for line integrals,

$$\int_{P_1}^{P_2} \vec{F} \cdot d\vec{r} = - \ln r \Big|_{P_1}^{P_2} = - [\ln r_2 - \ln r_1] \\ = - \ln(r_2/r_1), \quad \text{where } r_i = r(P_i).$$


Have to exclude pts where $r(P_i) = 0$, i.e., $P_i \neq (0, 0)$.

c) (i) C is placed so that its endpoints are equally distant from (0,0).

(ii) There is no maximum value

for $\int_C \vec{F} \cdot d\vec{r}$:

if we place C as shown (radially)




then $\frac{r_2}{r_1} \rightarrow 0$ as $P_2 \rightarrow (0, 0)$

$$\therefore \int_C \vec{F} \cdot d\vec{r} = - \ln\left(\frac{r_2}{r_1}\right) \rightarrow \infty \text{ as } P_2 \rightarrow (0, 0).$$


4 a) $\nabla(x^2y - 3xy^2) = \langle 2xy - 3y^2, x^2 - 6xy \rangle$

$$\therefore \int_C \vec{F} \cdot d\vec{r} = \int_C (2xy - 3y^2) dx + (x^2 - 6xy) dy$$



Using the F.T.: $\int_C \vec{F} \cdot d\vec{r} = x^2y - 3xy^2 \Big|_{(1/3, 0)}^{(0, 1/2)} \\ = 0 - 0 = 0$

Using the path:



$$C_1: x = x, y = 0$$

$$C_2: x = 0, y = y$$


$$\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_1} 0 \cdot dx = 0, \quad \int_{C_2} \vec{F} \cdot d\vec{r} = \int_{C_2} 0 \cdot dy = 0.$$

5) a) $\vec{F} = xy\hat{i} + x^2\hat{j}$

$$\frac{\partial(xy)}{\partial y} = x, \quad \frac{\partial(x^2)}{\partial x} = 2x$$

Since these are not equal, the field is not conservative.

b) Method 1:



$$C_1: y=0, x=x_1$$

$$C_2: x=x_1, y=y_1$$

$$f(x_1, y_1) = \int_{C_1+C_2} xy dx + x^2 dy$$

$$\int_{C_1} = \int_0^{x_1} 0 dx = 0$$

$$\int_{C_2} = \int_0^{y_1} x_1^2 dy = x_1^2 y_1$$

Since $dx=0$ on C_2

$$\therefore f(x_1, y_1) = x_1^2 y_1, \text{ or } \boxed{f = x^2 y}$$

$$\text{but } \nabla(x^2 y) = 2xy\hat{i} + x^2\hat{j} \neq \vec{F}$$

Method 2:

$$\frac{\partial f}{\partial x} = xy \text{ (from } \vec{F})$$

$$\therefore f = \frac{1}{2}x^2 y + g(y)$$

$$\therefore \frac{\partial f}{\partial y} = \frac{1}{2}x^2 + g'(y)$$

$$= x^2 \text{ (from } \vec{F})$$

$$\text{But } x^2 \neq \frac{1}{2}x^2 + g'(y)$$

for any $g(y)$, since

$$\frac{1}{2}x^2 \neq g'(y)$$

function of
x alone

function
of y alone

6) $\vec{F} = f(r)(x\hat{i} + y\hat{j})$

a) Using $\frac{\partial r}{\partial x} = \frac{x}{r}, \quad \frac{\partial r}{\partial y} = \frac{y}{r}$:

$$\frac{\partial}{\partial y} f(r)x = f'(r) \frac{\partial r}{\partial y} \cdot x = f'(r) \cdot \frac{yx}{r}$$

$$\frac{\partial}{\partial x} f(r)y = f'(r) \frac{\partial r}{\partial x} \cdot y = f'(r) \cdot \frac{xy}{r}$$

Since these are equal, $\vec{F}$ is conservative

b) $\nabla g(r) = r^n(x\hat{i} + y\hat{j})$

$$\otimes \therefore g'(r) \frac{\partial r}{\partial x} = g'(r) \cdot \frac{x}{r} = r^n x$$

$$\otimes g'(r) \frac{\partial r}{\partial y} = g'(r) \cdot \frac{y}{r} = r^n y$$

These will be both satisfied if

$$\frac{g'(r)}{r} = r^n, \text{ so } g'(r) = r^{n+1}$$

$$g(r) = \boxed{\frac{r^{n+2}}{n+2}}$$

[if $n = -2$,

$$g'(r) = \frac{1}{r}, \text{ so } g(r) = \boxed{\ln r}]$$

Lecture 23

Green's Theorem

4/8

- next 3 lectures



- Hypothesis 1 - simple curve, does not overlap (self intersection)
- no end points "closed"
 - piecewise, smooth $\begin{cases} x = x(t) \\ y = y(t) \end{cases}$
 - differentiable except at certain points + shapes

The curve divides plane into 2 parts

- ∞ large area exterior
- enclosed region, finite area R
- hard to prove, but we don't care

curve must be oriented in a positive oriented

- aka go counterclockwise
- but sometimes that is not clear
- so the interior is always on your left

$$\vec{F} = M\hat{i} + N\hat{j}$$

$\hookrightarrow$ must be differentiable in R

Green's theorem relates the line integral around the curve

$\oint$ - closed path

$$\oint M dx + N dy \quad - \text{could be work integral}$$

Green's Theorem

$$\oint_C M dx + N dy = \iint_R (N_x - M_y) dA$$

$$\oint_C \vec{F} \cdot d\vec{r} = \iint_R \text{curl } \vec{F} \cdot d\vec{A}$$

$$\text{curl } \vec{F} = N_x - M_y$$

ex $\oint 3x^2 y^2 dx + 2x^2(1+xy) dy$




doing $\iint$ may be hard

So try Green's theorem

$$\iint_R \underbrace{4x + 6x^2 y}_{N_x} - \left[\cancel{6xy} x^2 \right] dA$$

$$4 \iint_R x dA$$

- could use com or avg value
over




$$\text{Average of } x = \frac{\iint_R x dA}{\text{avg } A = \pi a^2}$$

$$\iint_R x dA = (\text{avg of } x) \cdot \pi a^2$$

$$= 4\pi a^3$$

Some are more easily calculated as a line integral



did by
vector
methods
before

$$\rightarrow \begin{cases} x = a(\theta - \sin\theta) \\ y = a(1 - \cos\theta) \end{cases} \text{ deriv of}$$

Area under 1 arch


Area by line integral

$$- \int y dx = \iint_R 1 dA$$

Want $N_x - M_y = 1$

$\int x dy$ or $\frac{1}{2}$ (both) $\leftarrow$ what is this for??

$$\oint_{C_1+C_2} -y dx = \int_{C_1} -y dx + \int_{C_2} -y dx$$



$$0 = \int_0^{\pi} -a(1-\cos\theta)$$

$$dx = -y d\theta$$

$$= - \int_0^{2\pi} a^2 (1-\cos\theta)^2 d\theta$$


calculate

$$= -a^2 \int_{2\pi}^0 \left(\underset{2\pi}{1} - \underset{+}{2\cos\theta} + \underset{+}{\cos^2\theta} \right) d\theta$$

$$\boxed{3\pi a^2}$$

3 times the area of the circle

- very mysterious when it came out
- important
- not obvious



"Vertically simple" Δ

entering + leaving once each

Proof

- calculate both sides of the theorem
- double integrals + line integrals

$$\oint_C M dx = - \iint_R \frac{\partial M}{\partial y} dA \quad ??$$

On C_2, C_4 - vertical pieces
will contribute nothing to line integral
since $x = a$ or $x = b$ and $dx = 0$

$$\int_{C_1} M dx + \int_{C_2} M dx$$

$C_1: x = x$ can change to t - but harder
 $y = g(x)$

$$\int_a^b M(x, g(x)) dx + \int M(x, h(x)) dx$$

$$\oint M dx = \int_a^b [M(x, g(x)) - M(x, h(x))] dx$$

$$\iint_R \frac{\partial M}{\partial y} dy dx = \int_a^b \int_{g(x)}^{h(x)} \frac{\partial M}{\partial y} dy dx$$

$$\begin{aligned} & \text{Inner } M(x, y) \Big|_{g(x)}^{h(x)} \\ &= M(x, h(x)) - M(x, g(x)) \end{aligned}$$

Outer = same thing

So same answer

Lecture 24

Flux, Green's Theorem in normal form

4/4

Review

$$\vec{F} = M\vec{i} + N\vec{j}$$

force field

$$\oint \vec{F} \cdot \vec{t} ds$$

$\vec{t}$
unit
tangent

tangential
form



$$\oint_C \vec{F} \cdot d\vec{r} = \iint_R (\text{curl } \vec{F}) dA$$

$$\oint M dx + N dy = \iint_R (N_x - M_y) dA$$

Green's
Theorem

work
energy


why??

but how does path relate to inside of region

$\vec{F}$

velocity field

- less common in 1st year
- common in fluid mechanics



velocity of incompressible fluid


constant over time

- not a function of t

$$\vec{F}(x, y)$$

Flux

- analog to work



- net flow rate across C

- cm^2/sec


- face into C ?

- $\oplus$ flux is left to right

$\ominus$ " " right to left

- net = $\oplus - \ominus$ flows

a)



$\vec{F}$ is constant

- does not vary w/ position

resolve F into 2 components


- perpendicular + parallel to lines

- only the perpendicular $\vec{F}$ counts as flux

- in the $\hat{n}$ (unit normal) direction


$$\text{Flux across } C = (\vec{F} \cdot \hat{n}) |PQ|$$

b)



but now $\hat{n}$ keeps changing direction
magnitude still the same though

Divide curves into little pieces



$$\sum_{i=1}^n \vec{F}(x_i, y_i) \cdot \Delta r_i = \text{total flux}$$

now as $n(\# \text{ of pieces}) \rightarrow \infty$

$$= \int_C (\vec{F} \cdot \hat{n}) ds$$

usual notation - not all may agree
flux across C

$$\hat{t} = \left\langle \frac{dx}{ds}, \frac{dy}{ds} \right\rangle \quad \text{unit tangent vector}$$

$$\hat{n} = \left\langle -\frac{dy}{ds}, \frac{dx}{ds} \right\rangle$$



Note its on the other side as compared to start of year

- 2 different conventions for separate problems


$$\int_C \vec{F} \cdot \hat{n} ds$$

flux

$$= \int_C \left(\frac{M dy}{ds} - \frac{N dx}{ds} \right) ds$$

But can never use s as a parameter
- only in formulas

$$\text{Symbolic flux interval} = \int_C M dy - N dx$$



$\vec{F}$ is always out

$$\vec{F} = x\hat{i} + y\hat{j}$$

Flux over C

$$x = a \cos \theta$$

$$y = a \sin \theta$$

$$dx = -a \sin \theta$$

$$dy = a \cos \theta$$

$$\vec{F} = x\hat{i} + y\hat{j}$$

$$\text{flux over } C = \int_C x dy - y dx$$

$$= \int_0^{2\pi} (a^2 \cos^2 \theta + a^2 \sin^2 \theta) d\theta$$

$$= 2\pi a \cdot a = 2\pi a^2$$

$\uparrow$ length of circle
 $\uparrow$ velocity flow

Green's theorem $\rightarrow$ flux
"normal form of Green's theorem"

* notice the order *

$$\oint_C -N dx + M dy = \iint_R M_x - (-N)_y dA$$

applied to this case

- only 1 Green's theorem, but multiple meanings

$$\oint M dy - N dx = \iint (M_x + N_y) dA$$

$$\oint \vec{F} \cdot \hat{n} ds = \iint_R (\text{div } \vec{F}) dA$$

↑
divergence

Flux over $C =$?? have to know divergence
- but is physical interpretation

So what is div ?

- $M_x + N_y$


- but what does it mean?



$$f(x, y) = \text{div } \vec{F}$$

$$f(x_0, y_0) = ??$$

$$\oint_C \vec{F} \cdot \hat{n} ds = \text{div}(x\hat{i} + y\hat{j})$$
$$= 2 \quad \text{why?}$$



By Green

$$f(x, y) = \text{div } \vec{F}$$

$$f(x_0, y_0) = \vec{F}$$

$$f(x, y) \approx f(x_0, y_0)$$

By Greens

$$\oint_{C_0} \vec{F} \cdot \vec{n} \, ds = \iint_{R_0} f(x_0, y_0) \, dA$$

$$= \underset{\text{div}}{f(x_0, y_0)} \text{ area } R_0$$

$$\frac{(\text{div } \vec{F})_{(x_0, y_0)}}{f(x_0, y_0)} = \frac{\text{net flux across } C_0}{\text{Area of } R_0}$$

= rate (cm^2/sec) to which fluid is added to tank at (x_0, y_0) (or $\ominus$ if leaking)

Recitation

4/12

Lectures

Green's Theorem

$$\oint_{C \text{ closed}} M dx + N dy = \iint_R N_x - M_y dA$$



Consequences for fields

$$\text{"work"} \oint_C \vec{F} \cdot d\vec{r} = \iint_R \text{curl } \vec{F} dA$$

$$\text{"flux"} \oint_C \vec{F} \cdot \hat{n} ds = \iint_R \text{div } \vec{F} dA$$


$$d\vec{r} \equiv \langle dx, dy \rangle = \langle x', y' \rangle dt$$

$$\hat{n} ds = \langle dy, -dx \rangle = \langle y', -x' \rangle dt$$

$\hat{n}$ = normal unit vector

ex 1 Compute a line integral in 2 different ways

$$\vec{F} = x\hat{i} + xy\hat{j}$$



$$\text{Compute } I = \oint_C \vec{F} \cdot d\vec{r}$$


a) directly

b) using Green's Theorem

didn't start Pset
due to cpl
so very bad at
this

So need to cut the curve into 3 pieces

$$\oint_C \vec{F} \cdot d\vec{r} = \int_{C_1} + \int_{C_2} + \int_{C_3}$$



C₁ $\int_{C_1} \vec{F} \cdot d\vec{r} = 0$ because $x=0$ on C_1
 so $f=0$ so $\int_{C_1} \vec{F} \cdot d\vec{r} = 0$

C₂ $\int_{C_2} \vec{F} \cdot d\vec{r} =$ need to parametrize
 $x(t) = t$ or $x(x) = x$
 $y(t) = 0$ or $y(x) = 0$
 $\uparrow$ Oliver $\uparrow$ Prof. Mattick

$$d\vec{r} = \begin{pmatrix} x' \\ y' \end{pmatrix} dt = \begin{pmatrix} 1 \\ 0 \end{pmatrix} dt \quad t \in [0,1] \quad x \in [0,1]$$

$$\begin{aligned} \int_{C_2} \vec{F} \cdot d\vec{r} &= \int_{t=0}^{t=1} \begin{pmatrix} t \\ 0 \end{pmatrix} \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} dt \\ &= \int_0^1 t dt \\ &= \left. \frac{t^2}{2} \right|_0^1 \\ &= \frac{1}{2} \end{aligned}$$

C₃ $\int_{C_3} \vec{F} \cdot d\vec{r} =$ needs parametrization
 $x(t) = \cos t$
 $y(t) = \sin t$
 - others are possible - this is best

$$\begin{aligned}
 \int_{C_3} \vec{F} \cdot d\vec{r} &= \int_0^{\pi/2} \langle \cos t, \cos t \sin t \rangle \cdot \langle -\sin t, \cos t \rangle dt \\
 &= \int_0^{\pi/2} -\cos t \sin t + \cos^2 t \sin t dt \\
 &= \left. \frac{\cos t^2}{2} + \frac{\cos t^3}{3} \right|_0^{\pi/2} \quad \text{same sort of identity} \\
 &= -\frac{1}{2} + \frac{1}{3} \\
 &= -\frac{1}{6}
 \end{aligned}$$

$$\begin{aligned}
 I &= \int_{C_1} + \int_{C_2} + \int_{C_3} \\
 &= 0 + \frac{1}{2} - \frac{1}{6} = \frac{1}{3}
 \end{aligned}$$

b)
$$I = \oint_C \vec{F} \cdot d\vec{r} = \iint \text{curl } \vec{F} \, dA$$

green's theorem

↓ what is it

$$= \iint \left(\frac{\partial}{\partial x} (xy) - \frac{\partial}{\partial y} (x) \right) =$$

$$= \iint_R y \, dx \, dy$$

$$= \int_0^1 \int_0^{\sqrt{1-y^2}} y \, dx \, dy$$

← look at picture 

$$= \int_{x=0}^1 \frac{y^2}{2} \Big|_0^{\sqrt{1-x^2}} dx$$

$$= \int_0^1 \frac{1-x^2}{2} dx$$


$$= \frac{x^2}{2} - \frac{x^3}{6} \Big|_0^1$$

$$= \frac{1}{2} - \frac{1}{6}$$

$$= \left(\frac{1}{3}\right)$$

✓ Same answer both methods

ex2



Set up integrals $\oint_C \vec{F} \cdot \hat{n} ds$

a) directly

b) using Green's theorem



a) $\int_{C_1} = 0$ because $y=0$ so $F=0$

$\int_{C_2} \vec{F} \cdot \hat{n} =$ needs parameters
what do we have?

$$x(t) = t$$

$$y(t) = 1 - t^2$$

$$x = x$$

$$y = 1 - x^2$$

that parametrized $-C_2$

$$\int_{C_2} \vec{F} \cdot \hat{n} \, ds = - \int_{-C_2} \vec{F} \cdot \hat{n} \, ds$$

$$\hat{n} \, ds = \langle y', -x' \rangle dt = \langle -2t, -1 \rangle \text{ in terms of } t$$

$$I = \int_{t=-1}^1 \langle (1-t^2)^2 + (1-t^2), 2t(1-t^2) \rangle \cdot \langle -2t, -1 \rangle dt$$

b) $I = \iint \operatorname{div} \vec{F} \, dA$

$$\begin{aligned} \operatorname{div} \vec{F} &= \frac{\partial}{\partial x} (x^2 + y) + \frac{\partial}{\partial y} 2xy \\ &= 2x \end{aligned}$$

Proving a theorem
- doing in recitation

Property if $\vec{F} = \langle M, N \rangle$ satisfies $M_y = N_x$ on $\mathbb{R}^2$ then $\vec{F}$ is conservative (aka its a gradient field) $\rightarrow$ f such that $\vec{F} = \nabla f$

Proof We suppose $\vec{F} = \langle M, N \rangle$ satisfies $M_y = N_x$

By Green's Theorem we get $\oint_C \vec{F} \cdot d\vec{r} = 0$
along closed path of gradient field

$$\begin{aligned} \text{(because } \oint_C \vec{F} \cdot d\vec{r} &= \iint_R N_x - M_y \, dA \\ &= \iint_R 0 \, dA = 0) \end{aligned}$$

• Thus if C_1, C_2 are continuous curves with same endpoints then $\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_2} \vec{F} \cdot d\vec{r}$

$$\begin{aligned} \text{(because } \int_{C_1} \vec{F} \cdot d\vec{r} - \int_{C_2} \vec{F} \cdot d\vec{r} \\ = \oint_{C_1 - C_2} \vec{F} \cdot d\vec{r} = 0) \end{aligned}$$

• We define a function f by $f(P) = \int_C \vec{F} \cdot d\vec{r}$ where C is going from $O \rightarrow P$

↳ (defined by preceding point)

$$\text{or rather } d\vec{r} = \vec{F}$$

Want to check

$$\frac{d}{dx} \int_C (x \cdot dy) = 1$$

Review in Lecture

4/13

$$\oint_C \vec{F} \cdot \vec{T} ds = \iint_A \text{curl } \vec{F} dA$$

$$d\vec{r} = \vec{T} ds = \langle x', y' \rangle$$

↖ tangent

$$\oint M dx + N dy = \iint_A (N_x - M_y) dA$$

Work form / tangential form

$$\oint_C \vec{F} \cdot \hat{n} ds = \iint_A \text{div } \vec{F} dA$$

$$\oint M dy - N dx = \iint_A (M_x + N_y) dA$$

normal form

Flux form



Green's Theorem

Lecture 25

Extension + applications of Green's Theorem

4/13

An example of Green's Theorem
- very important


Last recitation
- prof was wrong direction

Thm
$$\begin{cases} N_x = M_y \\ \text{curl } \vec{F} = 0 \end{cases} \rightarrow \vec{F} \text{ is conservative} \rightarrow \oint_C \vec{F} \cdot d\vec{r} = 0$$

(any simple closed C)
↓
So must be true in whole x, y plane

Proof = Green's Theorem

Example
$$\vec{F} = \frac{-y\hat{i} + x\hat{j}}{x^2 + y^2}$$



- perpendicular to normal of circle
- so tangential to circle

$$\begin{cases} \text{direction: tangential to circle} \\ \text{magnitude: } \frac{1}{r} = \frac{1}{a} \end{cases}$$

Calculate curl
(using chain rule)

Remember

$$\frac{\partial r}{\partial x} = \frac{x}{r} \quad \frac{x}{\sqrt{x^2+y^2}}$$

$$\frac{\partial r}{\partial y} = \frac{y}{r}$$

$$N_x = \frac{\partial}{\partial x} \left(\frac{x}{r^2} \right)^{cN}$$

$$= \frac{r^2 \cdot 1 - x \cdot 2r \cdot \left(\frac{x}{r} \right)}{r^4}$$

$$= \frac{r^2 - 2x^2}{r^4}$$

$$-M_y = -\frac{\partial}{\partial y} \left(\frac{y}{r^2} \right) = \frac{r^2 - 2y^2}{r^4}$$

Now add them

$$\frac{r^2 - 2x^2 - r^2 - 2y^2}{r^4}$$

$$= 2r^2 - 2r^2$$

$$= 0$$

Supposed to be a conservative field
- is it?

* practice


$$\begin{aligned}
 \oint_C \vec{F} \cdot d\vec{s} &= \underbrace{|\vec{F} \cdot \vec{t}|}_{\text{scalar product}} \cdot \text{length } C \\
 &= |\vec{F}| \\
 &= \frac{1}{a} 2\pi a \\
 &= 2\pi
 \end{aligned}$$

Paradox: Green's Theorem said 0
 But we got 2π
 and get 2π with $\sin + \cos$


Why???

So Green's Theorem not applicable
 - $\vec{F}$ is not defined at the origin
 - green's theorem does not apply

$\vec{F}$ is ^{magnetic} field of a wire coming out of page $\hat{z}$



unit DC current in $+\hat{z}/\odot$ dir



2π to go around

Problem - holes

Suppose $\text{curl } \vec{F} = 0$ in R , but R has "holes"

places where $\text{curl} \neq 0$

When no "holes" R is

"simply connected"

- can be shrunk to a pt w/o leaving the region



1 hole = "one connected"

$$\vec{F} = \frac{\langle -y, x \rangle}{r^2}$$

defined in whole $x-y$ plane
without $(0,0)$

$\text{curl} = 0$

"one connected" at $(0,0)$

Theorem



For any such C

$$\oint_C \vec{F} \cdot d\vec{r} = 2\pi$$

- can't go through pt b/c can't calculate line integral there

- but can be any path



← draw small second curve

- draw a line between them $\emptyset$

$$\underbrace{C_1 + D' + C_2' + D}_C$$

kinda overlaps when return on D
but think of it as a 2nd line
infinitely far away



So region is interior of C
- must walk in $(+)$ dir
with interior of left
 R is always on the left

$\vec{F}$ defined in R
 $\text{curl } \vec{F} = 0$ in R

Now Green's Theorem works!

$$\iint_R \text{curl } \vec{F} \cdot d\vec{r} = 0$$

this is
curl!

$$\int_C \vec{F} \cdot d\vec{r} = \oint_{C_1} + \int_{D'} + \oint_{C_2'} + \int_D$$

$\uparrow \quad \quad \quad \uparrow$
 Candles

So cut disappears

$$= \oint_{C_1} - \oint_{C_2} = 0$$


$$\text{So } \oint_{C_1} \vec{F} \cdot d\vec{r} = \oint_{C_2} \vec{F} \cdot d\vec{r} = \boxed{2\pi}$$

← that's what
we want

This is called "extended Green's Theorem"

(both work + flux forms)

If R has holes, bounded by closed curve



$$\int_{C_1 + C_2 + C_3} \vec{F} \cdot d\vec{r} = \iint_R \text{curl } \vec{F} \, dA$$


Proved i make cuts

Recitation

4/14

New stuff from last lecture: the holes

Remember $d\vec{r} = \vec{T} ds$
 $= \langle x', y' \rangle dt$
 $\hat{n} ds = \langle y', -x' \rangle dt$



ex1 $\vec{F} = \langle -y, x \rangle$

Compute $\int_C \vec{F} \cdot \hat{n} ds$ (flux) and $\int_C \vec{F} \cdot d\vec{r}$ (work)


- a) C is circle of radius a around 0
- b) C is line going from 0 to (a,b)

Hint Look at geometric properties of F

a) $\int_C \vec{F} \cdot \hat{n} ds$

will be 0

because flux is normal at every point



$\int_C \vec{F} \cdot d\vec{r}$

- do parametrization + calculation

= maximum

= product of the length

$\vec{F}$ is $\parallel$ to $d\vec{r} = \vec{T} ds$

$$\vec{F} \cdot d\vec{r} = |\vec{F}| \cdot |d\vec{r}| = |\vec{F}| ds$$

$$= \int_c \sqrt{y^2 + x^2} ds$$

$$= \int_c a ds$$


$$= a \int_c ds$$

= circumference of a circle

$$= a \times 2\pi a$$

$$= 2\pi a^2$$

b) $\int_c \vec{F} \cdot \hat{n} ds$



$$= \int_c -|\vec{F}| ds$$

Let's do by parametrization

$$x(t) = at \quad t \in [0, 1]$$

$$y(t) = bt$$

$$n ds = \langle y', -x' \rangle dt$$

$$= \langle b, -a \rangle dt$$

$$\int_c \vec{F} \cdot \vec{n} ds = \int_0^1 \langle -bt, at \rangle \cdot \langle b, -a \rangle dt$$


$$= \int_0^1 -(a^2 + b^2)t \, dt$$

$$-\left. \frac{(a^2 + b^2)t^2}{2} \right|_0^1$$

$$= -\frac{a^2 + b^2}{2}$$

$$\oint_C \vec{F} \cdot d\vec{r} = 0$$

because they are
⊥



Work = along curve

Flux = across curve


$$\oint_C \vec{F} \cdot d\vec{r}$$

$$\oint_C \vec{F} \cdot \hat{n} \, ds$$

think $F \rightarrow$ force field
velocity


ex2 $\vec{F} = \langle 2xy^2 - 2xy, 2x^2y - x^2 + y^2 \rangle$

Graduate
Student
Practicing
Teaching
:-)



Work done by F same along
both C_1 and C_2

We integrate over Region between C_1 and C_2



$$C = -C_1 - C_3 + C_2 + C_4$$

$$\oint \vec{F} d\vec{r} = -\int_{C_1} -\int_{C_3} + \int_{C_2} + \int_{C_4} \vec{F} d\vec{r}$$

$$= \oint_R \text{curl } \vec{F} dA$$

$$\vec{F} = \langle M, N \rangle$$

$$\text{curl } \vec{F} = N_x - M_y$$

$$= (4xy - 2x) - (4xy - 2x)$$

$$= 0$$

oh so one of those tricks
it will go to 0

$$\text{To show } \int_{C_3} \int_{C_4} \vec{F} d\vec{r} = 0$$

$$\int_{C_3} \vec{F} d\vec{r} =$$

$$\begin{pmatrix} x(t) = t \\ y(t) = 1 \end{pmatrix} \quad t \in [x_0, x_1]$$

$$d\vec{r} = \langle x(t)', y(t)' \rangle dt$$

$$= \langle 1, 0 \rangle dt$$

$$\int_{C_3} \langle 2t - 2t, 2t^2 - t^2 + 1 \rangle \cdot \langle 1, 0 \rangle$$


$$\int_{C_3} 0 dt$$

so 0

Similarly for C_4

$$\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_2} \vec{F} \cdot d\vec{r}$$

ex3 More General Case




To do this
make new curve C_3 that is completely
independent $\int_{C_1} \int_{C_3}$

ex4 Regions w/ holes

Suppose $\text{curl } \vec{F} = 0$ everywhere except at 2 points
 $(a, 0)$
 $(a+2, 0)$

$$\int_{C_P} \vec{F} \cdot d\vec{r} = 3 \quad \int_{C_Q} \vec{F} \cdot d\vec{r} = 1$$



Value of $\int_C \vec{F} \cdot d\vec{r}$ where C is circle of radius r around O ($r \neq a, a+2$)

if $r < a$ $\oint_C = 0$ b.
because $\oint_C = \iint_R \text{curl } \vec{F} = 0$

if $a < r < a+2$ $\oint_C = 3$

if $r > a+2$ $\oint_C = 4$

18.02 Problem Set 7 (DUE THURS. APR. 15 10:45 2-106)

Part I (15 pts.)

Lecture 23. Thurs. Apr. 8 Green's theorem (tangential form).

Read: 21.3 to middle p. 768 Work: 4D-1bc, 2, 3, 5

Lecture 24. Fri. Apr. 9 Flux; Green's Theorem in normal form.

Read: Notes V3, V4 Work: 4E-1ac, 2, 5 (without calculation); 4F-4, 5

Lecture 25. Tues. Apr. 13 Extensions and applications of Green's Theorem

Read: Notes V5, V6 (pp. 1,2) Work: 4G-1, 4, 5

Lecture 26. Thurs. Apr. 15 Triple integrals: rectangular and cylindrical coordinates

Read: Notes I.3, 20.5, 20.6

Part II (20 pts.)

Directions: Try each problem alone for 20 minutes. If you subsequently collaborate, solutions must be written up independently. It is illegal to consult problem sets from previous semesters. With each problem is the day it can be done.

Problem 1. (Thurs. 4 pts.: 1, 2, 1)

a) Let C be a simple, closed, smooth, positively oriented curve and let R be its interior. Show that

$$\frac{1}{3} \oint_C -y^3 dx + x^3 dy$$

is equal to the polar moment of inertia about the origin (cf. p. 725) of the plane region R , assuming its density $\delta = 1$.

b) Use part (a) to calculate the polar moment of inertia about the origin of the region inside the ellipse given parametrically by $x = a \cos t$, $y = b \sin t$, (a, b positive constants).

(Hint: Begin by sketching the ellipse. By symmetry it is enough to calculate the polar moment of inertia for the portion of the ellipse inside the first quadrant. This simplifies the boundary, and also permits the use of the formulas for $\int_0^{\pi/2} \sin^n t dt$ and $\int_0^{\pi/2} \cos^n t dt$ at the beginning of exercises 3B.)

c) Check your answer in part (b) by seeing what it gives when $b = a$, i.e., the ellipse becomes a circle of radius a . Then confirm this by calculating the polar moment of inertia of a circular disc of radius a , using a double integral in polar coordinates.

Problem 2. (Thurs. 3 pts.) Prove the other half of Green's theorem

$$\oint_C N(x, y) dy = \iint_R \frac{\partial N}{\partial x} dA$$

with the usual hypotheses, except assume the interior R of the curve is *horizontally simple*, i.e. is the region lying between the graphs of the smooth functions $x = g(y)$ on the left and $x = h(y)$ on the right, and the horizontal lines $y = a$ below and $y = b$ above.

if split through
day does
not seem
so long!

Problem 3. (Thurs. 3 pts.)Find the simple, closed, positively oriented curve C for which

$$\oint_C x^2(y+1) dx + 4x(1-y^2) dy$$

*A - need to find**- oh like my old area $1-x^2$*

has the maximum value. Indicate reasoning.

(Use Green's theorem; note that the integrand of $\iint_R$ takes on both positive and negative values.)*Want region for biggest integrand***Problem 4.** (Fri. 4 pts.: 3,1)Let $\mathbf{F} = 2xy(1-x^2)\mathbf{i} + y^2(3x^2-1)\mathbf{j}$ a) Show that the flux of $\mathbf{F}$ across any smooth curve $y = f(x)$ running from the line $x = -1$ to the line $x = 1$ has always the same value, i.e., the flux does not depend on the choice of curve.(Take two such curves C_1 and C_2 ; assume first that one lies entirely above the other and apply Green's Theorem in the normal form.)

b) Determine this constant value of the flux.

*not -n
b/c for $\oint$
So want from
-1 to 1*

$$4x^2 + y^2 = 0$$

$$4 - x^2 - y^2$$

$$4 - x^2 - 2y^2$$

Problem 5. (Fri. 2 pts.)A function $f(x, y)$ defined in a region S of the xy -plane is said to be **harmonic** in S if at every point of S it satisfies the two-dimensional form of **Laplace's equation**:

$$f_{xx} + f_{yy} = 0.$$

Prove that if the field $\mathbf{F}$ is the gradient of a harmonic function in S , the flux of $\mathbf{F}$ across any simple closed curve lying entirely inside S is zero.**Problem 6.** (Tues. 4 pts) Let

$$\mathbf{F} = \frac{-y}{x^2 + y^2} \mathbf{i} + \frac{x}{x^2 + y^2} \mathbf{j}.$$


Show that the flux of $\mathbf{F}$ across any simple closed curve C surrounding the origin is zero.

(Note that the field is not defined at the origin. Study and imitate the analogous argument in the notes involving the calculation of work done around a simple closed curve by a plane vector field whose curl vanishes, but which is not defined at the origin.)

Lecture 23 Green's Theorem Tangential Form4D-1b For the following fields F , curves C $\int_C \vec{F} \cdot d\vec{r}$

- directly as line integral
- green's theorem

$$F = x^2(\hat{i} + \hat{j}) = \langle x^2, x^2 \rangle$$


$$C = \text{rectangle}$$


So we want to do

$$\oint_C M dx + N dy$$

but have to do it for each section

$$\int_{C_1}^2 x^2 dx + \int_{C_2}^1 4 dy + \int_{C_3}^0 x^2 dx + \int_{C_4}^0 0 dy$$



$$\uparrow dy=0$$

$$y=0 \text{ \& does not matter}$$

$$x \text{ changing}$$

$$\cancel{x^2}$$

$$\uparrow dx=0$$

$$x=2$$

$$y \text{ changing}$$

$$\cancel{x^2}$$

$$\uparrow dy=0$$

$$x \text{ changing}$$

$$x^2$$

$$\uparrow x=0$$

$$x^2=0$$

and then it adds

(4)

Ok - this was last P-set right?

Now Green's Theorem - which is really simple to calculate

$$\iint_R N_x - M_y \, dA$$

$$N = x^2$$

$$N_x = 2x$$

$$M = x^2$$


$$M_y = 0$$

$$\iint_A 2x - 0 \, dA$$

⌈

now need to integrate over R

- again last P-set or one before that ⌈




$$\int_0^1 \int_0^2 2x \, dx \, dy$$

$$\left. \frac{2x^2}{2} \right|_0^2$$
$$\int_0^1 4 \, dy$$

④

Ok - so learned both before methods
- know how methods are different

c) $F = \langle xy, y^2 \rangle$ $C = y = x^2$ $0 \leq x \leq 1$



a) Directly

$$\int_C M dx + N dy$$

$$\int_{C_1} xy dx + y^2 dy + \int_{C_2} xy dx + y^2 dy$$

parametrize

$$\begin{aligned} y &= x^2 \\ x &= x \\ dx &= 1 \\ dy &= 2x \end{aligned}$$

$$\begin{aligned} x &= x \\ y &= x \\ dx &= 1 \\ dy &= 1 \end{aligned}$$

that's kinda
easy

$$\int x \cdot x^2 \cdot 1 + x^4 \cdot 2x + \int x \cdot x \cdot 1 + x^2 \cdot 1$$

now what?
or did I convert wrong? - or it would work when so I forgot definite integral

$$\int \vec{F} \cdot d\vec{r}$$

$$\int \vec{F} \cdot d\vec{r}$$

$$\int_0^1 \langle x^2 x, x^4 \rangle \cdot \langle 1, 2x \rangle + \int_0^1 \langle x x, x^2 \rangle \cdot \langle 1, 1 \rangle$$

$$\int_0^1 x^3 + 8x^5 + \int_0^1 x^2 + x^2$$

did not actually integrate

~~9~~ + ~~2~~

next

$$\frac{x^4}{4} + \frac{x^6}{3} \Big|_0^1$$

$$\frac{7}{12}$$

$$\left(-\frac{1}{12} \right)$$

$$\frac{2}{3} x^3 \Big|_0^1$$

$$-\frac{2}{3}$$

Don't forget to
actually integrate!

b) Green's Theorem

$$M = xy$$

$$N = y^2$$

$$\cancel{M_x = 1_y}$$

$$\cancel{N_y = 2y}$$

note backwards

$$N_x = 0$$

$$M_y = 1_x$$

$$\iint_R 0 - 1x \, dA$$

$$\int_{x^2}^x \int_0^1 -1x \, dy \, dx$$

$$-\frac{x^2}{2} \Big|_0^1$$

$$\int_{x^2}^x -\frac{1}{2} \, dy$$

$$-\frac{1}{2} y \Big|_{x^2}^x$$

$$-\frac{1}{2} x - \left[-\frac{1}{2} x^2 \right]$$

$$-\frac{x + x^2}{2}$$

$$\frac{x^2 - x}{2}$$

but need #

-switch integral x and y

$$\int_0^1 \int_{x^2}^x -x \, dy \, dx$$

$$-xy \Big|_{x^2}^x$$

$$-xx - [-xx^2]$$

$$-x^2 + x^3$$

$$\int_0^1 x^3 - x^2 \, dx$$

$$\frac{x^4}{4} - \frac{x^3}{3} \Big|_0^1$$

$$\frac{1}{4} - \frac{1}{3}$$

$$\frac{3}{12} - \frac{4}{12} = \left(-\frac{1}{12} \right) \quad \checkmark$$

2. Show that $\oint_C 4x^3 y \, dx + x^4 \, dy = 0$ for all closed C

Well isn't it a closed path - yeah by icon
But isn't the field not conservative, so it is not

Oh no - look what it is

Green's Theorem

$$M = 4x^3 y$$

$$N = x^4$$

$$N_x = 4x^3$$

$$M_y = 4x^3$$

$$\iint_R 4x^3 - 4x^3 = 0$$

I guess you just have to see that

3. Find the area inside the hypocycloid $x^{2/3} + y^{2/3} = 1$
by using Green's theorem
parametrize $x = \cos^3 \theta$
 $y = \sin^3 \theta$

Hint: $\vec{F} = \frac{1}{2} \langle y, x \rangle$

What's a hypocycloid?

WP

covered these before


$$x(\theta) = (R-r)\cos\theta + r\cos\left(\frac{R-r}{r}\theta\right) = M$$

$$y(\theta) = (R-r)\sin\theta - r\sin\left(\frac{R-r}{r}\theta\right) = N$$



ann

So how would I do this?



and why for area
isn't that normal double area
or do it for Flux

$$\iint_R (M_x + N_y) dA$$

still need to $\int$ over the area!

Use symmetric form for integrand
since parametrization of curve
does not favor x or y

∴
Symmetric
form

∴ What is this so called symmetric form?

$$\text{Area} = \frac{1}{2} \oint_C -y dx + x dy$$

$$\frac{1}{2} \int_0^{2\pi} 3 \sin^4 t \cos^2 t dt + 3 \sin^2 t \cos^4 t dt$$

∴
Don't get
their solution

Where in all world did they get that?
What were the original functions
they were using? The parameterizations?

$$\frac{3}{2} \int_0^{2\pi} \sin^2 t \cos^2 t dt$$

$$\begin{aligned} \text{Now using } \sin^2 t \cos^2 t &= \frac{1}{4} (\sin 2t)^2 \\ &= \frac{1}{4} \cdot \frac{1}{2} (1 - \cos 4t) \end{aligned}$$

$$\begin{aligned} \text{Now above} &= \frac{3}{8} \left(\frac{t}{2} - \frac{\sin 4t}{8} \right) \Big|_0^{2\pi} \\ &\quad \frac{3\pi}{8} \end{aligned}$$

Do not get whatsoever!

3

$$\text{Area} = \iint_R 1 \, dA = \oint \vec{F} \cdot d\vec{r}$$

if $|\vec{F}| = 1$
" $N_x - N_y$ "

again
Office
Hrs
Oliver

on
lecture
day 2
but marked
as day 1

need to find function
some easy fields prof gave us

$$\vec{F} = \langle 0, x \rangle \text{ or } \langle -y, 0 \rangle \text{ or } \langle -\frac{y}{2}, \frac{x}{2} \rangle$$

Use this trick to compute area

They gave parametrization

$$x = \cos^3 \theta \quad \theta \in [0, 2\pi]$$

$$y = \sin^3 \theta$$

$$\int_0^{2\pi} \langle 0, \cos^3 \theta \rangle \cdot \langle -3\sin \theta \cos^2 \theta, 3\cos \theta \sin^2 \theta \rangle \, d\theta$$

$$\int_0^{2\pi} 3 \cos^4 \theta \sin^2 \theta \, d\theta$$

-but how to $\int$ - no clear anti deriv

know

$$\int_0^{\pi/2} \cos^n \theta \, d\theta$$

cut integral into parts so get this

$$\int_0^{\pi/2} 1 - \cos^2 \theta$$

5. Show that the value $\oint_C xy^2 dx + (x^2y + 2x) dy$

around any square depends only on size
of square - not position

So is this another trick

$$M = xy^2$$

$$N = x^2y + 2x$$

$$M_y = x^2$$

$$N_x = y^2$$

$$N_x = 2xy + 2$$

$$M_y = 2xy$$

$$\iint_R x^2 - y^2$$

↑ get this right

$$\int_0^y \int_0^x x^2 - y^2 dx dy$$

← think did this wrong

$$\left. \frac{x^3}{3} - xy^2 \right|_0^x$$

$$\int_0^y \left(\frac{1}{3}x - y^2 \right) dy$$

$$\left. \frac{1}{3}xy - \frac{y^3}{3} \right|_0^y$$

$$\frac{y}{3} - \frac{y^3}{3}$$

$$\boxed{\frac{-y^3 + y}{3}}$$

← so what is y

Try again

$$\iint_R (2xy + 2) - \cancel{x^2} 2xy$$

$$\int_0^y \int_0^x 2xy + 2 - y^2 \, dy \, dx$$

$$\left. \frac{2x^2y}{2} + 2y - \frac{y^3}{3} \right|_0^x$$

$$x^3 + 2x - \frac{x^3}{3}$$

$$\int_0^y x^3 + 2x - \frac{x^3}{3} \, dx$$

$$\left. \frac{x^4}{4} + \frac{2x^2}{2} - \frac{x^4}{3 \cdot 4} \right|_0^y$$

$$\frac{y^4}{4} + y^2 - \frac{y^4}{12}$$

so how does that help

Note have

$$\iint_R 2 \, dA = 2(\text{area of } R)$$

Lecture 24 Flux, Green's Theorem in normal form

4E-1a Let $F = -y\hat{i} + x\hat{j} = \langle -y, x \rangle$

Recall what this is



tangent to circle all all times

thinks
rotation

And remember $\hat{n}$ is straight out
(normal to tangent vector on right)

So it will be $\perp$ at all points and thus $= 0$

✓ Flux

c) C is line from $(0,0)$ to $(1,0)$



$$\int_C \langle -y, x \rangle \cdot \langle 0, 0 \rangle = \int_C \langle -y, x \rangle \cdot \langle 0, 0 \rangle$$

~~$ds = \vec{T} dr = \langle x', y' \rangle$
 $\vec{T} \text{ tangent} = \langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y} \rangle$
Flux form~~

$$= \langle 0, 0 \rangle$$

$\hat{n} = -\hat{j}$ is just the normal vector to curve

$$F \cdot \hat{n} = x\hat{j} \cdot -\hat{j} = -x$$

$$\int_0^1 -x dx = \left(-\frac{1}{2} \right)$$


2. Let $\vec{F}$ be the constant vector field $\vec{i} + \vec{j}$
Where would you place a directed line segment C so flux is



- a) Maximal - in the same direction of the vector
(since dot product) aka northwest ☺
- b) Minimal - if opposite vector - southeast
- c) Zero - perpendicular to vector northeast
southwest
- d) -1 - if C has direction + magnitude of $-\vec{i}$ or $-\vec{j}$

$$\begin{aligned} (\vec{i} + \vec{j}) \cdot (-\vec{i}) &= -1 \\ (\vec{i} + \vec{j}) \cdot (-\vec{j}) &= -1 \end{aligned}$$

But why



but how is that $\ominus$?

Don't think of flux in terms of
Gaussian surface - just vector addition
Have to integrate, right?

d) What are values of min and max?

$$\frac{|F| \cdot |h| \cdot \cos \theta}{\sqrt{2}} = \frac{1 \cdot \cos 0}{\sqrt{2}}$$

$$\frac{|F| \cdot |h| \cdot \cos \theta}{\sqrt{2}} = \frac{1 \cdot \cos \pi}{\sqrt{2}}$$

5. without calculation: Let $\text{dir}(F) = \text{radially out}$
 $|F| = r^m$ m integer

a) Evaluate the flux of F across circle
 at origin radius a CCW



Always tangent to circle
 so 0?

No

$$a^m \cdot 2\pi a$$

$$2\pi a^{m+1}$$

← where is this from

- I thought it was $\perp$ to

OH $\parallel$ to $\hat{n}$

which means it does add

$\vec{r} \rightarrow \hat{n} \rightarrow F$ so does add since parallel

But you get the 2π from integrating?
 Not shown in solution

did in lecture w/ parametrization
 - good to see 2 forms of it

b) For what values of m will the flux be independent of a ?

$$a^m$$

$$a^1 = a \quad a^0 = 1$$

$$a^2 = a \cdot a \quad \text{always}$$

So when $m=0$

they say -1

but that is $\frac{1}{a}$ and thus

still depends on a


no

no in whole function

$$2\pi a^{m+1}$$

So m should be -1
for exponent $= 0$

ans
they
gave

4F-4 Verify Green's Theorem in normal form by calculating both sides and showing that they are = if $F = x^2 \hat{i} + xy \hat{j}$ and C 

Ok so normal form means for flux,

~~$$\int_C F \cdot \hat{n} = \langle x^2, xy \rangle \cdot \hat{n}$$~~

~~$$\int_{C_1} + \int_{C_2} + \int_{C_3} + \int_{C_4}$$~~

~~$$\int_0^1 \langle x^2, xy \rangle \cdot \langle \frac{1}{2}, -\text{something} \rangle$$~~

+ ~~$\int$~~

Note the other form

$$\oint -N dx + M dy$$

$$M = x^2$$

$$N = xy$$

$$C_1 \int_0^1 \underset{x=0}{-xy dx} + \underset{dy=0}{x^2 dy} = 0 \quad (\checkmark)$$

$$C_2 \int_0^1 \underset{dx=0}{-xy dx} + \underset{x=1}{x^2 dy} \quad \leftarrow \text{actually put it in}$$

$$\int_0^1 x^2 dy \rightarrow x^2 y \Big|_0^1 \rightarrow x^2 \rightarrow 1$$

$$C_3 \int_1^0 -xy dx + x^2 dy$$

$y=1 \quad dy=0$

$$\int_0^1 -x dx \quad \checkmark$$

$$-\frac{x^2}{2} \Big|_0^1$$

$+x \quad +\frac{1}{2}$ ~~x~~ should always have π

$$C_4 \int_1^0 -xy dx + x^2 dy$$

$dx=0 \quad x=0$

$$\int_0^1 x^2 dy$$

$$x^2 y \Big|_0^1$$

$$-x^2 \quad \emptyset$$

$\nearrow$ thought error again

$$S_2 \quad 0 + \cancel{x^2} + y - \cancel{x^2} \quad 1 + \frac{1}{2}$$

$(7) \quad \left(\frac{3}{2}\right)$


Now other way after screwing that up fairly bad

$$\iint_R M_x - (-N)_y$$

$$M = x^2 \rightarrow M_x = 2x$$

$$N = xy \rightarrow -N = -xy \rightarrow (-N)_y = -x$$

$$\iint_R 2x - x \, dA$$



$$\int_0^1 \int_0^1 3x \, dx \, dy$$

$$\left. \frac{3x^2}{2} \right|_0^1$$

$$\int_0^1 \frac{3}{2} \, dy$$

$$\left. \frac{3y}{2} \right|_0^1 = \left(\frac{3}{2} \right) \text{ much better}$$

5. Calc the $\text{div } \vec{F}$ for $\vec{F} = r^n (x\vec{i} + y\vec{j})$
 $\text{curl } \vec{F}$ $= \langle r^n x, r^n y \rangle$

Simplify differentiation $r_x = \frac{x}{r}$ $r_y = \frac{y}{r}$

$$M = \frac{x}{r} x = \frac{x^2}{r} \frac{y^2}{\sqrt{x^2+y^2}}$$

$$N = \frac{y}{r} y = \frac{y^2}{r} \frac{y^2}{\sqrt{x^2+y^2}}$$

$$\text{Well } r = \sqrt{x^2+y^2}$$

$$N_x = \frac{-x y^2}{(x^2+y^2)^{3/2}}$$

$$M_y = \frac{-x^2 y}{(x^2+y^2)^{3/2}}$$

$$\text{curl } \vec{F} = \frac{-xy^2}{(x^2+y^2)^{3/2}} - \frac{x^2y}{(x^2+y^2)^{3/2}}$$

$$\frac{-xy^2 + x^2y}{(x^2+y^2)^{3/2}}$$

$$\text{div } \vec{F}$$

$$M_x = \frac{x(x^2 + 2y^2)}{(x^2+y^2)^{3/2}}$$

$$(-N)_y = \frac{-y(y^2 + 2x^2)}{(x^2+y^2)^{3/2}}$$

$$\text{div } \vec{F} = \frac{x(x^2 + 2y^2)}{(x^2+y^2)^{3/2}} - \left(\frac{-y(y^2 + 2x^2)}{(x^2+y^2)^{3/2}} \right)$$

$$\frac{x^3 + 2xy^2 + y^3 + 2x^2y}{(x^2+y^2)^{3/2}}$$

Here they explain how they got the hint

$$r = (x^2 + y^2)^{1/2}$$

$$r_x = \frac{1}{2}(x^2 + y^2)^{-1/2} \cdot 2x = \frac{x}{r} \quad \text{coh that was pre-derived}$$

-not to make deriv easier

$$\text{So } M_x = r^n x$$

$$N = r^n y$$

So by chain rule

$$M_x = r^n + nr^{n-1} \cdot x \cdot \frac{x}{r} \\ = r^n + nr^{n-2} x^2$$

$$M_y = r^n + nr^{n-1} \cdot y \cdot \frac{y}{r} \\ = r^n + nr^{n-2} y^2$$

$$\begin{aligned} \text{div } F &= M_x + M_y && \in \text{I thought it was } -(-M)_y \\ &= 2r^n + nr^{n-2}(x^2 + y^2) \\ &= r^n(2+n) && \in \text{flux, area} \end{aligned}$$

$$N_x = nr^{n-1} \cdot \frac{x}{r} \cdot y$$

$$M_y = nr^{n-1} \cdot \frac{y}{r} \cdot x$$

$$\begin{aligned} \text{curl } \vec{F} &= N_x - M_y \\ &= nr^{n-1} \frac{xy}{r} - nr^{n-1} \frac{xy}{r} \\ &= 0 && \in \text{work, line} \end{aligned}$$

Lecture 25 Extension + applications of Green's Theorem

46-1

Using the criterion of this section - tell which are conservative or not

each problem
is so slow
but whole thing
is fast
- since staying on
fast

- so this is with holes
- but what is conservative

conservative = can use any line and get same answer

But how do I tell from looking at it?

a)

$$(y^2 + 2) dx + 2xy dy$$

~~that~~ O.K. right $\rightarrow \text{curl } \vec{F} = 0$ if conservative

$$M = y^2 + 2$$

$$N_x = 2y$$

$$N = 2xy$$

$$M_y = 2y$$

$$\int 2y - 2y = 0 \quad (\text{conservative}) \quad \checkmark$$

$$\vec{F} = \nabla f$$

b) $x(\cos y) dx + y(\cos x) dy$

$$M = x(\cos y)$$

$$N_x = -x y \sin x$$

$$N = y \cos x$$

$$M_y = -x \sin y$$

$$\int -x y \sin x - [-x y \sin y]$$

-y

$$\int -xy(\sin x + \sin y) \quad \text{not conservative}$$

$$-y \sin x - x \sin y$$

c) $\frac{x\uparrow + y\uparrow}{\sqrt{r^2 - 1}}$ this seems like it will work
looks familiar

$$r^2 = x^2 + y^2$$

$$M = \frac{x}{\sqrt{x^2 + y^2 - 1}} \cdot x(x^2 + y^2 - 1)^{-1/2}$$

$$N = \frac{y}{\sqrt{x^2 + y^2 - 1}} \cdot y(x^2 + y^2 - 1)^{-1/2}$$

$$M_x = \frac{-xy}{(x^2 + y^2 - 1)^{3/2}}$$

$$M_y = \frac{-xy}{(x^2 + y^2 - 1)^{3/2}}$$

these are kinda fun

So 0 so field conservative
Does not matter what path you take
but $\mathbb{R}^2$ is exterior of unit circle
- not simply connected $\leftarrow ?$

d) $\frac{x dx + y dy}{\sqrt{1 - r^2}}$

$$M = \frac{x}{\sqrt{1 - x^2 - y^2}}$$

$$N = \frac{y}{\sqrt{1 - x^2 - y^2}}$$

$$M_x = \frac{xy}{(1 - x^2 - y^2)^{3/2}}$$

$$M_y = \frac{yx}{(1 - x^2 - y^2)^{3/2}}$$

what is the difference

Same $\rightarrow$ conservative
interior of unit circle
is simply connected

e) $\sqrt{x} \mathbf{i} + \sqrt{y} \mathbf{j}$

$$M = \sqrt{x}$$

$$N = \sqrt{y}$$

$$N_x = 0$$

$$M_y = 0$$

conservative

and R is in the first quadrant

is simply connected

F is a gradient field

I don't get all of this how R must be
Simply connected. I know no holes
but what does in side of circle mean

4. Even though the field of example 1

$$\vec{F} = xy \hat{i} + x^2 \hat{j} \quad \text{not a gradient field}$$

Show that $\oint \vec{F} \cdot d\vec{r} \neq 0$

around every simple closed curve

So this is work/curl

$$M = xy$$

$$N = x^2$$

$$M_x = 2x$$

$$M_y = 1x$$

$$d_x = y$$

$$d_y = 0$$

or is it that $d\vec{r} = 2\pi r$

$$\text{so } \oint \langle xx, x^2 \rangle \cdot 2\pi r$$

$$\text{or } \oint xy \cdot y + x^2 \cdot 0$$

use the right form

$$\int xy^2$$

$$\boxed{\iint_R 2x - x \, dA}$$

$$= \underset{\uparrow}{x} (\text{area of } R)$$

x component of center of mass


Region symmetric w/ respect to y axis
COM on y-axis

$$\text{So } \overbrace{x=0} \quad \text{and} \quad \iint x \, dA = 0$$

5. Which of the following regions are simply connected?

- the ones w/o holes
- again, is there anything else?
- oh it can be shrunk to one point

Wolfram
alpha



a) Half plane above x axis

- does going off to ∞ matter
- if not simply connected ☒ yes simply connected


b) Plane minus line segment $(0,0) \rightarrow (0,1)$

No $\rightarrow$ hole ☒

a circle surrounding R is included
but not the interior
so like



c) Plane - $(+) x$ axis




d) Plane - entire x axis



No (✓)

e) Region $r > 0$, $0 \leq \theta < \theta_0$




yes (✓)

f) Region b/w 2 concentric circles



~~yes~~ No - if hole its a no
even if they ask for region around

g) Region b/w 2 branches of $xy = 1$
 $y = \frac{1}{x}$



No - not connected

yes for some reason - they
must have different picture
than I do

Part 2

#1. Let $\oint_C$ be a simple, closed, smooth, $\oplus$ oriented curve

$$\frac{1}{3} \oint_C -y^3 dx + x^3 dy$$

is = to the polar moment of inertia

$$I_z = \iint_R r^2 \delta(x, y) dA$$

$$\iint_R r^2 dA$$

so what

Show give = polar moment of inertia

$$I_z = \iint_R r^2 dA$$

$$\text{Now } \iint_R N_x - M_y dA = \iint_R r^2 dA = \frac{1}{3} \oint_C -y^3 dx + x^3 dy$$

$$\text{So } N_x - M_y = r^2$$

$$N_x - M_y = x^2 + y^2$$

so am
I messing
up what =
what

$$M = -y^3$$

$$N = x^3$$

$$N_x = 3x^2$$

$$M_y = -3y^2$$

$$\frac{1}{3} \iint_R \underset{\uparrow}{3x^2} - \underset{\uparrow}{-3y^2} dA$$

$$\text{Wait } x^2 - y^2$$

$$\iint_R x^2 + y^2 dA = \iint_R r^2 dA \quad \checkmark$$

b) Use part a to calculate the polar moment of inertia
 $x = a \cos t$ & given parametrization
 $y = b \sin t$
 $t \text{ constants}$

Hint: start by sketching the ellipse. By symmetry
 only need portion of ellipse of 1st quad

$$\int_0^{\pi/2} \sin^n t \, dt$$


$$\int_0^{\pi/2} \cos^n t \, dt$$

$$\frac{1}{3} \oint -y^3 dx + x^3 dy$$

↑ don't know original x and y
 but what to do because M, N
 leads to $Nx - My$ which is $\oint x^2 + y^2$

Also do we know what the region is?
 oh oh ellipse

$$\int_0^x \int_0^y x^2 + y^2$$



$$\int_0^{b \sin t} \int_0^{a \cos t}$$

$$a^2 \sin^2 t + b^2 \cos^2 t \, dt$$

$$\int_0^{a \sin t} (a^2 \sin^2 t + b^2 \cos^2 t) dt$$

$$t \left(\frac{a^2}{2} + \frac{b^2}{2} \right) - \frac{\sin t \cos t (a^2 - b^2)}{2} \Big|_0^{a \sin t}$$

Dependent limit - unsolvable

Unless they want to $\int_0^{\pi/2}$ but I was not doing polar

$$\frac{\pi (a^2 + b^2)}{4}$$

$$\int_0^{b \sin t} \frac{\pi (a^2 + b^2)}{4}$$

No I think we wanted a

$$t \left(\frac{a^2}{2} + \frac{b^2}{2} \right) - \frac{\sin t \cos t (a^2 - b^2)}{2} \Big|_0^a$$

$$\frac{a (a^2 + b^2)}{2} - \frac{(a^2 - b^2) \sin a \cos a}{2}$$

Does not look like it either

Back to above

$$\frac{\pi(a^2+b^2)}{4} \cdot b \sin t$$

rr

Does not seem right either

try

try again

$$= \frac{1}{2} \oint -y^3 dx + x^3 dy$$

$$= \frac{1}{2} \int_0^{\pi/2} -b^3 \sin^3 t (-a \sin t) dt + a^3 \cos^3 t (b \cos t) dt$$

$$= \frac{ab^3}{2} \int_0^{\pi/2} \sin^4 t dt + \frac{a^3b}{2} \int_0^{\pi/2} \cos^4 t dt$$

$$= \frac{ab^3}{2} \int_0^{\pi/2} \sin^4 t dt + \frac{a^3b}{2} \int_0^{\pi/2} \cos^4 t dt$$

$$= \frac{ab^3}{2} \left(\frac{3!!}{4!!} \right) + \frac{a^3b}{2} \left(\frac{3!!}{4!!} \cdot \frac{\pi}{2} \right)$$

the
rule

$$= \left(\frac{\pi}{16} (ab^3 + a^3b) \right) \cdot 4 \text{ for full ellipse}$$

$$= \frac{\pi}{4} (ab^3 + a^3b) \quad \checkmark$$

- c) Check your answer to part b, by seeing $b=a$
aka a circle of radius a
Then confirm polar moment of inertia


If $b=a$, then

$$\frac{\pi}{4} (a^4 + a^4)$$

$$\frac{2a^4\pi}{4} = \frac{a^4\pi}{2} \quad \checkmark$$

$$I = \int_0^{2\pi} \int_0^a r^2 \cdot r \cdot dr d\theta$$

$$\frac{a^4\pi}{2}$$



$$= \int_0^{2\pi} \int_0^a r^3 dr d\theta$$

$$= \int_0^{2\pi} \frac{1}{4} r^4 \Big|_0^a d\theta$$

$$= \frac{1}{4} \int_0^{2\pi} a^4 d\theta$$

$$= \frac{1}{4} \cdot a^4 \cdot 2\pi$$

$$= \frac{a^4\pi}{2}$$


✓ confirms

✓


2. Prove the other half of Green's Theorem

$$\oint_C N(x, y) dy = \iint_R \frac{\partial N}{\partial x} dA$$

w/ usual hypothesis
except assume interior R is horizontally simple



Was totally confused till
saw this picture bc we
did in recitation



Want to integrate over Region
between C_1 and C_3

$$C = -C_1 + C_3 + C_2 - C_4$$

$$\oint \vec{F} \cdot d\vec{r} = \int_{-C_1} \vec{F} \cdot d\vec{r} + \int_{C_3} \vec{F} \cdot d\vec{r} + \int_{C_2} \vec{F} \cdot d\vec{r} - \int_{C_4} \vec{F} \cdot d\vec{r}$$

$$= \iint_R \text{curl } \vec{F} dA$$

$$= \iint_R (N_y - M_x) dA$$

know that $\int_{C_1} \int_{C_3} \vec{F} \cdot d\vec{r} = 0$

Never learned
how to write
a proof -
or even what
one is

$$x = x \quad dx = 1$$

$$y = \text{fixed \# constant} \quad dy = 0$$

$$\int M dx + N dy$$

$$\int M dx$$

some function of $y = \text{constant}$ either a or b

$$So = 0$$

So it does not matter

C_1 and C_2 are in different directions
So are -

And are over a conservative field
so it does not matter which path you take

$$\int_{C_1} \vec{F} \cdot d\vec{r} = - \int_{C_2} \vec{F} \cdot d\vec{r}$$

is there more I have to do

show area of inside

count up the edges


3. Find the simple, closed, $\oplus$ oriented curve C for which $\oint$ has max value

$$\oint x^2(y+1) dx + 4x(1-y^2) dy$$

Use Green's Theorem

Note it has both $\oplus/\ominus$ values

office hrs
oliver



Want the biggest integrand

Don't want $\ominus$ because that will be $\ominus$ so want from -1 to 1

The above is the flux

max flux = max area

but then what are the constraints?

$$\begin{aligned} \oint F \cdot dr &= \oint_C x(x+1) dx + 4x(1-y^2) dy \\ &= \oint M dx + N dy \end{aligned}$$

Know the rules

$$= \iint_R (N_x - M_y) dA$$

$$\begin{aligned} N &= 4x - 4xy^2 \\ M &= x^2y + x^2 \end{aligned}$$

then can write this

$$= \iint_R (4 - 4y^2 - x^2) dA$$

and find it

don't get
how
try
again

So we know

$$\text{curl } \vec{F} = 4 - 4y^2 - x^2$$

$$4 - 4y^2 - x^2 = 0$$

$$\frac{1}{4}x^2 + y^2 = 1$$

$$\left[\frac{x^2}{4} + \frac{y^2}{1} = 1 \right]$$

$\therefore$ R is biggest inside ellipse ✓

$$\text{Max curl will} = \text{max work} = \oint \vec{F} \cdot d\vec{r}$$

4. $\vec{F} = 2xy(1-x^2)\vec{i} + y^2(3x^2-1)\vec{j}$
 $\langle 2xy - 2x^3y, 3x^2y^2 - y^2 \rangle$

a) Show that the flux of $\vec{F}$ across any smooth curve $y=f(x)$ running from $x=-1$ to $x=1$ has same value

So basically field conservative

$$M = 2xy - 2x^3y$$

$$N = 3x^2y^2 - y^2$$

$$M_x = 2y - 6x^2y$$

$$N_y = 6x^2y - 2y$$

$$\iint_R (2y - 6x^2y + 6x^2y - 2y)$$

See next page

0
So field is conservative

Take 2 such curves C_1 and C_2 assume one lies entirely above the other
 Green's Theorem in normal form
 aka flux form




So this is basically the other problem $\sim 90^\circ$ and similar analysis works (by symmetry) not exactly

1

b) Determine flux?

Isn't flux 0
by what I showed



$C_1 + C_3$ are 0
by symmetry

But we only want flux of one curve

But when I calculated it came out
as 0 ✓

Conservative if $M_y = N_x$
which for us
 $M_y = -N_x$

Oh so I did do it right. But what next?

Well you know

$$\int_{C_1} \vec{F} \cdot d\vec{r} = \int \vec{F} \cdot d\vec{r}$$

↑
only bc -

5. A function $f(x, y)$ defined in a region S of the xy plane is said to be harmonic in S if every point of S it satisfies the 2D form of Laplace's equation

$$f_{xx} + f_{yy} = 0$$

Prove that if $\vec{F}$ is the gradient of a harmonic function in S the flux of $\vec{F}$ across any simple closed curve lying entirely in S is 0

do not
really get
about what is
what for gradient
non gradient
fields

? Did we learn about "harmonic functions"
- or they are teaching here

Is it that flux is 0 if inside conservative field?

Lets test $f = x^2 + xy$

$$\begin{array}{ll} f_x = 2x + y & f_{xx} = 2 \\ f_y = x & f_{yy} = 0 \end{array} \quad \text{not harmonic}$$

$$f = x^2 + y^2$$

$$\begin{array}{ll} f_x = 2x & f_{xx} = 2 \\ f_y = 2y & f_{yy} = 2 \end{array} \quad \text{not harmonic}$$

$$f = x^2 - y^2$$

$$\begin{array}{ll} f_x = 2x & f_{xx} = 2 \\ f_y = -2y & f_{yy} = -2 \end{array} \quad \text{harmonic}$$

Now $\vec{F} = \nabla f$

F is the gradient of f

$$\vec{F} = \langle 2x, -2y \rangle$$

$$M = 2x$$

$$M_x = 2$$

$$N = -2y$$

$$N_y = -2$$

$$\iint 2 + -2 = 0$$

Oh - its the same thing

$$\vec{F} = \langle f_x, f_y \rangle$$

$$M_x = f_{xx}$$

$$N_y = f_{yy}$$

so its $\iint_R f_{xx} + f_{yy}$

And if it $= 0$ then no flux ✓
field conservative

↑ —————> ? Am I making the right conclusions?


6.
$$\mathbf{F} = \frac{-y}{x^2+y^2} \hat{i} + \frac{x}{x^2+y^2} \hat{j}$$

Show that flux of $\mathbf{F}$ across any simply closed C surrounding origin $= 0$

Need to
look closely
at all of
the assumptions

Note $\rightarrow$ field is not defined at origin
- Study notes

How does this differ from notes?



$$\oint_C = \oint_{C_1} + \underbrace{\oint_{O'}}_{\text{cancel}} + \underbrace{\oint_{O'}}_{\text{cancel}} + \oint_{C_2}$$

$$\oint_{C_1} - \oint_{C_2} = 0$$

$$\int_{C_1} \vec{F} \cdot d\vec{r} = \int_{C_2} \vec{F} \cdot d\vec{r} = 2\pi$$

So is $dr = 2\pi n$?

Oh field is the same as example

$$\mathbf{F} = \frac{\langle -y, x \rangle}{r^2}$$

$$= \iint_R \text{curl } \vec{F} d\vec{A}$$

$$\iint_R (N_y + M_x) dA$$

$$N = x(x^2 + y^2)^{-1}$$

$$M = -y(x^2 + y^2)^{-1}$$

$$N_y = -x(x^2 + y^2)^{-2} \cdot 2y$$

$$M_x = y(x^2 + y^2)^{-2} \cdot 2x$$

$$\text{flux} = \iint_R N_y + M_x$$

$$= \iint_R \left(\frac{-2xy}{(x^2 + y^2)^2} + \frac{2xy}{(x^2 + y^2)^2} \right) dA$$


$$= \iint_R 0 dA$$

$$= 0 \quad \checkmark$$

1) a) $\oint_C \frac{1}{3}(-y^3 dx + x^3 dy)$ (P2)C
 $= \iint_R (x^2 + y^2) dA$, by Green's theorem:
 polar mom. of inertia of R about origin $\frac{\partial}{\partial x}(\frac{x^3}{3}) = x^2$
 $\frac{\partial}{\partial y}(-\frac{y^3}{3}) = -y^2$

b) $x = a \cos t$ $dx = -a \sin t dt$
 $y = b \sin t$ $dy = b \cos t dt$

$\therefore \frac{1}{3} \oint_{C_1+C_2+C_3} -y^3 dx + x^3 dy = \frac{1}{3} \int_{C_1} -y^3 dx + x^3 dy$


 since integrand is 0 on C_2 ($x=0, dx=0$) and C_3 ($y=0, dy=0$)

$\int_{C_1} \frac{1}{3}(-y^3 dx + x^3 dy) = \frac{1}{3} \int_0^{2\pi} (ab^3 \sin^4 t + a^3 b \cos^4 t) dt$

$= (ab^3 + a^3 b) \cdot \frac{1}{3} \cdot \frac{3!!}{4!!} \cdot \frac{\pi}{2} \quad \frac{3!!}{4!!} = \frac{3 \cdot 1}{4 \cdot 2}$

$= ab(a^2 + b^2) \cdot \frac{\pi}{16}$ Ans: $\boxed{ab(a^2 + b^2) \cdot \frac{\pi}{4}}$


(multiply by 4 to include all 4 quadrants)

c)  Putting $b=a$ in part (b) gives polar m.o.i. $= \frac{2a^4 \pi}{4} (= \frac{\pi a^4}{2})$

Calculating directly:

$= \int_0^{2\pi} \int_0^a r^2 \cdot r dr d\theta$

$= 2\pi \cdot \frac{a^4}{4}$, which checks.

2)  $x=g(y)$ $x=h(y)$
 $y=b$ $dy=0$
 $y=a$ $dy=0$
 so $\int_{C_2}=0, \int_{C_4}=0$

$\therefore \oint N dy = \int_{C_1} N dy + \int_{C_3} N dy$
 $= \int_a^b N(g(y), y) dy + \int_a^b N(h(y), y) dy$ (*)

Double integral: integrate along horizontal lines:

$\iint_R \frac{\partial N}{\partial x} dx dy = \int_a^b \int_{g(y)}^{h(y)} \frac{\partial N}{\partial x} dx dy$

Inner: $N(x, y) \Big|_{x=g(y)}^{x=h(y)} = N(h, y) - N(g, y)$

Outer: $\int_a^b (N(h, y) - N(g, y)) dy$,

which agrees with (*).

$\left(\int_a^b N(g(y), y) dy = - \int_a^b N(h, y) dy \right)$

3) By Green's theorem, (R)C
 $\oint_C x^2(y+1) dx + 4x(1-y^2) dy$

$= \iint_R 4(1-y^2) - x^2 dA$

$= \iint_R 4 - x^2 - 4y^2 dA$

To maximize the integral, R should be the region where $4 - x^2 - 4y^2 \geq 0$

i.e., where $x^2 + 4y^2 \leq 4$

This is the region inside the ellipse C:

$x^2 + 4y^2 = 4$



So this C is the closed curve that maximizes the line integral.

4 a)  Use Green's theorem (normal form):

$$\text{div } \vec{F} = \frac{\partial}{\partial x}(2xy - 2x^2y) + \frac{\partial}{\partial y}(y^2(3x^2 - 1))$$

$$= 2y - 6x^2y + 6yx^2 - 2y = 0$$

By Green's theorem:

$$\oint_C Mdy - Ndx = \int_{C_1} + \int_{C_2} + \int_{C_3} + \int_{C_4} = 0$$


(or $\vec{F} \cdot \hat{n} ds$) ($C_2' = C_2$ oppositely oriented)

$C_1: x=1, dx=0, \int_{C_1} Mdy = \int_{-1}^1 2 \cdot 1 \cdot y(1 - (1)^2) dy = 0$

$C_4: x=-1, dx=0, \int_{C_4} Mdy = 0$

$$\therefore \int_{C_1} Mdy - Ndx = - \int_{C_2'} Mdy - Ndx = \int_{C_2} Mdy - Ndx$$

If C_1 and C_2 intersect, choose a C_3 above both of them.


 Then by the previous:

$$\int_{C_1} Mdy - Ndx = \int_{C_3} Mdy - Ndx = \int_{C_2} Mdy - Ndx$$

b) Use as the path  $y=0, dy=0$
 $x=x$

$$\int_{C_1} Mdy - Ndx = \int_{C_1} 0 dx = 0, \text{ since } N(x, 0) = 0.$$

5 Let the function $f(x, y)$ be harmonic in S , and suppose $\vec{F} = \vec{\nabla} f = \langle f_x, f_y \rangle$

 Flux of $\vec{F}$ across C

$$= \oint_C f_x dy - f_y dx$$

(by Green's theorem in normal form)

$$= \iint_R (f_x)_x + (f_y)_y dA$$

$$= \iint_R (f_{xx} + f_{yy}) dA$$

$$= 0, \text{ since } f \text{ is harmonic.}$$

6 $\text{div } \vec{F} = 0$ by calculation: (except at the origin, where $\vec{F}$ is undefined)

$$\frac{\partial}{\partial x} \left(\frac{-y}{x^2 + y^2} \right) = -y \cdot \frac{\partial}{\partial x} (x^2 + y^2)^{-1}$$


$$= -y \cdot \frac{-2x}{(x^2 + y^2)^2}$$

$$\frac{\partial}{\partial y} \left(\frac{x}{x^2 + y^2} \right) = x \cdot \frac{\partial}{\partial y} (x^2 + y^2)^{-1}$$

$$= x \cdot \frac{-2y}{(x^2 + y^2)^2}$$

These have opposite signs, so their sum is 0.

(Thus $\oint_C \vec{F} \cdot \hat{n} ds = \iint_R \text{div } \vec{F} dA = 0$ for any C not surrounding the origin, but we want a C which does surround the origin.)



Surround C with a large circle C_1 centered at the origin.

By the extended Green's theorem (normal form)

$$\oint_{C_1 + C'} \vec{F} \cdot \hat{n} ds = \int_{C_1} \vec{F} \cdot \hat{n} ds - \int_C \vec{F} \cdot \hat{n} ds$$

$$= \iint_R \text{div } \vec{F} dA = 0$$

$$\therefore \oint_C \vec{F} \cdot \hat{n} ds = \oint_{C_1} \vec{F} \cdot \hat{n} ds$$

However, the integral on the right is 0: for at each point of the circle, (say it has radius a)

$$\vec{F} = \frac{-y\hat{i} + x\hat{j}}{a^2}, \text{ which is tangent to the circle, } \therefore \perp \text{ to the normal, } \therefore \vec{F} \cdot \hat{n} = 0$$

$$\hat{n} = \frac{x\hat{i} + y\hat{j}}{a} \quad \text{(or note that } \vec{F} \cdot \hat{n} = 0 \text{ from these two formulas)}$$

[or you can calculate

$$\oint_{C_1} Mdy - Ndx \text{ using } \begin{cases} x = a \cos \theta \\ y = a \sin \theta \end{cases} \text{ and get 0}]$$

Lecture 26

Triple Integrals + Rect + Cylindrical coords

4/15

- Exam a week from today
- vector integral calculus
 - Green's Theorem
 - triple integrals


Really need to study →

Now moving into space (3D)


Hard → setting up
figuring out equation of shape
limits

Coordinate systems


- rectangular



- cylindrical



- spherical



ρ is like r = distance from origin

θ = ~~ϕ~~ from x-axis
(in xy plane)

ϕ = phi
angle from z axis
co-latitude

$\phi = 0$ north pole

$\phi = \pi/2$ equator

Look at the grid lines for a coord system

- but grid lines are surfaces
- level surfaces of the coordinates

rect $x = c_1$) vert planes
 $y = c_2$
 $z = c_3$ horiz plane

cyl $r = c_1$ cylinder of radius c_1 , axis: z -axis
 $\theta = c_2$ like a page in a book ↑ both sides ↖ top view
 $z = c_3$ horiz plane

sph $\rho = c_1$ sphere centered at origin
 $\theta = c_2$ same; like a page in a book = both sides
 $\phi = c_3$ half cone / nappe

Need to know how to convert between

- must memorize or be able to derive




can remember
the triangle

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \end{aligned} \quad \rightarrow \quad \begin{aligned} r &= \rho \sin \phi \\ z &= \rho \cos \phi \end{aligned}$$


$$dV = dx dy dz$$

a little rectangular box
infinitesimally small



θ is the longitude

$$\text{Volume} - dV = dz r dr d\theta$$




think of large melon

first cut slice b/w 2 book pages
then divide up into values of ρ
- if mixed rest



$$dV = \rho^2 \sin \varphi d\rho d\varphi d\theta$$

Cone



$$\begin{aligned} \text{spherical } \varphi &= \alpha \\ \text{cylindrical } \frac{z}{r} &= \frac{h}{a} & z &= \frac{h}{a} r \\ \text{rect coord } z &= \frac{h}{a} \sqrt{x^2 + y^2} \end{aligned}$$


Lecture 27

Spherical Coords, Gravitational Attraction

4/16

Doing Triple Integrals

- harder than we will do so encounter all aspects
- read the book w/ simple examples
- no time for simple examples



← aux thing to use when calculating


↪ why did prof draw R like this
wouldn't it be the shadow

D = volume

R = area

$$\iiint_D f(x, y, z) dV$$

$$= \iiint f dz$$



integrate from where enters ($z = g(x, y)$)
to where it leaves ($z = h(x, y)$)

Use all the vertical lines in the shadow of D

- which is where x, y are in R

$$= \iiint_{g(x,y)}^{h(x,y)} f dz \cdot dA$$

0, Volume element

- spherical $\rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$

1, Equations for $g(x,y)$ $h(x,y)$

2, Finding R

- most students this is mistifying the most
- intersection of two surfaces

3, Put in limits on 3 integral signs

- Use symmetry of the figure to simplify the limits

- some are trivial

$$\int_0^{\pi/2} \sin^n x \, dx \quad \int_0^{\pi/2} \cos^n x \, dx \quad \left| \begin{array}{l} \text{hard to } \int \text{ from} \\ \text{scratch} \end{array} \right.$$

- a lot of problems will have some symmetry



In each problem one of them is harder than the others

ex 1

Paraboloid $z = x^2 + y^2$
Plane $z = 2y$


Find volume enclosed between them

level curves




$$z = y^2$$

(cross section of both)



Find R



(x_1, y_1) in R

$\Leftrightarrow$ there is a z ,
such that (x, y, z)
lies in intersection of 2
surfaces

Find boundary of R

$\downarrow$ if and only if

$$\begin{cases} z = x^2 + y^2 \\ z = 2y \end{cases} \text{ have solutions}$$

Points that satisfy both equations
- non linear ILS problem

eliminate z

only where (x, y) satisfies
 $x^2 + y^2 = 2y$


single most
essential
step $\rightarrow$

Boundary of R

$$x^2 + y^2 - 2y \rightarrow x^2 + (y-1)^2$$

complete square

- circle



to integrate over this region $\rightarrow$ natural to use polar coords



$$r = 2 \sin \theta$$

$$\text{vol} = \iiint 1 \, dz \cdot r \, dr \, d\theta \quad \leftarrow dV \text{ in cylindrical coordinates}$$

$$\int_0^\pi \int_0^{2 \sin \theta} \int_{x^2+y^2}^{2y} 1 \, dz \, r \, dr \, d\theta$$

$\theta \quad r \quad z$

Don't do fancy set-ups
just answer the q

In general set up + eval problems different
to minimize mistakes

$$\int_0^{\pi} \int_0^{2\sin\theta} \int_{r^2}^{2r\sin\theta} dz \cdot r \, dr \, d\theta$$

Now inner, middle, outer integrals

Inner

$$zr \Big|_{r^2}^{2r\sin\theta}$$

$$r [2r\sin\theta - r^2]$$

Middle

$$\frac{2}{3} r^3 \sin\theta - \frac{r^4}{4} \Big|_0^{2\sin\theta}$$

$$\frac{16}{3} \sin^4\theta - \frac{16}{4} \sin^4\theta$$

$$\frac{16}{12} \sin^4\theta$$

Outer

Use symmetry




$$2 \int_0^{\pi/2} \frac{4}{3} \sin^4\theta \, d\theta$$

$$\left[\frac{8}{3} \right]_0^{\pi/2} = \frac{3 \cdot 1}{4 \cdot 2} \cdot \frac{\pi}{2} = \boxed{\frac{\pi}{2}}$$

π if even

This HW might be longer than others

Gravitational attraction in spherical coords



$$\vec{F} = \langle F_x, F_y, F_z \rangle$$

Calculate F_z - total z -comp
force on the mass at 0

$$|\vec{F}| = \frac{G \cdot \int \frac{dm}{r^2} \cos \phi}{\cos \phi}$$

$$F_z = \frac{G dm}{r^2} \cos \phi$$

$$\text{Total } F_z = \iiint_0 \frac{G \cos \phi}{r^2} \delta(x, y, z) dV$$


$\delta(x, y, z)$ usually constant though

for spherical coords - small miracle

$$\iiint_0 \frac{\cos \phi}{r^2} \cdot r^2 \sin \phi \, dr \, d\phi \, d\theta$$

Problem: find gravitational attraction of
solid sphere
radius = a
 $\delta = 1$
on a point circumference

Place point on origin



= eq of sphere in spherical coordinates?

ρ = length of rod

ρ, φ coordinates

Don't memorize - be able to derive

$$\rho = 2a \cos \varphi$$

$$\oint \iiint \cos \varphi \sin \varphi \, d\rho \, d\varphi \, d\theta$$

$$\oint_0^{2\pi} \int_0^{\pi/2} \int_0^{2a \cos \varphi} \cos \varphi \sin \varphi \, d\rho \, d\varphi \, d\theta$$

Practice for exam 3

Ex1. Let $f(x, y, z) = x + \sqrt{x^2 + y^2} + x^2 + y^2 + z^2$. Express this function in terms of

- a) cylindrical coordinates.
- b) spherical coordinates.

Ex2. Let S be the solid in first quadrant between planes $z = 1$ and plane $x + y + z = 2$.

- 1) Set up limits for computing the integral of a function $f(x, y, z)$ over S : $\int \int \int f dz dy dx$.
- 2) Suppose the density of S is given by a function $\delta(x, y, z)$. What is the function f to integrate if one want to compute:
 - a. The volume of S .
 - b. The Mass of S .
 - c. The moment of inertia around y axis.
 - d. The moment of inertia line defined by $x = 0, z = 1$.
 - e. The x coordinate $\bar{x}$ of the center of mass.
 - f. The gravitational attraction exerted by S on a unit mass placed at origin.

Ex3. Set up limits of integration for $\int \int \int_S f dV$, where S is


- a) Cylinder of radius a and height 2 (Cylindrical).
- b) Part of the unit sphere (around origin) in first octant (Cylindrical, Spherical)
- c) Part of unit sphere (around origin) above plane $z = 1/2$ (Rectangular, Cylindrical, Spherical)
- d) Part of unit sphere (around origin) above cone $z = \sqrt{x^2 + y^2}$ (Rectangular, Cylindrical, Spherical)
- e) Ball of radius a around $Q = (0, 0, a)$ (Cylindrical, Spherical).
- f) Part of that ball above plane $z = a$ (Cylindrical, Spherical).
- g) Intersection of unit sphere around origin and unit sphere around $(0, 0, 1)$ (Cylindrical).

Recitation Review for Test 3

4/21


Lecture Coordinate Systems

Cylindrical



$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta \\z &= z \\ \theta &\in [0, 2\pi]\end{aligned}$$

Spherical



$$\begin{aligned}x &= \rho \sin \phi \cos \theta \\y &= \rho \sin \phi \sin \theta \\z &= \rho \cos \phi \\ \rho &= \text{distance from center} \\ \theta &\in [0, 2\pi] \\ \phi &\in [0, \pi]\end{aligned}$$

$$dV = dx dy dz = r dr d\theta dz = \rho^2 \sin \phi d\rho d\theta d\phi$$

$$dM = \delta dV$$

Gravitational attraction of P_2 over P_1

$$\vec{F} = \frac{G m_1 m_2}{d^2} \frac{\vec{P_1 P_2}}{|\vec{P_1 P_2}|} = \frac{G m_1 m_2}{d^3} \vec{P_1 P_2}$$

ex 1 $f(x, y, z) = x + \sqrt{x^2 + y^2} + \underbrace{x^2 + y^2 + z^2}$

Questions
on sheet


a) Cylindrical coords

$$f(r, \theta, z) = r \cos \theta + r + r^2 + z^2$$

b) spherical coords

$$\begin{aligned} f(\rho, \theta, \phi) &= \rho \sin \phi \cos \theta + \sqrt{(\rho \sin \phi \cos \theta)^2 + (\rho \sin \phi \sin \theta)^2} \\ &\quad + (\rho \sin \phi \cos \theta)^2 + (\rho \sin \phi \sin \theta)^2 + (\rho \cos \phi)^2 \\ &= \rho \sin \phi \cos \theta + \rho \sin \phi + (\rho \sin \phi)^2 + (\rho \cos \phi)^2 \\ &= \rho \sin \phi \cos \theta + \rho \sin \phi + \rho^2 \end{aligned}$$

ex 2



want top triangle

a) set up limit over that volume

$$\int_x \int_y \int_z f \, dz \, dy \, dx$$

Draw shadow of region in x-y plane



project volume onto xy plane
(get rid of z)


eliminates z in $\begin{cases} x+y+z=2 \\ z=1 \end{cases}$

Shadow region bounded by $x+y+1=2$
 $x+y=1$



the more automated way to do it

$$\int_0^1 \int_0^{1-x} \int_1^{2-x-y} \dots dz \quad \leftarrow \text{sdn for } z$$



b) If you want Volume
function f you integrate = 1

$$\text{Mass} \rightarrow f = \delta(x, y, z)$$


c) Moment of inertia solid around y axis
- know formulas from lecture

$$MI = \iiint \text{distance}^2_{\text{from } y \text{ axis}} \delta \, dV$$

$$\text{distance of } P=(x, y, z) \text{ from } y\text{-axis} = \sqrt{x^2 + z^2}$$

$$f = (x^2 + z^2) \delta$$

d) distance of $P(x, y, z)$ from line $= \sqrt{x^2 + (z-1)^2}$



e) $\bar{x} = \frac{1}{M} \iiint_V \underbrace{x \delta}_{f} dV$

f) Gravitational attraction

$$\vec{F} = \frac{G m_1 m_2}{d^2} \frac{\vec{P}_1 \vec{P}_2}{|\vec{P}_1 \vec{P}_2|} = \frac{G m_1 m_2}{d^3} \vec{P}_1 \vec{P}_2$$

x component of gravitational attraction

$$\downarrow \text{x component of } \vec{F} = \vec{F} \cdot \vec{\uparrow} = \frac{G m_1 m_2}{d^3} \vec{P}_1 \vec{P}_2 \cdot \vec{\uparrow}$$

$$f = \frac{G m_1 m_2}{d^3} \vec{P}_1 \vec{P}_2 \cdot \vec{\uparrow}$$

$m_1 = 1 \Rightarrow$ unit mass

$m_2 =$ mass given by small cube $= dm = \sigma dV$


$$f = \frac{G \cdot 1 \cdot \sigma}{(x^2 + y^2 + z^2)^{3/2}} x$$

$$\vec{P}_1 \vec{P}_2 = \langle x, y, z \rangle$$

$$\vec{P}_1 \vec{P}_2 \cdot \vec{\uparrow} = x$$


Then integrate, if required

3.



$$\int_0^{2\pi} \int_0^a \int_0^2 f dV$$

always do
this order! (I thought it was other way)




Cylindrical coords
One $\frac{1}{8}$ of a circle "unit sphere"

cylindrical

$$\int_0^{\pi/2} \int_0^1 \int_0^{\sqrt{1-r^2}} r \, dz \, dr \, d\theta$$

$$\begin{aligned} x^2 + y^2 + z^2 &= 1 \\ r^2 + z^2 &= 1 \end{aligned}$$

↑
from the shadow



spherical

$$\int_0^{\pi/2} \int_0^{\pi/2} \int_0^1 r^2 \sin\theta \, dr \, d\theta \, d\phi$$

← better

c)



rectangular


1. find shadow region

2. eliminate z

$$\begin{cases} x^2 + y^2 + z^2 = 1 \\ z = \frac{1}{2} \end{cases}$$

$$\Rightarrow x^2 + y^2 = \frac{3}{4}$$

Shadow is disc of radius $\frac{\sqrt{3}}{2}$



$$\int_{-\frac{\sqrt{3}}{2}}^{\frac{\sqrt{3}}{2}} \int_{-\sqrt{\frac{3}{2}-x^2}}^{\sqrt{\frac{3}{2}-x^2}} \int_{-\frac{1}{2}}^{\sqrt{1-x^2-y^2}} dz$$

Each time you fix something

Cylindrical

- Very similar in that need to find shadow region

$$\int_0^{2\pi} \int_0^{\frac{\sqrt{3}}{2}} \int_{-\frac{1}{2}}^{\sqrt{1-r^2}} dz$$

Spherical

$$\int_0^{2\pi} \int_0^{\phi_{\max}} \int_0^1 \rho \, d\rho \, d\phi \, d\theta$$



$$\sin(\phi_{\max}) = \frac{\sqrt{3}}{2}$$

$$\phi_{\max} = \frac{\pi}{3}$$

$\phi_{\max} =$

$$\int_0^{2\pi} \int_0^{\pi/3} \int_{\uparrow}^1$$

$$z = \frac{1}{2}$$

$$\rho \cos \phi = \frac{1}{2}$$

$$\rho = \frac{1}{2 \cos \phi}$$

$$\int_0^{2\pi} \int_0^{\pi/3} \int_{\frac{1}{2 \cos \phi}}^1$$

Seems
easier than
I thought


Test Review

04/1

4/21



$$\oint \vec{F} \cdot \vec{n} \, ds = \iiint_R \text{div } \vec{F} \, dV$$




$$\oint_C = \oint_{C'} + \iiint_R \text{div } \vec{F} \, dV$$

||

0

if $\text{div} = 0$

parallel = tangent there



$$\int \vec{F} \cdot d\vec{r} = 0$$

$$\int_C \frac{|\vec{F}|}{|\vec{F}|} \cdot \vec{n} \, ds = \iint_C = \text{length} = \sqrt{2}$$

18.02 Exam 3 Thurs. Apr. 21, 2010 11:05-11:55

Directions:

1. There are 3 sheets, printed on both sides: eight problems in all.
2. Do all the work on these sheets; use the blank part below if truly necessary. Write down enough to show you are not guessing.
3. No books, notes, calculators, use of cell-phones, etc.
4. Please don't start until the signal is given; stop at the end when asked to; don't talk until your paper is handed in.
5. When the exam starts, read through the exam and start with what you are surest of.
6. Fill out the information below now.

Name Michael Plasmeier e-mail@mit.edu theplaz

Recitation teacher Oliver Rec. hour 12

$$\int_0^{\pi/2} \sin^n \theta d\theta = \int_0^{\pi/2} \cos^n \theta d\theta = \frac{(n-1)!!}{n!!} A, \quad \text{where } A = \begin{cases} \pi/2, & n \text{ even} \\ 1, & n \text{ odd} \end{cases} \quad \text{and } n!! = n(n-2)(n-4) \dots$$

Score 76
passing 60

pg.1 16
pg.2 11
pg.3 6
pg.4 8
pg.5 11
Total 52

Problem 1. (10 pts: 4; 3,3)

a) Write down in xy -coordinates the vector field F whose vector at (x,y) points towards the origin and has unit length.



$F = \text{dir} \cdot \text{length}$

$\frac{-x}{\sqrt{x^2+y^2}} \hat{i} - \frac{y}{\sqrt{x^2+y^2}} \hat{j}$

$\frac{-x\hat{i} - y\hat{j}}{\sqrt{x^2+y^2}}$

b) (No calculation required or expected): For the oriented path C consisting of the line segment from $(0,2)$ to $(0,1)$, followed by the quarter-circle (center $(0,0)$, radius 1) from $(0,1)$ to $(1,0)$:

(i) find the value of $\int_C F \cdot dr$ on the path C


(ii) find the flux of F across C .

3 $\therefore$ Only C_1 matters, because $C_2 \perp$ so it = 0

$\int_0^2 M dx + N dy$

$\int_0^2 \frac{-y}{\sqrt{0^2+y^2}} dy$

$\int_1^2 1 dy = (1)$



2

$\therefore$ Flux = $\int M dy - N dx$

Only C_2 matters because $\perp$

$\int_0^{\pi/2} F \cdot dr$

$\frac{1}{4} \cdot 2\pi r = \frac{\pi(1)}{2}$

$r=1$

stupid

Problem 2. (10) Use Green's theorem to evaluate $\oint_C (y^3 + 2y) dx + x(3y^2 - 1) dy$ over the positively oriented square C having vertices at the four points ± 1 on the x - and y -axes.

7

$M = y^3 + 2y$

$N = 3xy^2 - x$

$N_x = 3y^2 - 1$

$M_y = 3y^2 + 2$

$\iint_R (3y^2 - 1) - (3y^2 + 2) dA$

$\int_{-1}^1 \int_{-1-x}^{1-x} (6y^2 - 1) dy dx$

$y = -1-x$

$\left[\frac{6y^3}{3} - y \right]_{-1-x}^{1-x}$


$(1-x)^2(1-x)$

$(1^2 - 2x + x^2)(1-x)$

$1 - 2x + x^2 + x - 2x^2 + x^3 - x^3 + 3x^2 - 3x + 1$

$2(1-x)^3 - (1-x) - [2(-1-x)^3 - (-1-x)]$

$2(-x^3 + 3x^2 - 3x + 1) - 1 + x - 2(-1-x)^3 - 1 - x$



oh it was $N_x - M_y$

- and flux is $M_x + N_y$
- confused item

they just multiplied
- 3 = area

$-3 \cdot (\sqrt{2})^2$

2 know length of side
-6

Problem 3. (20 pts: 5,10,5)
where a and b are constants.

Let $F = ay(x-y) \mathbf{i} + (x^2 - bxy + 3y^2) \mathbf{j}$,

a) Prove that if F is conservative, then $a = 2$, $b = 4$.

$$N_x = M_y$$

$$M = 2xy - 2y^2$$

$$N = x^2 - 4xy + 3y^2$$

$$M_y = 2x - 4y$$

$$N_x = 2x - 4y$$

$$2x - 4y = 2x - 4y$$

$N_x = M_y$ so conservative

b) Let $a = 2$, $b = 4$.

Using a systematic method (show work), find a function $f(x,y)$ such that $F = \nabla f$.

integrate along path

stupid thought mistake



$$\int_{C_1} M dx + N dy + \int_{C_2} M dx + N dy$$

flipped
work $\int M dx + N dy$
flux $\int M dy - N dx$

$$\int_0^x \frac{x^2 - 4xy + 3y^2}{x=0} dx + \int_0^y \frac{2xy - 2y^2}{x^2 - 4xy + 3y^2} dy$$

$$\int_0^x 3y^2 dx + \int_0^y 2xy - 2y^2 dy$$

$$3xy^2 \Big|_0^x + \frac{2xy^2}{2} - \frac{2y^3}{3} \Big|_0^y$$

should have

but don't have
 x, y pts!
so ok

$X - 5$

$$x_1^2 y_1^2 - 2x_1 y_1^2 + y_1^3$$

c) Continue to use $a = 2$, $b = 4$, and using the earlier parts of this problem:

evaluate $\int_C F \cdot dr = 0$ along the path C given by: $x = \tan t$, $y = \sin 2t$, as t runs from 0 to $\pi/4$.

(This is easy, but you must show the work.)

Fundamental Theorem of calculus line integrals
if gradient field so any path is the same if gradient field
 $\int_C F \cdot dr = f(x,y) \Big|_P^2$

$$t=0 \rightarrow (0,0)$$

$$t=\pi/4 \rightarrow (1,1)$$

well gradient field so any path is the same if gradient field
could use path above
at $\pi/4$ $x = \tan \pi/4 = 1$
 $y = \sin 2\pi/4 = 0$

$$4(1)(0)^2 - \frac{2}{3}(0)^3 = 0$$

$$4x^2 - 2xy^2 + y^3 \Big|_{0,0}^{1,1} = 0$$

$\textcircled{0}$

6/20

ran out of time


Problem 4 (20 pts: 5, 10, 5, 0)

Verify Green's theorem in the tangential (work) form for the vector field $\mathbf{F} = -y^3 \mathbf{i} + x^3 \mathbf{j}$ and the closed curve C consisting of the two segments of the x - and y -axes connecting the origin with the points $(a, 0)$ and $(0, a)$, plus the quarter-circle given by $x = a \cos \theta$, $y = a \sin \theta$, $0 \leq \theta \leq \pi/2$, $a > 0$.

That is, showing work:

- (a) apply Green's theorem;
 (b) evaluate the line integral (over all of C);
 (c) evaluate the double integral;
 (d) spend the next twenty minutes trying to make them agree.

(In calculating the line integral you will need the formulas on the cover sheet.)



$$\oint_C M dy - N dx$$

flux form

$$dx = -a \sin \theta$$

$$dy = a \cos \theta$$

parametrizing
 correction w/ θ

curved not calculate

$$\int_0^{\pi/2} \int_0^a (-a^4 \sin^4 \theta - a^4 \cos^4 \theta) r dr d\theta$$

Green's not stated

$$a = r$$

why double integral?

$$-a^4 \left(\frac{r^2}{2} \right) (\sin^4 \theta - \cos^4 \theta) \Big|_0^a$$

Now need to do each leg
 + Green's theorem

$$-\frac{a^6}{2} (\sin^4 \theta + \cos^4 \theta)$$

$$\int M dy = \int_0^a -y^3 dy = -\frac{a^4}{4}$$

$$\int -N dx = \int_0^a x^3 dx = \frac{a^4}{4}$$

$$-\frac{a^6}{2} \int_0^{\pi/2} \sin^4 \theta + \cos^4 \theta d\theta$$

both incorrect

$$2 \frac{(4-1)!!}{4!!} \cdot \frac{\pi}{2}$$

$$\int N_x + M_y = \int 3x^2 - 3y^2$$

$$2 \frac{3(3-2)}{4(4-2)(4-4)} \cdot \frac{\pi}{2}$$

? ran out of time

separate sheet


$$-\frac{a^6}{2} \cdot \frac{4-6}{16-8} \cdot \frac{\pi}{2} = \frac{-a^6 3 \pi}{16}$$

Problem 5. (10) Let $f(x, y)$ be a function with continuous second derivatives that satisfies the equation

$$f_{xx} + f_{yy} = 0.$$

Show that the field $\mathbf{F} = \nabla f$ has the same flux across any two differentiable paths C_1 and C_2 which start at the same point P_1 , end at the same point P_2 , and don't intersect.

because it is a gradient field



$$\text{curl } \vec{F} = 0$$

separate sheet

$$N_x + M_y = 0$$

$$\iint_R 0 = 0$$

$$N_x = f_{yx}$$

$$M_y = f_{xy}$$

$$\left. \begin{array}{l} N_x = f_{yx} \\ M_y = f_{xy} \end{array} \right\} \text{ if } f_{yx} + f_{xy} = 0 \text{ then } f_{xx} + f_{yy} = 0$$

create closed surface

Use Green's Theorem


$$\iint M_x + N_y$$

Problem 6. (10 pts.: 8,2)

Let D be the domain of finite volume in the first octant cut off by the plane $x + y + z = 1$.

a) Set up an iterated triple integral in rectangular coordinates $\iiint f(x, y, z) dz dy dx$ for the total mass of D , if the density function is given by $\delta = z$. (Give the integrand and limits.)

b) Then take only the first step in evaluating it, i.e., reduce it to a double iterated integral in x and y .



Oh that was wrong

it was all 1

$$\text{Mass } \iiint \sigma dv = \int_0^{1/3} \int_0^{1/3(1-z)} \int_0^{1/3(1-z-y)} z dx dy dz$$

oppo

set x constant


$$y = 1 - z$$

$$\int_0^{1/3} \int_0^{1/3(1-z)} 2 dx dy dz$$

$$\int_0^1 \int_0^{1-x} \int_0^{1-x-y} z dz dy dx$$

$$\int_0^{1/3} \int_0^{1/3(1-z)} z \left(\frac{1}{3} - \frac{1}{3}z - \frac{1}{3}y \right) dy dz$$

Problem 7. (10) Let V be the finite domain in xyz -space lying between the paraboloid $z = x^2 + y^2$ and the plane $z = 9$. Taking density $\delta = 1$, set up an iterated triple integral in cylindrical coordinates (provide limits and integrand, but do not evaluate) giving its moment of inertia about the z -axis.



mass

$$\int_0^{2\pi} \int_0^3 \int_{r^2}^9$$

just base




$$x^2 + y^2 = r^2$$

$$dz \, r \, dr \, d\theta$$


did wrong order

$$\int_0^9 \int_{r^2}^9$$

if want distance take $\sqrt{r^2}$
 not $\sqrt{r}$



drew wrong



Problem 8. (10) Let D be the "filled ice-cream-cone" in 3-space whose top surface is formed by a portion of the sphere of radius a centered at the origin, and whose bottom surface is a portion of the circular cone having as axis the z -axis, and whose vertex angle is 60° . Set up a triple iterated integral in spherical coordinates which gives the volume of D (provide limits and integrand, but do not evaluate).

$$\iiint \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$$



$$\int_0^{2\pi} \int_0^{60^\circ} \int_0^a$$

$$\rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$$

can just say



$$\sin 60^\circ = \frac{a}{r}$$

$$\frac{a}{\frac{\sqrt{2}a}{2}} = \frac{2a}{\sqrt{2}a} = \frac{2\sqrt{2}}{2}$$



$$\tan 60^\circ = \frac{a}{r}$$

$$\tan 60^\circ = \sqrt{3}$$

4) a) $\oint -y^3 dx + x^3 dy = \iint_A 3x^2 + 3y^2 dA$

think I screwed it up again

~~Flux $\oint M dy - N dx$ ← what I wrote~~
~~work $\oint M dx + N dy$ ← what it asked for~~

$\oint F_T dx + F_T dy$

$\oint M dx + N dy$

$\vec{F}_x$

tangent

normal

work	$\oint M dx + N dy$	$\stackrel{GT}{=} \iint (N_x - M_y) dA$
flux	$\oint M dy - N dx$	$\stackrel{or}{=} \iint (M_x + N_y) dA$

Both
Always
work

grad
field or
not

the fundamental problem I had on the test was screwing up these $\oint$

$\oint$


$\oint M dx + N dy = \iint (N_x - M_y) dA$
 $M = F_T$ one equation

b)

Now break into parts

$C_1 = y = 0 \quad dy = 0 \quad S = 0$

$C_2 = x = 0 \quad dx = 0 \quad S = 0$



→

$$C_2 = \begin{aligned} x &= a \cos \theta & dx &= -a \sin \theta \\ y &= a \sin \theta & dy &= a \cos \theta \end{aligned} \quad \text{✓ had this}$$

$$\int_C = \int_0^{\pi/2} -a^3 \sin^3 \theta (-a \sin \theta) + a^3 \cos^3 \theta (a \cos \theta) d\theta$$

↗
not double
integral
just along
line

$$a^4 \int_0^{\pi/2} (\sin^4 \theta + \cos^4 \theta) d\theta$$

$$a^4 \left(\frac{3 \cdot 1}{4 \cdot 2} \frac{\pi}{2} + \frac{3 \cdot 1}{4 \cdot 2} \frac{\pi}{2} \right)$$

$$\frac{3\pi}{8} a^4$$

what did I screw up here

remember $!!_1 =$ every other one

$$n=4$$

$$n-1=3$$

$$(n-1)!!_1 = (4-1) \cdot (4-1-2) = 3 \cdot 1$$

$$n!!_1 = 4 \cdot 2$$

this was close but I
screwed it up

No it was actually right
except I did not do 16-8
really stupid!

c) Here double integral

$$= \iint_R 3x^2 + 3y^2 dA$$

but w/ these #

$$\int_0^{\pi/2} \int_0^a 3r^2 \cdot r dr d\theta$$

- use polar

$$= \frac{3r^4}{4} \Big|_0^a \cdot \frac{\pi}{2}$$

$$\frac{3\pi}{8} a^4$$

$$5. \nabla f = \langle f_x, f_y \rangle$$

did not know these
rules

flux of $\vec{\nabla} f$ over C

$$= \int_C M dy - N dx$$

$$\uparrow M = f_x \quad \uparrow N = f_y$$

$$\oint M dy - N dx$$

or div

$$\iint_R f_{xx} + f_{yy} dA$$

$$\int_{C_1'} + C_2 = -\int_{C_1} + \int_{C_2} = 0$$

$$\int_{C_1} = \int_{C_2}$$

yeah that it is a - proof

1) a) $\vec{F} = -\frac{\langle x, y \rangle}{\sqrt{x^2 + y^2}}$

b) $\int_C \vec{F} \cdot d\vec{r} = 1 + 0 = 1$
on C_1 , on C_2

flux over $C = 0 + \frac{\pi}{2} \cdot 1 = \frac{\pi}{2}$
on C_1 on C_2

2) $\oint_C (y^3 + 2y) dx + x(3y^2 - 1) dy$
 $= \iint_R (N_x - M_y) dA$
 $= \iint_R (3y^2 - 1) - (3y^2 + 2) dA$
 $= \iint_R -3 dA = -3 \cdot (\sqrt{2})^2 = -6$

3) $\frac{\partial M}{\partial y} = a(x - 2y)$ $\frac{\partial N}{\partial x} = 2x - by$

a) $a(x - 2y) = 2x - by \Rightarrow a = 2$
 $b = 2a = 4$

b) Method 1:

$f(x, y) = \int_{C_1 + C_2} 2y(x - y) dx + (x^2 - 4xy + 3y^2) dy$

$\int_{C_1} = 0 + 0$ (since $y=0, dy=0$)

$\int_{C_2} = 0 + \int_0^1 (x^2 - 4x, y + 3y^2) dy$
($dx=0$)

$= x_1^2 y_1 - 2x_1 y_1^2 + y_1^3$

Method 2:

$f_x = 2y(x - y)$ $f_y = x^2 - 4xy + g(y)$

$f = yx^2 - 2y^2x + g(y) = x^2 - 4xy + 3y^2$

$\therefore g' = 3y^2, g = y^3$

$f = yx^2 - 2y^2x + y^3$ (+c unnecessary)

c) $t=0 \Rightarrow (x, y) = (0, 0)$ use
 $t=\pi/4 \Rightarrow (x, y) = (1, 1)$ F.T. Calc for
 line integrals
 $yx^2 - 2y^2x + y^3 \Big|_{(0,0)}^{(1,1)} = 0 - 0 = 0$

4) $\oint_C -y^3 dx + x^3 dy = \iint_R 3x^2 + 3y^2 dA$

b) On $C_1: y=0, dy=0 \int_{C_1} = 0$

$C_3: x=0, dx=0 \int_{C_3} = 0$

$C_2: x = a \cos \theta, y = a \sin \theta$
 $dx = -a \sin \theta, dy = a \cos \theta$

$\int_{C_2} = \int_0^{\pi/2} -a^3 \sin^3 \theta (-a \sin \theta) + a^3 \cos^3 \theta (a \cos \theta) d\theta$

$= a^4 \int_0^{\pi/2} (\sin^4 \theta + \cos^4 \theta) d\theta =$

$= a^4 \left(\frac{3 \cdot 1}{4 \cdot 2} \cdot \frac{\pi}{2} + \frac{3 \cdot 1}{4 \cdot 2} \cdot \frac{\pi}{2} \right) = \frac{3\pi a^4}{8}$

c) $\iint_R 3x^2 + 3y^2 dA = \int_0^{\pi/2} \int_0^a 3r^2 \cdot r dr d\theta$
 (use polar coords) $= \frac{3r^4}{4} \Big|_0^a \cdot \frac{\pi}{2}$
 $= \frac{3\pi a^4}{8}$


5) $\vec{\nabla} f = \langle f_x, f_y \rangle$

Flux of $\vec{\nabla} f$ over $C = \int_C M dy - N dx$
 (any C) $M = f_x, N = f_y$

$\oint_{C_1 + C_2} M dy - N dx = \iint_R (f_{xx} + f_{yy}) dA = 0$
 $\text{div}(\vec{\nabla} f)$

$\int_{C_1 + C_2} = -\int_{C_1} + \int_{C_2} = 0, \therefore \int_{C_1} = \int_{C_2}$

6)
 $\int_0^1 \int_0^{1-x} \int_0^{1-x-y} z dz dy dx$
 Inner: $\frac{1}{2} z^2 \Big|_0^{1-x-y}$
 $\therefore \int_0^1 \int_0^{1-x} \frac{1}{2} (1-x-y)^2 dy dx$




(cross-section)

$\int_0^{2\pi} \int_0^3 \int_{r^2}^9 r^2 \cdot dz \cdot r dr d\theta$

moment of inertia about z-axis

dA in polar coords



eqn cone: $\varphi = \pi/6$

eqn cap: $\rho = a$

$$\text{Vol} = \int_0^{2\pi} \int_0^{\pi/6} \int_0^a \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$$

integrand = dV in spherical coords


Triple Integrals

From Book Chap 20.5

4/20

Same pattern as double integral (20.2)

Review } - which was over a region



Add the blocks

$$\iint_R f(x, y) dA$$

Must be bounded by a rectangle
- Not off to ∞

Add up the area of each rectangle

The function is the height
- volume = height $\cdot$ area of base

End Review L


$f(x, y, z)$ defined in 3D region R

Divide R into many small rectangular boxes along coords
- add volumes

Take limit as diagonal of boxes $\rightarrow 0$

$$\Delta V = \Delta x \Delta y \Delta z$$

$$dx dy dz$$



function must be continuous, etc, etc

Region R is thought of as a solid
body $\delta = \delta(x, y, z)$

$$M = \iiint_R \underbrace{\delta dV}_{\text{element mass}}$$

Moments

Moments of inertia

$$M_{yz} = \iiint x \delta dV$$

$$I_x = \iiint_R (y^2 + z^2) \delta dV$$

$$M_{xz} = \iiint y \delta dV$$

$$I_y = \iiint_R (x^2 + z^2) \delta dV$$

$$M_{xy} = \iiint z \delta dV$$

$$I_z = \iiint_R (x^2 + y^2) \delta dV$$

$$\bar{x} = \frac{M_{yz}}{M} \quad \bar{y} = \frac{M_{xz}}{M} \quad \bar{z} = \frac{M_{xy}}{M}$$

center of mass if δ constant

Evaluate w/ iteration


$$a \leq x \leq b$$

$$y_1(x) \leq y \leq y_2(x)$$

$$z_1(x, y) \leq z \leq z_2(x, y)$$

$$\int_a^b \int_{y_1(x)}^{y_2(x)} \int_{z_1(x, y)}^{z_2(x, y)} f(x, y, z) \, dz \, dy \, dx$$

Can choose any order you want
however



Lecture 28

Vector Fields in 3-space, flux surface S

4/23

Integration shortcut

$$\int_{a_2}^{b_2} \int_{a_1}^{b_1} f(x) g(y) dy dx$$

product

each function uses 1 variable

limits are constants

- not functions

- all of them

Can do 2 integrals separately

$$= \int_{a_1}^{b_1} f(x) dx \cdot \int_{a_2}^{b_2} g(y) dy$$

Same for triple integrals $\iiint$

Now extend vector calculus to space

- already did tripple integrals
- Green's theorem
- flux
- work

Need vector fields in 3-space


$$\vec{F} = M \hat{i} + N \hat{j} + P \hat{k} \quad (\text{or } P, Q, R)$$

are each functions of 3 variables

$$F = M(x, y, z) \hat{i} + N(x, y, z) \hat{j} + P(x, y, z) \hat{k}$$

Examples

gravitational fields
electrostatics
electromagnetic
velocity
fluid flow




$$\vec{F} = \frac{\langle x, y, z \rangle}{\rho} \cdot \frac{k}{\rho^2} = \frac{k \langle x, y, z \rangle}{(x^2 + y^2 + z^2)^{3/2}}$$

direction magnitude


in 3 space flux + work are 2
totally different concepts
each needs like 3-4 lectures

flux $\rightarrow$ flow

2D



3D flux across surface



Draw a little piece of territory on surface



If tangent = no flux
so only interested in component $\perp$
to surface

But what direction is positive?

- you must specify
- not automatic for 3D

Count up all the little surface areas

Flux through a little piece
 $\vec{F} \cdot \hat{n}$

(component of vector in the direction)
you are interested in) dot product

$$(\vec{F} \cdot \hat{n}) dS$$

(surface, not area
called - but same thing)

$$= \vec{F} \cdot d\vec{S}$$

$$\uparrow d\vec{S} = \hat{n} dS$$

Flux through Surface

$$= \iint_S \vec{F} \cdot d\vec{S}$$


surface integral for flux
of $\vec{F}$

The General formulas are useless
Need examples

$$\vec{F} = y\vec{j}$$

S: right hemisphere
of radius a
center at origin

Draw useful 2D pictures
Not complex 3D pictures



Must calculate
not intuition

(not direction of normal vector)

We need $\vec{F}, \hat{n}, dS$

$$\vec{F} = y\vec{j} \quad \text{known}$$

$$\hat{n} = \langle x, y, z \rangle$$

← same for all spheres
picked radially out (easier)

$$dS =$$

← "is a formula for
but its stupid" - Mattuck

Remember $dV = \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\theta$
Very small thickness $d\rho$

$$dS = a^2 \sin \varphi \, d\varphi \, d\theta$$

2 coord systems are fighting

$$\vec{F} \cdot \hat{n} = \frac{y^2}{a}$$

Convert y to spherical coordinates



$$y = r \sin \theta$$

$$r = a \sin \varphi$$

$$\vec{F} \cdot \hat{n} = \frac{y^2}{a} = \frac{(a \sin \varphi \sin \theta)^2}{a}$$

$$\text{flux} = \iint (\vec{F} \cdot \hat{n}) dS$$

$$= \int \int \underbrace{a \sin^2 \varphi \sin^2 \theta}_{\vec{F} \cdot \hat{n}} \underbrace{a^2 \sin \varphi d\varphi d\theta}_{dS}$$

$$= a^3 \int_0^{\pi} \int_0^{\pi} \sin^3 \varphi \sin^2 \theta d\varphi d\theta$$

now we can use our shortcut
but we could make it even easier
if have $\frac{1}{4}$ hemisphere (in 1st octant)

$$= 4a^3 \int_0^{\pi/2} \int_0^{\pi/2} \sin^3 \varphi \sin^2 \theta d\varphi d\theta$$

$$\text{Flux} = 4a^3 \int_0^{\pi/2} \sin^3 \varphi \, d\varphi \cdot \int_0^{\pi/2} \sin^2 \theta \, d\theta$$

$$= 4a^3 \cdot \frac{2}{3} \cdot \frac{1}{2} \frac{\pi}{2}$$

$$= \frac{2}{3} \pi a^3$$

$$= \underset{\substack{\uparrow \\ \text{area} \\ \text{hemisphere}}}{2\pi a^2} \cdot \underset{\substack{\uparrow \\ \text{velocity}}}{\frac{a}{3}}$$


Recitation

4/26

New material

Surface Integrals

$$\text{Def } \iint_S f(P) ds = \lim \sum f(P_i) \Delta s_i$$



* Flux of 3D field across S


$$\text{Flux} = \iint_S \vec{F} \cdot d\vec{s}$$
$$d\vec{s} = \hat{n} ds$$

$\hat{n}$ = unit vector
⊥ to surface
 ds = area


Today no parametrization

Field in
3D


→ Ex | Give flux of field $\vec{F}(x, y, z)$
which is velocity field of rigid
rotation of angular velocity ω
a) axis = y axis around
b) around a line through origin
direction $\vec{u}$



What is magnitude of field? $|\vec{F}(x, y, z)|$




$$\begin{aligned} \text{velocity} &= \text{radius} \cdot \omega \\ &= \text{distance to } y \text{ axis} \cdot \omega \end{aligned}$$



$$\text{distance} = \sqrt{x^2 + z^2}$$

$$|\vec{F}| = \omega |\vec{PQ}| = \omega \sqrt{x^2 + z^2}$$



$$\begin{aligned} \vec{F} &\perp |\vec{PQ}| \\ \vec{PQ} &= \langle x, 0, z \rangle \end{aligned}$$

$$\begin{aligned} \vec{F} &= \omega \langle z, 0, -x \rangle \\ &\quad \uparrow \text{to get right magnitude, adjust} \end{aligned}$$

Check sign is right

$$|\vec{F}| = \omega |\vec{PQ}|$$

$$\vec{F} \perp \vec{u}$$

$$\vec{F} \perp \vec{PQ}$$



↑ take cross product to
get something in right direction

$$\vec{F} = \vec{u} \times \vec{OP}$$

↑ what should coefficient be?
just w
for amplitude


$$|\vec{u} \times \vec{OP}| = |\vec{u}| \times |\vec{QP}| = |\vec{QP}|$$

In fact $\vec{F} = w \vec{u} \times \vec{OP}$
because $\vec{OP} = \vec{OQ} + \vec{QP}$
and $\vec{u} \times \vec{OQ} = \langle 0, 0, 0 \rangle$

Surface Integrals w/o Parametrization

ex2 What is the volume of flux $\phi = \iint_S \vec{F} \cdot d\vec{s}$

$S =$ horizontal rectangle w/ coord
 $(0,0,0) (0,1,0) (2,0,0) (2,1,0)$




must define:
oriented upward

a) $\vec{F} = xy\hat{i} + z\hat{j}$
b) $F(x,y,z) = \hat{i}$
c) $F(x,y,z) = xy\hat{i}$

Use some geometric property to simplify

Perpendicular to surface is vertical

$$d\vec{S} = \hat{k} dS$$



$$a) \iint_S \vec{F} \cdot d\vec{S} = \iint_S 0 dS = 0$$

Horizontal - no partical going across

$$b) \iint_S \vec{F} \cdot d\vec{S} = \iint_S dS = \text{Area} = 2 \cdot 1 = 2$$

Same direction

$$\hat{k} \cdot \hat{k} dS = dS$$

$$c) \iint_S \vec{F} \cdot d\vec{S} = \iint_S xy dS = \iint_S xy dx dy$$

$xy \hat{k} \cdot \hat{k} dS$
 $= xy dS$

the small area is
this


$$= \int_0^2 \int_0^1 dy dx$$

$$= 1$$

ex 3 More w/ symmetries

Compute $\iint_S \vec{F} \cdot d\vec{S}$

a) $\vec{F}(x, y, z) = \langle x, y, z \rangle$




Surface = sphere radius a
at origin



b) $\vec{F}(x, y, z) = \langle 0, 0, z \rangle$
 $S = \text{sphere}$

c) $\vec{F}(x, y, z) = \langle 0, 0, z \rangle$
 $S = \text{upper half of sphere}$

d)



direction of vector $\perp$ surface $= \vec{F}$

$$\iint_S \vec{F} \cdot d\vec{S}$$

$$\overset{||}{|\vec{F}|} |d\vec{S}| = \sqrt{x^2 + y^2 + z^2} ds = a ds$$

$\uparrow$ because $\vec{F} \parallel d\vec{S}$
 $\uparrow$ parallel to

$$\iint_S a ds$$

$$= a \cdot \text{Area}$$

$$= 4\pi a^3$$

b) New field is only part of previous one

Symmetry
of
field

$$\langle x, y, z \rangle = \langle x, 0, 0 \rangle + \langle 0, y, 0 \rangle + \langle 0, 0, z \rangle$$

$$\iint_S \langle x, y, z \rangle d\vec{S} =$$

$$\iint_S \langle x, 0, 0 \rangle d\vec{S} + \iint_S \langle 0, y, 0 \rangle d\vec{S} + \iint_S \langle 0, 0, z \rangle d\vec{S}$$

They should be all =

By symmetry

$$\iint_S \langle x, 0, 0 \rangle d\vec{S} = \iint_S \langle 0, y, 0 \rangle d\vec{S} = \iint_S \langle 0, 0, z \rangle d\vec{S}$$

Hence each = $\frac{1}{3}$ total

$$\iint_S \langle 0, 0, z \rangle d\vec{S} = \frac{1}{3} 4\pi a^3$$

c) Now symmetry of the surface



$$\iint_H \langle 0, 0, z \rangle \cdot d\vec{S}$$

$$= \frac{1}{2} \iint_S \langle 0, 0, z \rangle d\vec{S}$$

✓ disk
Bottom surface of sphere does not count
- problem does not care

$$\iint_H \langle 0, 0, z \rangle d\vec{s} = \frac{1}{2} \iint_S \langle 0, 0, z \rangle d\vec{s}$$